PROBLEM OF PRODUCTION

To the many inquiries which have been made as to the cost of these cabinets and where they may be obtained, no satisfactory answer can be made. Two large concerns have made overtures looking to their manufacture. In order to secure the widest possible educational effect, it is desirable that the exhibits be sold almost at cost to insure their general use. Is it practicable to attain this end? Why should not the Board of Trustees of the Association, the Scientific Exhibit Committee or other regularly constituted representative undertake the supervision of this undertaking? Let them ascertain through experts the lowest figure at which these cabinets, equipped, can be made. Then secure an agreement from some reputable firm to market them at this price.

The Association's endorsement would insure large sale. Let the manufacturer pay the Association a royalty to cover the expense of proper supervision of the data that go into the cabinets. These would need to be varied to suit localities and states. Such supervision would insure the introduction of such matter only as

effective cooperative aid to health authorities and, therefore, would prove of great service in the march of preventive medicine—a field in which the Association, as an organization, has as yet done but little.

311 Newton Claypool Building.

A CASE OF SYPHILIS RESEMBLING PITYRIASIS RUBRA PILARIS

HOWARD FOX, M.D.

NEW YORK CITY

The following case of miliary papular syphilis is of interest on account of its very close resemblance to the papular stage of pityriasis rubra pilaris. When seen in August, 1908, at the Vanderbilt Clinic, the patient, a widow of 30, presented a general eruption of the face, trunk and extremities. This consisted, for the most

would be prudent and sane; sufficiently instructive to the lay mind on tuberculosis and other sanitary questions, but so tempered as not to excite ridicule or unnecessarily alarm the people on such matters. It would offer the Association an opportunity to render most

A case of miliary papular syphilis in which the eruption simulated pityriasis rubra pilaris.

part, of small, conical, pin-head papules, capped with horny scales. There was no tendency for the lesions to become confluent and there were no signs of itching. On the face there were erythematous and scaly lesions that might have occurred with either of the diseases in question. There was no history of any genital sore, no glandular enlargement and no lesions of the mouth. The palms were thickened and scaly, but did not present the appearance characteristic of the early palmar syphilide. While the eruption simulated a pityriasis rubra pilaris very closely, it was noticed that there was a slight tendency of the papules to follicular grouping so characteristic of miliary papular syphilide. Further, the appearance of the backs of the phalanges generally noted in pityriasis rubra pilaris was absent.

When the case was first seen I had not begun to work with the Wassermann test, and a clinical diagnosis alone was possible. To show the great difficulty in diagnosis it may be said that a distinguished dermatologist of the city pronounced the case to be unquestionably one of pityriasis rubra pilaris. The patient was kept under observation for two weeks, at the end of which time the lesions began to show a tendency to become papulosquamous, and she was then put on anti-syphilitic treatment; within a month the greater part of the eruption had disappeared. The patient then became somewhat irregular in attendance at the clinic,

and a few weeks later developed a rather severe iritis. For the privilege of reporting this case I am indebted to Dr. George T. Jackson.
616 Madison Avenue

AN OPERATING-TABLE

D. TOD GILLIAM, M.D.
COLUMBUS, O.

The special features of this table are:

1. An extension rod which slides in and out at the foot of the table.
2. The leg-holders, which fit into the posts and which can be set at any height by means of a set-screw.
3. A revolving instrument tray which is attached to the leg-holder can be raised or lowered at will.
4. The body-lift, plain, or provided with movable side brackets to fix the kidney during operation and to keep the patient from rolling.
5. The shoulder-supports, small, neat and movable. They are hinged at the sides of the table and can be adjusted immediately for use or turned under the table out of the way.
6. A support for rectal treatment or for maintaining the knee-chest posture. This is effected by a strap extending from one leg-holder to the other at a proper height; the patient kneels on the shelf covered by a blanket, his body thrown forward over the strap.

An improved operating-table.

7. The shield between the anesthetist and operator, which fits snugly around the end of the table when not in use, and which can be raised to the vertical or any desired angle and fixed by a set-screw when needed.

The extension rod at the end of the table has many uses:

1. In vaginal or rectal operation a towel thrown over it protects the sutures from contact with the drop.
2. Drawn out a little further it makes a foot-rest for the patient in examination or treatment.
3. It holds a douche-pan in vaginal or rectal operations.
4. Pulled out so as to allow the legs to be lowered through it and strapped to the drop, the same douche-pan, placed between the end of the extension rod and the patient's knees, acts as a receptacle for soiled sponges in abdominal operations.
5. By placing the shelf from under the table on it and adjusting the extension rod to the length of the patient, a full-length table is obtained exactly fitted to his stature.

Every attachment is an integral part of the table and always at hand, there being no pieces to lay aside and gather up when needed. Also there are no hinged leaves or other unstable parts to the table.

50 North Fourth Street.

A PROTECTOR FOR USE IN THE CUTANEOUS TEST FOR TUBERCULOSIS

R. L. SUTTON, M.D.
KANSAS CITY, MO.

In applying the cutaneous test for tuberculosis, the scarified surfaces, unless covered, are subjected to irritation and possible infection from without.

Protection usually is secured by the use of a soft, smooth dressing, as a linen or silk handkerchief, applied as a bandage directly over the wound. While this method serves very well in many instances, it is open to some objections. The cloth comes closely in contact

Protector for use in the cutaneous tuberculosis test.

with the excoriated surfaces and is liable to irritate, causing misleading inflammation; the dressing is difficult to keep in place, especially on the forearm (the part commonly selected); and the progress of the reaction can not be observed without removing the bandage.

I have found an oblong felt protector, like the one shown in the accompanying illustration, of considerable value. It is made on the same principle as the widely used vaccination pads, and the openings are covered with thin, transparent celluloid. The dressing is firmly held by four short, transverse strips of adhesive plaster.

518 Commerce Building.

Special Article

SEVENTH ANNUAL SUMMARY OF FOURTH OF JULY INJURIES

For the seventh consecutive year we are presenting statistics of injuries received during the celebration of the Fourth of July, with particular reference to tetanus resulting from these injuries. Considerable effort has been made to secure reliable data, and all serious cases have been carefully investigated, so that, so far as the figures go, dependence may be placed on them. We have received thousands of letters from physicians in all parts of the country, reporting cases which otherwise could not have been included in the statistics. There are doubtless thousands of other cases not reported, although chiefly minor injuries. Our thanks are due for the many careful reports which have made these statistics more complete and the figures more reliable. The data are presented in the same manner as heretofore, in order that comparison may be made.

MORE CASES OF TETANUS

There were 150 tetanus cases this year, almost double the record of last year, when 76 cases were reported. This is the largest number since 1903, when there were 415 cases. It is significant to note that the number of blank cartridge wounds correspondingly increased from 816 last year to 1,095 this year, and that the states having the largest numbers of blank cartridge injuries have