of a more fibrous nature than in children, having lost to a degree the fish-worm character and being more in furrows and folds which run in an anterior and posterior direction. Somehow the real size is deceptive and there is always more of the tissue than would appear to be from examination.

These patients are apt to be dismissed in clinics with sprays and gargles, and in private practice with sprays and atomizers and other forms of what should be called "office suggestion." After a course of such treatment, patience ceases to be a virtue and the sufferer prefers to continue his unpleasant existence as it is rather than submit to the experiments which experience has taught him are of no avail.

In an average practice there should be four or five well-defined cases in a year.

The case from which this photograph was taken is one of the most instructive ones that I have recorded and it is presented as illustrating a type and also as a lesson in humility, as the underlying pathologic condition was overlooked by several good general physicians and by at least one specialist of reputation. When recognized it was not relieved by the first operation and for no other reason than that it seemed too simple for the exercise of

Second adenoid growth removed. The one removed the day before was fully as large.

careful technic. The accompanying illustration shows the real size of the specimen, with the old-fashioned curette which I have now discarded.

Patient.—E., female, aged 38, single, well developed; home in Nebraska, temporary climatic resident of Colorado. (When an eastern physician exhausts his resources the patient is sent to Colorado for climate regardless of what the disease is.) She said that she had a dry mouth, slight deafness, frequent desire to clear the throat, which was not relieved by the almost constant hawking; ears hurt when nose was blown hard. At one time a douche was prescribed which brought on an attack of excruciating pain in ears which have been worse ever since.

Examination.—This showed the presence of adenoids, "granular pharyngitis," hypertrophied inferior turbinate on one side and the throat full of ropy mucus.

Treatment.—Only enough of the turbinate was removed to give clear space and the adenoids were supposed to be thoroughly curetted. The next day the voice was still muffled and examination showed the mass in the nasopharynx apparently as large as before although the quantity removed the day previously was fully as large as the specimen shown in the illustration. Local anesthesia was used as before and the mass removed under the inspection with the mirror by one sweep of the curette.

Remarks.—Evidently the tumor was bifurcated and the anterior portion was pushed forward into the posterior nares at the first operation. At the completion the patient at once remarked: "Doctor, you have removed something that has been stopping up my nose for years." She remained three weeks for treatment which consisted of massaging a strong solution of one of the silver salts into the nasopharyngeal membranes with use of the Eustachian catheter every third day.

Before operation the tuning-fork test was minus 30 and at the time of returning home three weeks later it was minus 15, both ears being about the same. A letter received two months later stated that the throat had given no trouble and that the mouth and ears were well.

Adenoids in adults are not common, as they usually disappear before maturity, but when they do not their character changes from those of childhood.

The patients are really sufferers and think that there is no relief because the "catarrh is in the system." There is no value in sprays and applications or internal treatments. Results of complete operation are satisfactory and permanent.

SUPPURATIVE ACCESSORY SINUS DISEASES IN WHICH AN INTRANASAL OPERATION IS INDICATED *

HANAU W. LOEB, A.M., M.D.

Professor of Diseases of the Ear, Nose and Throat in St. Louis University
ST. LOUIS

The great majority of the cases of suppuration of the accessory sinuses of the nose are amenable to intranasal treatment at some period in the course of the disease. Suppuration of the ethmoid labyrinth responds most readily to this form of treatment, with that of the maxillary, sphenoid and frontal, following in the order mentioned. Intranasal treatment may be of no avail in suppuration of any of these cavities, however, if the process has been long-continued, or if the infection is rapid and virulent. Between these two extremes lies the field of intranasal surgery as applied to the suppurative conditions which involve the accessory sinuses of the nose.

The methods by which good results are obtained are various, and for their success depend largely on the personality and equipment of the individual operator; for a method which is satisfactory in the hands of one thoroughly familiar with the technic may be without favorable result in the hands of another who is accustomed to a different plan of operative procedure. Inasmuch as this paper comprehends conditions rather than individual methods, only casual reference will be made to the latter where the discussion requires it.

Besides discussing processes which call for intranasal lessons which anatomy teaches with respect to surgical treatment, however, it will be the endeavor to show the treatment of the accessory sinus through the nose.

To bring this more clearly into view, there are here presented lateral reconstructions of twenty heads, showing forty groups of sinuses with their relations shown to one another and to the floor of the nose, the turbinates, and the external contour of the nose and upper lip. The frontal, sphenoid and maxillary sinuses, and the contour of the nose and upper lip are shown in unbroken lines, the posterior ethmoid in dotted lines, the anterior ethmoid in alternate dots and dashes, the infundibulum in broken lines, and the turbinates in feathered lines.

* Read in the Section on Laryngology and Otology of the American Medical Association, at the Sixtieth Annual Session, held at Atlantic City, June, 1909.

This makes it possible to see the exact relations of each factor that is brought up for consideration. The nasal orifices of the sphenoid and frontal sinuses are indicated in black dots. The method employed in reconstruction has already been fully described in previous papers on the subject.

MAXILLARY SINUS

An intranasal operation ordinarily suffices for the relief of simple, uncomplicated suppuration of the maxillary sinus, provided there is no dental cause. Even then, relief may follow the intranasal operation if there is no active process going on in the teeth or alveolus. Furthermore, good results may be obtained by combining with the nasal operation the removal of diseased roots of teeth.

While a simple puncture through the nasointral wall is often sufficient, as a rule the best results are obtained only when an extensive resection of this wall is made. This naturally requires the removal of a portion of the inferior turbinate, in this way affording ample space for irrigation and drainage. The well-known tendency of all these openings to contract, and, in many instances, to close altogether, further calls for a large opening which would, at least, do no harm.

Considering the variability in the size and shape of the maxillary sinus, it is always important to be sure that the opening is made into the sinus and not into the tissues anterior or inferior to it. If the opening is made too near the anterior extremity of the inferior turbinate, it may escape the sinus altogether; as, for instance, in heads II left, III right, IV right, VII, VIII left, IX, X right, XI left, XII, XV, XVI, XVII, XVIII, XIX, and XX. In these heads the distance between the anterior extremity of the inferior turbinate and that of the maxillary sinus is more than 5 mm. This distance is greatest in VII right, IX, XV left, XVI left, XVII, XVIII, and XX.

In IX the maxillary sinuses are so small that only a careful operator who has anticipated the existing state of affairs would have been successful at the first attempt.

In the main, the inferior turbinate courses along the nasointral wall in a horizontal direction, but in a number of instances it inclines sharply downward and backward from its anterior attachment. This is particularly well shown in X right, XIV, XVIII and XX, in which it reaches the level of the superior wall of the sinus, or even higher. In only one instance, IX, does the inferior border of the inferior turbinate reach a lower level than the sinus, though in several cases the distance between them is not great: VI left, XII, XVI, and XVII right.

The position of election for opening the sinus, as ascertained by a study of these twenty heads, is just below the attachment of the middle portion of the inferior turbinate. This enters the sinus in every instance, and, in the main, at its most dependent part—at least, so far as possible, by using the inferior turbinate as a guide.

When the inferior turbinate has the great inclination already mentioned and the floor of the nose is widely separated from the turbinate attachment the portion of the nasointral wall penetrated under this rule will be high above the floor of the sinus, so that it will be advisable to remove the wall below this for quite a considerable distance, easily determinable at the time of operation.

FRONTAL SINUS

In some acute cases, even when associated with ethmoiditis, operation through the nose will afford sufficient drainage and thereby result in cure. Even in the more chronic cases relief follows resection of the adjacent ethmoid cells and irrigation through the nasofrontal canal with or without enlargement. Although claims are made that well-marked cases of frontal sinus suppuration may be relieved by the various intranasal methods, it must be admitted that ordinarily nothing short of an external radical operation suffices.

A study of the reconstructions here presented throws some light on the accessibility of the frontal sinus to nasal operative procedures, and, at the same time, on the anatomic variations which make the operation uncertain, difficult or impossible. These comprehend the distance between the nasal opening of the frontal sinus and the anterior wall of the nose, the presence of an ethmoid cell or nasofrontal canal which reaches a high elevation, and, what can not be shown in these illustrations, septal irregularities, congenital narrowness, or new growths.

XXI

Frontal sinus (sf), anterior ethmoid cell (cea) and posterior ethmoid cell (cep), extending over the roof of the orbit; dnf, nasofrontal duct; lc, cribriform plate; nos, nod, optic nerves; c/s, left internal carotid.

In the following a probe would readily enter the frontal opening without any likelihood of entering any other cell with which the frontal might be confused; of course, assuming that the operator understands the technique: I right, III right, IV, V, VI right, VII, VIII, X, XII, XIII right, XVI, XVII, XVIII, XIX right, and XX.

In only twenty-five out of forty instances, therefore, could the probing be accomplished with a reasonable degree of accuracy, and in several of these cases one who is not an expert might be deceived by the entrance of the probe into one of the adjacent ethmoid cells or nasofrontal canal. This can be more or less obviated by making a radiograph which would show how high upward the frontal extends.

In I left, II left, IX, XI, XIII left, XIV, XV, XIX left, an adjacent ethmoid cell reaches so nearly the superior level of the frontal that it would be practically impossible to state whether the probe was in it or

In these figures, the frontal, sphenoid, maxillary sinuses and the contour of the nose and upper lip are indicated by unbroken lines, the posterior ethmoid by dotted lines, the anterior ethmoid by alternate dots and dashes, the infundibulum by broken lines and the turbinates by feathered lines.

in the frontal, and in II right and VI left the naso-frontal canal occupies a similar relation. It is thus evident that there must be some doubt in almost half the cases as to whether the frontal sinus is being treated when intranasal operative measures are directed toward relief of frontal sinus suppuration. It must be admitted, however, that a careful study of properly taken radiographs will reduce this possibility of error considerably, but this involves lateral as well as anterior views, perhaps with probe *in situ*.

The intranasal operation has, therefore, a limited field of service, and is subject to much likelihood of error. There is, however, this saving clause—that, even if the operation is directed toward an adjacent cell, so much bone and other tissues will be removed that drain-

age from the frontal is promoted and the condition is relieved.

We do not anticipate a good result in those cases in which the frontal extends over the orbit and is affected in this portion.

ETHMOID CELLS

Ethmoid suppuration, when uncomplicated, practically always responds to intranasal treatment, for the entire ethmoid labyrinth may be exenterated by operation through the nose. The most anterior cells can not, sometimes, be freely opened and curetted on account of their nearness to the nasal bone. Hajek asserts, however, that it can always be done if the middle turbinate has been properly resected. There are occasionally

anomalies of development of the ethmoid cells, which render the operation either difficult or unavailing. For instance, in XXI, it will be seen how both the anterior and the posterior ethmoidal cells extend, in this particular case, over the roof of the orbit. In this position the cells could not be properly exenterated through the nose and would probably escape observation in a radical frontal operation.

Of course, there is always some danger in working in the dark, as it were, and the process may be most marked in the most inaccessible portion of the cells, which extend farthest from the seat of operation. Yet,

it must be said that, as in all suppurative conditions, drainage does a great deal toward affording relief, and the operative field being in the most dependent portion of the cells, drainage is apt to be secured.

The truth of Hajek's contention—that if the middle turbinate is resected the most anterior of the ethmoid cells may be exenterated—is borne out by the figures. In only nine instances—I, III left, VII, VIII left, XIV left, XV left, XVIII right—does an ethmoid cell extend as far forward as the anterior extremity of the middle turbinate; and in most of these instances the distance is insignificant, certainly not sufficient to inter-

ferred with the operation. The cells extend upward considerably in most of the heads, more than 10 mm. above the superior attachment of the turbinate in I left, II, III, V, VI, VII, IX, XI, XIII, XIV, XV, XVI, XIX and XX. This distance is very considerable in I left, II, III left, VI, XIII left, XIV, XV left, XVI and XX. In I left an ethmoid cell extends upward anterior to the frontal for a distance of 17 mm., while in IX right an ethmoid lying immediately behind the frontal reaches a higher level than the frontal.

These facts are important only in that they show how high up the ethmoid cells extend, thereby accounting for the recurrence of the polypi even after the floor of the cells has been removed; they emphasize the im-

portance of thorough exenteration of the diseased part. It goes without saying that, however radical one may be in this operation, respect must be had for the cranial wall of the sinus.

SPHENOID SINUS

In uncomplicated cases, and those in which the ethmoid alone is involved with the sphenoid, the nasal route is the only one to be advocated. There is a growing disposition, even among those who formerly thought that the external operation was always to be advised, to rely more on the intranasal method.

The nasal operation can be of service only by the curettement and the improved drainage occasioned by the enlargement of the nasal opening. Whether this be

confined to the pars nasalis, or, as Hajek suggested, made to include the pars ethmoidalis, the endeavor is the same.

The location of the nasal opening of the sphenoid is such that in any suppurative process affecting it there is a quantity of pus more or less constantly present. Practically all the sphenoids here shown have the opening so far from the floor that this must be quite general. In most comparison will show that even extensive resection of the anterior wall will fail to reach the floor. The drainage which results from this operation, however, provided the opening be permanent, will generally be followed by disappearance of the pus.

The distances between the sphenoid orifice and the tip of the nose and the anterior border of the middle turbinate are given in the table:

No.	DISTANCE FROM SPHENOID ORIFICE		Anterior Border of Middle Turbinate.	
	Anterior Tip of Nose.		Right.	Left.
I	77	77	36	31
II	77	74	43	38
III	64	69	33	30
IV	75	72	35	33
V	72	74	30	31
VI	75	80	37	44
VII	88	93	37	39
VIII	69	72	29	38
IX	70	71	37	32
X	72	70	38	33
XI	88	89	37	33
XII	91	88	41	34
XIII	77	77	38	35
XIV	84	82	39	34
XV	86	79	41	31
XVI	71	71	30	35
XVII	77	77	41	39
XVIII	80	80	50	40
XIX	77	77	42	36
XX	87	92	37	40

For the first, the variations are from 64 to 93, the usual (leaving out the five shortest and longest) from 71 to 88, the average 78.

In regard to the distance between the orifice and the anterior border of the middle turbinate, the variations are from 29 to 50, the usual (leaving out the shortest and longest five) from 31 to 41, the average 37. This gives a fair idea of the length of the probe and operative instrument required.

In a previous paper I called attention to the greater likelihood of optic nerve involvement, in cases in which the nerve lies near the level of the orifice, for this would signify that the pus would be separated from the optic nerve by a thin bony wall only, instead of the air, which would intervene if the opening was at a lower level. Whether this relation can be ascertained by radiographic study is still to be determined, with fair prospects of success. At any rate, sphenoid suppuration with only the slightest suspicious eye symptoms calls for immediate operation.

I wish, again, to call attention to an observation made before this body two years ago, with reference to the four heads reconstructed up to that time, that in cases in which the middle turbinate has an inclination downward and backward the floor of the sphenoid lies at a lower level than the ethmoid, or at its level. Compare I, II, V, VI, X, XI, XII left, XIV, XV left, XVII, XVIII left, XIX and XX. It will be noted that this is confirmed in every instance, except XV right, in which the sphenoid is only a little pit 2 by 4 by 2 mm.—and the left sphenoid, which occupies the space usually given over to the right, follows the rule. A glance at XV, right, shows that, even though the sphenoid be small, it lies considerably nearer the level of the floor than that of the roof of the ethmoid.

In the other instances in which the middle turbinates run horizontally backward the floor of the sphenoid lies above that of the ethmoid. The exact value which this will have will depend on the further confirmation of the observation and on the subsequent development of its utility.

Humboldt Building.

Ligation of the Jugular in Sinus Phlebitis.—R. Botey of Barcelona says that he will present a communication on this subject at the International Medical Congress, and the *Archives Internat. de Laryngologie*, August, has, rather prematurely, published it. He states that he has encountered 8 cases of thrombophlebitis of the lateral sinus in the course of 404 mastoid operations, and that the 4 patients who survived the phlebitis had not had the jugular ligated, while this was done in the 4 other cases. He used to ligate as a matter of course, but his own experience and review of the literature have convinced him that in most cases ligation is useless at best, and is liable to do actual harm. He advocates curetting the sinus in case of otitic thrombosis, without the Gerhardt sign, curetting from the knee to the bulb, without ligating the jugular. When the jugular is manifestly affected, and can be felt as a hard cord, it should be opened, ligated over the clavicle and resected. Pulsation in the sinus is not a sign of phlebitis, he adds, but is merely a transmitted pulsation from the brain tissue.