uncomplicated typhoid fever is to be recollected. At all events, in view of the foregoing facts, the use of vaccines in the treatment of this disease should be somewhat restricted according to conditions and indications presented by individual cases. Their routine use, however, is not to be encouraged, but it is obvious that an anti-endotoxin serum would, by neutralizing the soluble endotoxins, constitute the ideal method of treating typhoid fever. Such a serum has been prepared by MacFadyen, and favorable results have already been reported from its use, when administered early in typhoid cases. It is not improbable that the vaccine treatment of typhoid fever, except as indicated below, will give place to the method of establishing passive immunity by means of an anti-endotoxin. In view of the marked discrepancy among clinical observers as to the size and repetition of the doses, there would seem to be great necessity for an especial consideration of these aspects of the question of therapeutic inoculations in the future.

Vaccine therapy in the treatment of typhoid fever, however, should receive a more extended trial than hitherto, more particularly in the earlier stages of mild types of the disease, before being rejected. Finally, in the present state of our knowledge, the value of vaccines for the following purposes must be conceded: (1) as a means of prophylaxis; (2) in suitable cases when continued during convalescence, to prevent relapses; (3) to combat local infections with the typhoid bacillus, as for example, bone suppurations which arise in the period of convalescence, and (4) for the removal of the typhoid bacilli from the feces and urine in the case of typhoid-carriers.¹⁰

I desire to express my thanks to Dr. A. A. Stevens, who at my suggestion, treated three of the cases reported above with the results of blood examinations before and after the inoculations, in the wards of the Protestant Episcopal Hospital of Philadelphia, and to Dr. Robert B. Walker for furnishing one of the cases belonging to my series from the Philadelphia Hospital.

1005 Walnut Street.

THE FUNDAMENTAL PRINCIPLES UNDERLYING THE TREATMENT OF WOUNDS ON THE BODY SURFACE

CHAUNCEY E. TENNANT, M.D.
DENVER

Steadily through the maze of a multiplicity of methods, there has crystallized the present treatment of surface traumatic lesions. Based on the knowledge of definite pathologic and histologic processes and confirmed by the shorter periods of disability and lessened distortion of function, modern treatment of surface injuries may be reduced to two fundamental principles: *moist hyperemia* and the *open-air treatment*; in other words, the use of a dressing capable of giving sustained heat and moisture, or one so constructed that it will not displace the protective covering of coagulated serum, yet will protect the wound from further damage.

The wide latitude of pathologic possibilities in surface lesions, varying as they do from a microscopic skin puncture attended with violent constitutional toxemia and death, on the one hand, to extensive abrasions and lacerations with no constitutional manifestations on the

other, has, in a measure, justified any course which has had as its result a satisfactory functional termination.

Until recently, little thought was given to the individual factor of lowered bacterial resistance, or the influence of the general state of nutrition on cell-regeneration. But now, with the more careful and detailed consideration of the possibilities of lowered resistance and cell nutrition, there has been a stimulus for the introduction of methods calculated either to intensify locally, or to increase, the number of antibactericidal bodies at the various parts of the organism where needed, or to engorge the tissues with nutritive plasma for the revitalization of the traumatized cell.

Surface lesions of the body widely differ in character, as burns, incised or crushing wounds, etc., and they may also vary from the small pin-point puncture to the extensive loss of continuity which is limited only by the size and force of the destructive agent. But so far as

Fig. 1.—Method of applying large and copious wet dressings.

the end-results are concerned, the only difference is their type and degree of infection. Therefore the plan of treatment must be early outlined, if the most successful results are to be obtained. It is well known that in burns, skin-grafts and other superficial traumas which remain sterile, the dry method of treatment gives the most satisfactory results both as to convalescence and function, while the moist hyperemia is the treatment especially adapted to infected wounds, where the organism has lodged itself in the subcutaneous tissues and lymphatics.

As a rule, the infection is either of the staphylococcic or streptococcic type; the former invariably localizes in a small area, while the latter is most likely to lodge in the subcutaneous tissues and extend rapidly along the lymphatics. Where resistance is fair and early excited, the chemotactic disturbance is quickly manifested with temperature and other clinical phenomena of inflammation. In either type of infection, it is extremely impor-

¹⁰. Irwin, S. T., and Houston, T.: *Lancet*, London, Jan. 30, 1909, p. 311.

tant, as soon as it becomes evident that a pathogenic organism has lodged in the subcutaneous tissues and that the resistance force is low, that these natural forces be augmented by either the active or passive hyperemic process, or both. Every effort must be immediately made to create an overwhelming quantity of polymorphonuclear neutrophils and leukocytes to inhibit and digest the invading organisms before they have time to extend into the surrounding structures.

In all injuries involving a break in the skin, the first question to be determined is the presence or absence of infection, and if there is infection, whether or not the individual has sufficient resistance to destroy the organisms quickly providing moist hyperemia is properly used.

Fig. 2.

Fig. 3.

Figs. 2 and 3.—Sheet of light aluminum to which is fastened a strip of poultry wire.

The time is not long past when all these wounds were considered as primarily infected, and vigorous steps were immediately taken to rid the host of the unwelcome guest. Active antiseptics were freely used, most of which were destructive alike to tissue and invading organism, and, as a result, tissue necrosis with moist surfaces, sloughs, granulations and large cicatrices was the common sequence.

At the present time, the practice of putting every injury in wet dressings, or even the use of copious washings, is uncalled for; especially as the coagulated serum is extremely soluble in water, and such procedure means the removal of Nature's protective layer of coagulation from the surface, and a portal left open for infection.

The care of these cases, requires of course, considerable judgment, but why modern surgery should find any more reason for vigorous and persistent cleansing of these surface lesions than in the case of gunshot wounds of the extremity seems a question worthy of consideration.

If it is good practice to put gunshot wounds up in sterile dressings and wait for further evidence of infection, why should not the same principle apply to other reasonably sterile and easily accessible traumas of the surface? Where the history of the case shows a strong probability of there having been infectious material introduced into the wound, then great care should be used and proper application of the wet dressings made, with full knowledge of the fact that Nature's protective barrier must be sacrificed in the use of moisture.

We know that wherever infections occur, the use of moist dressings properly applied, quickly raises the polymorphonuclear neutrophilic blood content, which

Fig. 4.—Copper or aluminum coated fly-screen bound with adhesive tape.

Fig. 5.—Adhesive straps are then passed over the screen, holding it firmly to the adhesive sheeting.

soon overwhelms the invading organism. Where no infection exists, the dry dressing, properly applied permits of surface coagulation of the exudates, which forms both a splint and an impervious barrier to further infection. Repair is very prompt in these latter cases if destructive antiseptics have not been used. Here the moderate application of sterile warm water and a bland soap, followed by open-air dressings, is usually all that is necessary to accomplish rapid and uncomplicated repair. Even in the greasy dirt-begrimed hands of the machinist, where it is not possible to wash the wound, the application of either turpentine or the tincture of iodine is very practical and effective, and may be safely followed by the dry dressing.

The day of the local applications of salves and dressing powders in emergency surgery, fortunately, has passed. If the indications for the treatment of these cases are properly interpreted, and the principle underlying the use of moist hyperemia or open air is fully

comprehended and applied, the profession at large will greatly reduce the number of general septicemias. Where infection is definitely present, free incision must of course be made, and followed by the application of very large and copious wet dressings, consisting of, first, the gauze covering the site of the lesion and near-by tissues, then several layers of cotton, and finally the oiled silk, moisture and bandage (Fig. 1).

If a finger only is involved, the entire hand should be included in the dressing, so that the full benefit of the hyperemia may be secured. When the oiled silk has been well fastened down, a last small opening should be left, through which the hot fluid is poured until the dressings are supersaturated. After this, the oiled silk should be completely closed, to retain both moisture and heat. A splint should then be applied to keep the part at rest. Extreme care must be exercised that the solutions poured into the dressings are not too hot, for extensive burns may easily be produced, which are often more serious to care for than the original lesion.

Fig. 6.—A very satisfactory dressing in skin grafting and varicose ulcers.

Several solutions may be used to moisten these dressings, the simplest and safest of which is the hot sterile physiologic salt solution. Equal parts of alcohol and saturated solution of boric acid make a solution which is ideal for use in early dressings, because of its local stimulating effect on the tissues, which produces an active local hyperemia. Thiersch's solution (salicylic acid, one-half dram, boric acid, 2 drams and a quart of sterile water), so well recommended by the late Nicholas Senn, is another safe and effective solution for use in the wet dressings.

Unless the dressings are large enough to retain both moisture and heat for long periods of time, they are worse than useless.

Dressings should be changed once in twelve to twenty-four hours, depending on the amount of discharge, but the solution should be repeated as often as necessary. A change of the solution is often of great advantage, especially if the alcohol has first been used, since the other named solutions are less stimulating yet almost as satisfactory as hyperemic agents. After a certain period tissues often refuse to respond to further stimulation.

Where infection has not occurred, and is not likely to occur, it is much better that no dressings should come directly in contact with the wound, especially if it is extensive and moist. It has long been recognized that burns and traumas of the face, where dressings are not applied, soon heal with but little inconvenience to the patient, and a minimum of cicatrix. The same principle applied to other parts of the body works equally well. However, to devise successful methods for the application of treatment to these parts has been a difficult task, and the means at present used are almost as many as the surgeons applying them.

Two methods which I have been using for the last four years may be of interest, since they have been extremely satisfactory in my work (Figs. 2 and 3).

One is the use of a sheet of light aluminum as a base on which is fastened a cage made of poultry wire of any desired mesh for enclosing an extremity. The aluminum is light, clean, and easily punctured, yet firm enough to hold the poultry-wire top to the desired position. This frame is extremely light and comfortable, and easily applied. It is especially useful in extensive burns of the extremities.

The other is the use of aluminum-coated metal fly-screen, cut to the desired size, bound with adhesive tape, and set on strips of adhesive felt sheeting, which have first been cut, fashioned and attached to the skin about the outer margin of the wound (Figs. 4 and 5).

Adhesive straps are then passed over the screen, holding everything firmly to the part, yet nothing touches the wound (Fig. 6).

It is then covered with gauze dressings and snugly bandaged.

This last method has, in my experience, been very satisfactory in skin grafting, and in varicose ulcers of the leg. In the latter lesion, one may apply any form of medicament first, then cover with wire netting as described.

612 Empire Building.

SOME EFFECTS OF BRIGHT LIGHT ON THE EYES *

J. HERBERT PARSONS, F.R.C.S.

Surgeon, Royal London Ophthalmic Hospital; Assistant Ophthalmic Surgeon to University College Hospital

LONDON, ENGLAND

It is my first and most pleasing duty to express to you my high appreciation of the very great honor you have done me in inviting me to deliver this address to so distinguished an audience. It has been my good fortune to have met many of you on the occasions of your pilgrimages to the Old World. Moorfields and Vienna have long been the Mecca and Medina of ophthalmologists, and we at Moorfields are proud of the sacred fire which has been handed down to us by our predecessors. No branch of science, however, can flourish on the records of the past, and we in our turn must make pilgrimages to newer shrines, there to gather fuel to keep the sacred flame alight. I would fain believe that you in the past have learnt much from us; we too have learnt much from you. Thus in the brotherhood of science may we cement the brotherhood of race which is our inalienable heritage.

* Read in the Section on Ophthalmology of the American Medical Association, at the Sixty-First Annual Session, held at St. Louis, June, 1910.