this work will be of considerable help to the general operating surgeon.

DR. YANDELL HENDERSON, New Haven, Conn.: The work of Carrel at the Rockefeller Institute has shown a great difference between the effects of interference with the circulation on the side of the vein and on the side of the artery. It seems to me that in such experiments as those of Drs. Eisen-drath and Strauss, care should be taken to note whether both artery and vein were shut off, or whether partial compression was used, shutting off the vein entirely and the artery partially. If the vein were compressed completely and the artery only incompletely, the outflow would be shut off and the renal blood-vessels (both arteries and veins) would be distended by the full arterial pressure. The work of Carrel shows that many organs stand interference with their arterial supply fairly well, but are greatly harmed by interference with their venous drainage.

DR. H. GIDEON WELLS, Chicago: Calcification of the tubules is usually very extensive in experiments with the kidneys of rabbits, and one has to consider the fact that the blood of these animals contains more calcium than does that of either the carnivora or the omnivora. The results of such experiments have been applied to the problems of general calcification. This, I believe, is entirely and fundamentally erroneous, for the reason that in the case of the calcification of the necrosed tubules of the rabbit's kidney there is the same process as in calcification of dead organic material in the urine, in which there is a solution of certain organic and inorganic salts, differing from calcification occurring within the body, when the calcium comes from the circulating blood.

A point brought out with regard to Litten's work is that he found no definite relation between the functional changes in these cells and the anatomic alterations. We are too little inclined to bear in mind the entire discrepancy between functional and histologic changes. Although this lack of correspondence of structure and function has been frequently mentioned, yet one so frequently sees it disregarded that I do not hesitate to emphasize this point. The opposite condition also holds good, namely, that extreme anatomic alterations are not necessarily productive of defective functional capacity. In Washington, in May, I presented before the American Association of Pathologists and Bacteriologists a paper describing the effects of the most extensive possible fatty degeneration that I had been able to produce in the liver. I found that the liver-cells in these cases showed as great a power to destroy uric acid, which is difficult to oxidize, as do the cells in the normal liver. Therefore, cells may show extensive histologic changes and yet be functionally competent; and may also show no histologic changes and be totally incompetent.

DR. DAVID C. STRAUSS, Chicago: In the beginning of the paper I did not care to refer to the literature, because there is so much of it, and most of it does not apply to temporary compression, but to permanent ligation. In regard to the functional effect of temporary compression of the renal vessels, I found mention of some experimental work done by Ludwig on dogs. He found that temporary compression of the renal pedicle often caused temporary anuria and albuminuria that remained for hours, or even days; but this did not always follow. In our paper we simply wanted to refer to the anatomic changes produced. As Dr. Wells has pointed out, the functional changes ought also to be studied; however, this is simply a preliminary study. With reference to whether in our experiments there was complete compression of the vein but not of the artery, I would say that the clamp used by us was very strong and was protected by rubber, so that there was little chance of failure to compress the artery entirely. Moreover, in every case I felt the artery in order to be sure that there was no pulsation between the clamp and the kidney, and in some cases put on a second clamp to be more certain. In regard to Dr. Wells' remarks concerning calcification, I did not quite understand the point he made about the deposition of calcium being the same as in the urinary bladder.

DR. H. GIDEON WELLS, Chicago: There is simply a precipitation of calcium as from a sponge. It is not comparable to the calcification of the bronchial glands, for instance.

DR. DAVID C. STRAUSS, Chicago: Litten, in his article, points out that the chief reason why there is so much more calcification in these temporary compression experiments than in experiments in which permanent ligation is done, is the fact that the blood-vessels are not destroyed by the brief temporary compression, and that the circulation through the renal vessels is reestablished. This allows further deposition of calcium to occur; whereas, when the vessels are ligated permanently, the only source of calcium deposit is from the collateral circulation. We did not observe functional changes. They ought, however, to be observed.

THE ETIOLOGY OF EPITHELIOMA *

ARTHUR E. HERTZLER, M.D.
ROSEDALE, KAN.

I. THE DEVELOPMENT OF FIBROUS TISSUE

The result of some studies in the development of fibrous tissue as observed in wound-healing form a necessary introduction to the consideration of the behavior of connective tissue in its relation to developing epithelium. It will be necessary, therefore, to present in outline some researches already published.¹

The first change in the healing of any wound is the exudation of a colorless serum. Within a few minutes this material begins to coagulate, forming bands extending from one wound surface to another. These bands become fully formed after a few hours. They take the direction most effectual for union. I speak of this process as coagulation because the fibers so formed stain by Weigert's fibrin stain and stain red by Mallory's stain, as is characteristic for fibrin (and Mallory's fibroglia fibrils), and because those things which prevent the coagulation of blood prevent the formation of these fibrils. We have then a tinctorial and a physiologic test in agreement that the process here observed is identical with that of blood coagulation. These fibrils after a few days lose their affinity for the Weigert and the red in Mallory's stain and stain blue with the latter and are, therefore, fibrous tissue. It remains to determine that the fibers staining blue with Mallory's stain after three to six days are the same as those which stain red with Mallory's and blue with Weigert's stain during the first few days.

Two facts are available for this proof. In a large series of experiments in which corn-pith was stitched to a peritoneal surface the fibrin bundles formed over the surface geometrically ("in a mathematical manner"—Kulmann). These results are constant.

After the change in tinctorial reaction has occurred, the fibers have an identical arrangement. If, therefore, the fibrin fibers are replaced by fibrous tissue the arrangement is identical. That they are not replaced finds its chief argument, however, in that a part of a single fiber may stain blue while the remainder still takes the red stain. These results also are constant. From these facts I conclude that the initial process in wound healing is the formation of fibrin identical with that formed in the coagulation of blood, and that the fibrin fibrils are converted into fibrous tissue fibrils.

* Read in the Section on Pathology and Physiology of the American Medical Association, at the Sixty-first Annual Session, held at St. Louis, June, 1910.

1. Hertzler, A. E.: Pseudoperitoneum, Varicosity of the Peritoneum and Sclerosis of the Mesentery, *THE JOURNAL A. M. A.*, Jan. 29, 1910, p. 351.

II. CHANGES IN CONNECTIVE TISSUE IN ARTIFICIALLY INDUCED EPITHELIAL CELL PROLIFERATION

When Sudan III in olive oil is injected beneath the epithelium, epithelial cell proliferation takes place to such an extent as to simulate very closely a beginning epithelioma. The influence of the Sudan oil, according to Fischer, is to act as an attractin. No attention has been paid to the fact that the connective tissue in the region of the developing epithelium loses its specific tinctorial reaction to connective tissue stains and that epithelium does not proliferate in regions where this change in the connective tissue has not taken place. From study of sections of specimens obtained soon after the oil has been injected and for short intervals afterward, it seems that the connective tissue changes precede the epithelial proliferation. The epithelial proliferation, too, takes place after the use of those dyes only which induce these changes in the connective tissue, namely those of the OH group. The changes in the connective tissue are the reverse of those changes which take place when fibrin changes to fibers, that is to say, the changes are from blue to red instead of red to blue. I am not certain that I have succeeded in completing the reverse process. Fibers occur which stain red by Mallory's stain, but I have not succeeded in reversing the process far enough to get a positive reaction with Weigert's fibrin stain. As a corollary to this line of evidence it may be mentioned that epithelium does not develop in wound-healing in the presence of adult connective tissue. The use of Sudan in the treatment of chronic ulcers has been supposed to act by stimulating the epithelium to proliferation. The connective tissue changes produced by the Sudan when used as an ointment are identical with those induced when the dye is injected. I conclude from such studies that the changes in the connective tissue are probably the essential factor in the production of the epithelial proliferation. When the healing of chronic ulcers is attempted by surgical means we empirically bring about the same condition which we produce by the Sudan, that is, fibrin is supplied, or in other words, we cause the chemical reaction of fibrous tissue to approach that of fibrin.

III. THE CONNECTIVE TISSUE CHANGES IN EPITHELIOMA

In advance of epithelial proliferation in beginning cancer the connective tissue undergoes changes similar to those artificially produced in the Sudan experiments. It no longer stains in a characteristic manner by Van Gieson and stains a paler blue by Mallory's stain. The reaction to Van Gieson is identical with that given by the connective tissue in the Sudan experiments. The results are so constant as to be most valuable in the determination of the clinical character of epithelial cell proliferations. The chief matter of interest, however, is the occurrence about certain cell-nests of small bundles of fibrils which correspond to my fibrin fibrils and to Mallory's fibroglia fibrils. These are found in small numbers in a great variety of tumors. Whether these fibrils result from a reverse change from blue to red or whether they are newly formed is difficult to determine, I am disposed to regard the latter as probable; in the first place, because this corresponds to Mallory's opinion, and in the second place, because one cannot distinguish the change from a degeneration process. In certain rapidly growing tumors granular material reacting to Weigert's stain is found which corresponds to the granular fibrin of the physiologists, which, as stated in my previous paper, does not develop into fibrous tissue.

This explains the paucity of stroma in certain rapidly growing tumors.

In exact harmony with these changes in epithelioma and in the Sudan experiments are the changes observed in x-ray dermatitis and beginning epitheliomas resulting from the exposure to the x-rays. These lesions seem to supply the missing link between the Sudan experiments and spontaneous epithelioma. In the x-ray lesions, investigators are in agreement that the changes are primarily in the connective tissue. We have in each of these conditions, the Sudan experiments, the ray proliferations and spontaneous cancer, cell proliferation and changes in the connective tissue and an attempt to determine which is primary is not an easy matter. In the ray tumor and particularly in the Sudan experiments the changes in the connective tissue precede epithelial cell proliferation and there is much collateral evidence that the same is true in cancer.

In the summary of such collateral evidence we are reminded that it was the theory of Thiersch that the primary factors lay in the connective tissue. The fact that blood-vessels are not formed in granulation tissue after the fibrin stage has passed has seemed to me to be in accord with the observation already made that epithelial cells do not develop in the presence of normal connective tissue. These changes in the connective tissue are attended by round-cell and leukocytic infiltration. That cancer is most apt to develop at sites where chronic irritation is most apt to occur is perhaps the most constant factor in cancer etiology. The evidence of inflammatory reaction is present before the epithelial proliferation and the change in the connective tissue occur; in the presence of such irritation it seems but logical to assume as a working hypothesis that the changes in the connective tissue stand in a causal relationship. The x-ray cancer and the epithelial cell proliferation produced by Sudan can be pretty definitely demonstrated as being constantly preceded by connective tissue changes. If the hypothesis that the primary change occurs in the connective tissue be accepted, the nature of that primary change is yet to be determined. In these studies several points have appeared which are of interest. The fibrin fibrils occur about cancer nests, but their causal relation to the developing epithelial cell cannot be determined. The fact that the leukocytes are present in these processes preceding cancer development may in some remote way have to do with the process of fibrin formation. It would seem, however, that this is expressive of reaction against the invading epithelium, just as fibrin, then fibrous tissue forms in granulomas and not as a causal factor. Returning to the experiments with Sudan, we are reminded that those dyes only which have the power of combining with lipoids *in vivo* have this power and that they are all dyes belonging to the OH group. The negation of the properties of the lipoids may permit the processes of coagulation—that is, fibrin formation—to become active.

We know also that cancer cells have a predilection for metastatic formation in certain situations. These sites of predilection correspond in a rough way with the class of phenomena above considered.

These facts have a constant harmony which makes a specific significance probable. In a previous paper² I attempted to follow out on clinical grounds some of the circumstances under which cancers develop. In most situations irritations plus an alkaline secretion are coexistent. This crude hypothesis may have some value as a

2. Hertzler, A. E.: The Etiology of Epithelioma; A Laboratory and Clinical Study, *THE JOURNAL A. M. A.*, Feb. 8, 1908, p. 425.

working basis though a comprehensive statement cannot be entered into at this time. We may assume perhaps that a chemical balance has been broken. The latter would seem to be necessary in the explanation of the formation of metastasis, while the former might obtain as a preliminary factor in the development. I will remind you that epithelial metastases occur most frequently in those regions where leukocytes are most abundant (the lymph-glands) or where this prefibrin substance is most abundant (in subendothelial tissues). Metastasis, according to this hypothesis, becomes very closely related to the relation of leukocytes to blood coagulation. The relation of the leukocytes to chronic inflammations and the etiology of epithelioma halts at the same point. We must look to the physiologist for more light.

The relation of the fibrin fibers, of which I speak, to the fibroglia fibrils of Professor Mallory requires a word. They react to the same dyes and are found in the same situations, but I am at a loss to know how to go about establishing proof of their identity. Identity cannot be proved by fibrils produced otherwise than in wound-healing where the age of the fibrils can be accurately determined.

Studies in fibrils less than forty-eight hours old have convinced me of their identity. I desire to say a word about their relation to the cells which he believes produces them. I cannot agree that they may be produced by the cell, for the simple reason that they develop before the advent of the cells and in situations inaccessible to cells. They bridge areas not possible for cells to traverse. Their arrangement according to mathematical laws corresponds to the assumption that they are precipitated out of an amorphous exudate, while the arrangement cannot be explained by the assumption that they are the product as such of the cells; furthermore, at no time can any organic connection between the fibrils and the cell be demonstrated. What influence the cell may have in changing them from fibrin to fiber or what part the cell may have in the production of the material forming the amorphous exudate out of which the fibrils are precipitated by the action of some other factor, I cannot enter into here.

SUMMARY

1. The formation of fibrin is the first step in the formation of connective tissue.
2. Epithelium develops in the presence of fibrin only.
3. In the artificial production of epithelium by Sudan III the first changes are in the subepithelial connective tissue.
4. In *x*-ray cancer the epithelial development is preceded by changes in the connective tissue which are tinctorially identical with those which take place in the Sudan experiments.
5. The changes in the connective tissue in beginning epithelioma are tinctorially identical with those in the Sudan experiments and in *x*-ray cancer.
6. The factor most constant in cancer etiology is chronic irritation. Chronic irritation produces the changes above enumerated.
7. The sequence of phenomena above enumerated suggests as a working hypothesis that the etiology of cancer is associated with the lessened acidophilic properties of the connective tissue whereby it approaches fibrin in tinctorial reaction.

402 Argyle Building.

A FURTHER CONSIDERATION OF THE QUESTION OF PERMANENT DRAINAGE IN CERTAIN CASES OF PANCREATITIS *

LE GRAND GUERRY, M.D.
COLUMBIA, S. C.

At the Virginia Hot Springs meeting of the Southern Surgical Association, I read a paper entitled "A Plea for the More General Use of Cholecystenterostomy in Certain Cases of Pancreatitis." I said in part that I believed that the operation advocated was thoroughly consistent with the nature of the diseased process, namely, a chronic inflammation of the pancreas, whether the etiologic factor were gall-stones or infection independent of and apart from stone formation.

From what I have been able to read in books and the current literature, and have observed in the actual work of many large clinics, the operation of cholecystenterostomy in its relation to chronic or interstitial pancreatitis is not as generally in use by surgeons in this country as it should be.

The whole proposition hinges on this point: It appears that quite a number of cases of chronic pancreatitis require more or less permanent drainage to effect a symptomatic and physiologic cure; that if the pancreatic inflammation has progressed to a certain stage the operation of cholecystotomy does not afford drainage for a sufficient length of time to allow the trouble to subside; indeed, there is a point reached when the pancreatitis itself is incurable, although the symptoms due to insufficient bile drainage may be relieved by operation. In other words, if the pancreatitis which started as interstitial lasts long enough, the islands of Langerhans become involved and we then have the so-called inter-acinous pancreatitis with its resultant diabetes.

When we consider the comparatively small size of the bile channels, the very low blood-pressure under which bile is secreted, the slight force behind to force it on to the intestinal tract, it is easy to understand that in those cases in which the common duct passes through the head of the pancreas a comparatively slight obstruction would be sufficient to cause the damming back and absorption of the bile. I shall briefly relate a case which is typical of quite a number to serve as a concrete illustration.

M., a man aged 38, married, several years ago had an attack of pain in the gall-bladder region with very slight jaundice, pain occasionally radiating under right shoulder, slight fever, indigestion, etc.; these attacks would occur at frequent intervals; there was a gradual weakening and loss of flesh; no sugar in the urine. The diagnosis was gall-stones or pancreatitis (chronic) or both. At the operation a slightly distended gall-bladder with no stones was found; the bladder was full of the characteristic ropy, tarry, black bile with well-marked colon-bacilli odor to it. The pancreatic head was markedly enlarged. Operation consisted in cholecystotomy after the common duct had been thoroughly explored and a probe passed into the duodenum to insure the patency of choledochus; drainage was continued for three months, and during this time the patient not only gained 20 pounds in weight, but was completely restored to health. This is a sentence from a letter received three months after the wound had been closed: "Dear Doctor: I have just had another attack very much worse than any before my operation; what shall I do?" On my advice, the patient returned to the hospital, at which time cholecystenterostomy was done, with the result that he has been permanently relieved from all his symptoms and is now in perfect health.

* Read in the Section on Surgery of the American Medical Association, at the Sixty-first Annual Session, at St. Louis, June, 1910.