hyperacidity, for insufficiency of the pylorus or spasm of the pylorus. We must consider that these are merely symptoms and our examination has just begun with the diagnosis of dyspepsia. We are simply on the way toward a diagnosis. Recently I operated on a perforation of the duodenum, and the physician said that that was his second case within a month in persons who had been for two or more years under the care of a gastroenterologist for dyspepsia. It is not an easy matter to make a diagnosis, even when we go at the case very systematically. I have under my care at present a patient sent to me for ulcer of the duodenum. I had gone to work on the diagnosis by ruling out one thing after another, and after her eleven days of absolute constipation and a previous history of disturbance at times, I came to the conclusion that this was only a case of eyestrain. I have during the year every month some cases of dyspepsia due to eyestrain sent in for operation for gall-stones, duodenal ulcer, cancer of the pylorus, and my position is always this: So long as eyestrain does cause such symptoms in a certain proportion of cases, let us find the cases. That is a reasonable ground to take. Few of our ophthalmologists are competent to make this diagnosis. In the case referred to the patient's eyes have been examined by a most competent ophthalmologist. I asked him a few days ago whether he worked this case out with reference to imbalance of the eye muscles, and he told me he had not. We must not forget that similar symptoms are caused in a certain proportion of cases by gall-stones. Cases of gall-stones are often found at the deadhouse, in which it is said that the patient had no trouble from the presence of such stone—who knows? Such patients may have suffered for months or years or even a lifetime and with no diagnosis made, even though they told the physicians that they had gall-stones, and he said that it was not so.

DR. ARCHIBALD MACLAREN, St. Paul, Minn.: The pseudoperitonitis which Dr. Hertzler so well described seems to me to need immense consideration. The peritoneum does form a false membrane over each abscess, but Nature, it seems to me, is well able to take care of that pseudoperitoneum, as is well seen in so many operations for pelvic inflammatory disease. I remember years ago in the early history of abdominal surgery, where in a certain case the wall of a large pelvic abscess had been stitched to the peritoneal incision. Six months later I had to open that abdomen for some other cause and found nothing but a single thin fiddle string left. The dissection and stitching up of these flaps may occasionally be a good thing, but usually it is rather meddlesome surgery. Dr. J. N. Jackson recently described a false membrane which forms on top of the cecum and which, in his judgment, is to be dissected off. I have frequently seen the same result produced by simply dividing that membrane and loosening up the colon.

Dr. Deaver is unquestionably correct in regard to the difficulties of diagnosis in abdominal troubles. The wonder to me is that any court of law ever goes right; because these people come to us and try to tell their story, and there is no reason why they should not do so. And still, after half an hour's time spent in cross-examination, they will often say: "Why, I did have an attack of inflammation of the bowels two or three years ago." You can manufacture testimony for these people without any trouble, and the great difficulty is to avoid doing this very thing. In regard to gallstones, which have given no symptoms. When the patient is recovering from the operation where slumbering gallstones have been discovered and removed one can go back and always get a history, without any question, of gallstone dyspepsia so well described by Dr. Deaver.

DR. A. E. HERTZLER, Kansas City, Mo.: Dr. MacLaren's statement with regard to the "string" is correct; we all know that such strings produce intestinal obstruction. I want particularly to call attention to the reason why some adhesions are temporary and some permanent. As I stated in the beginning, there is the ordinary phenomenon of blood coagulation; when that process goes on undisturbed permanent adhesions result. In cases in which there are processes that ordinarily go on to precipitation the fibrin is not absorbed. In the last type referred to by Dr. MacLaren, the treatment

is simply that of any varicocele—the ligation of the veins. Such ligation will relieve the congestion, just as ligation of the veins does in obstructive phenomena in other places.

DR. J. B. DEAVER, Philadelphia: It seems to me now to be a question of the education of the public, just as in the matter of appendicitis. If we will educate the public they will say as they do now in appendicitis: "Doctor, I want my appendix taken out. Doctor, I want my abdomen opened if you think there is a chronic ulcer." We must do with the gastroenterologists as we did with the gynecologists, set them aside. We must do with the gentlemen who are washing out stomachs and electrocuting them, acting on the minds of the patients as we did with the gynecologists, and it leaves, therefore, only the internist and the surgeon to handle these patients.

Another class of men who have only a very limited sphere are the roentgenologists. If we wait until a lesion is demonstrable by the Roentgen ray, it is too late for surgery to accomplish anything. We must recognize the early symptoms and not be afraid of advising abdominal exploration. I have heard men who held high positions say that it is a terrible responsibility. We must get that out of our heads. We must operate on our patients early if we wish for results.

RESULTS OF THREE YEARS' EXPERIENCE IN BACTERIAL IMMUNIZATION *

B. A. THOMAS, A.M., M.D.

Assistant Instructor in Surgery, University of Pennsylvania;
Assistant Surgeon, Out-Patient Department, University Hospital;
Surgeon to the Children's Surgical Dispensary, Presbyterian Hospital; Surgeon to the Surgical Dispensary
St. Christopher Hospital

PHILADELPHIA

Two years ago in a contribution¹ read before the Philadelphia County Medical Society on the subject of opsonins, I made the following statements: 1. "In the heralded unlimited field of therapy, where the opsonic index is ordained to play a star rôle, its utility seems destined to pass into oblivion, not only because of its inconstancy in agreement with the clinical symptomatology, but especially because of its impracticability." 2. "Although my experience thus far with artificial auto-inoculations has been most gratifying and my results in close harmony with those of other observers, I am disposed to believe that there is a tendency to overvaluation even in this method of therapy." To-day I am more than ever convinced of the correctness of that stand and I doubt very much if at present Wright himself believes, literally, his statement promulgated a few years since, that "by means of these bacterial vaccines we have, in the power of raising the antibacterial power of the blood, with respect to any invading microbe, out of all comparison, the most valuable asset in medicine."

It is ever a fact in medicine that when truly important discoveries are announced the pendulum of applicability and utility, at first, either swings too far or not far enough. Valuable, indeed, as bacterin therapy has proved itself to be in an extensive category of affections to which the human organism is subject, it was and still is, in a measure, the victim of the former evil. Time and experience have gradually shaped the destiny of opsonotherapy or bacterial immunization, and the practitioner conversant with the current medical literature

* Read in the Section on Pharmacology and Therapeutics of the American Medical Association, at the Sixtieth Annual Session, held at Atlantic City, June, 1909.

* From the William Pepper Laboratory of Clinical Medicine of the University of Pennsylvania.

1. Variability and Unreliability in the Determination of the Opsonic Index, THE JOURNAL A. M. A., Oct. 12, 1907, xlix, 1249.

has already formed a more or less correct judgment on the merits of biologic therapeutics.

That inoculation by bacterin, in properly selected cases, plays an extremely important rôle in convalescence is indisputable. That failure to obtain expected results is due often to ignorance or a misapplication of this particular kind of treatment in contraindicated cases, thereby unjustly placing active immunization in disrepute with many clinicians, is unquestionable. I dare say that were a ballot taken of average physicians as to whether or not active immunization should be employed in a given case of septicemia, a positive vote would be cast. From my experiences nothing could be more erroneous. The defenses of the organism have already been exhausted and broken down, which accounts for the occurrence of the bacteriemia. How illogical, therefore, the introduction of an antigen, in the hope of stimulating these body cells to the production of antibodies, when their supply has already been consumed! We must not expect the impossible from bacterin therapy. The condition can be likened to the overzealous, but indiscreet jockey, who vainly hopes to win in the close finish race by relentlessly flogging his exhausted thoroughbred. Scientifically considered from the standpoint of anaphylaxis, it is not improbable, in cases of bacteriemia, that the inoculation of bacterin may not only do no good, but actually result harmfully. The organism is already in a state of hypersusceptibility induced by an overwhelming autogenous antigen, when, if the theory of Vaughan be correct, albumin in the body is broken up into the toxic and non-toxic group. At the time of the heterogenous injection of dead bacteria, the body attacks the albumin so vigorously and releases so much of the toxic group that it is overcome by the poisonous radical. On the other hand, late in the course of the disease, if the patient survives, it must be recalled that the inoculation of dead bacteria may act beneficially. Such has been the experience of Rosenow² in chronic septic endocarditis. I have had no success in this class of cases, and when improvement or cure has ensued I have been inclined to attribute it to Providence rather than to bacterin therapy.

Finally, overdependence on the opsonic index as a guide, and carelessness or inattention to certain clinical signs and symptoms, be they ever so slight and seemingly trivial in nature, or an irrepressible enthusiasm to "push the treatment" as soon as improvement becomes manifest, invariably leads to disappointment, if not to utter failure and disaster. This is particularly true of tuberculous affections, in which a prolonged, tedious and cautious course of treatment of six to twelve months' duration must be instituted.

The unanimous interest in and practical importance of active immunization make it the imperative duty of every practitioner, individually and collectively, to determine as early as possible precisely in what class of affections bacterin therapy shall and shall not be utilized. The problem is shrouded in many difficulties owing to the variability in the application of the treatment, so that results necessarily differ. For instance, some observers rely implicitly on the opsonic index as the guide in dosage; others maintain that the clinical symptomatology is all-sufficient; some pin their faith to autogenous bacterins; others claim that the stock preparations are equally or more meritorious; some advocate inoculations, comparatively, of large size; others, minimum dosage; some draw their conclusions following bio-

logic therapy alone; others refuse to discard old measures of proved worth and insist on associating bacterin therapy only with the former; some are gifted with excellent judgment and possess keen powers of observation, while others are indifferent, careless and overenthusiastic or prejudiced. These disadvantages can be overcome only by a persistent effort on the part of the profession to continue the search for the truth by the multiplication of case reports.

It is very easy in noting results following an attempt at immunization to become either an optimist or a pessimist, depending on whether one is an advocate or a doubter of the method. In this report it has been my avowed purpose to communicate the results of all cases, good, bad and indifferent, of all cases treated, in the belief that only by such statements can we ever hope to solve this question in its entirety. Certainly, the report of sporadic cases, with only brilliant results, from time to time, may lead to erroneous conclusions.

It has been my fortune to employ bacterin therapy in 106 various affections, among which the following bacteria were isolated for the preparation of autogenous bacterins. In a few instances, notably in gonorrheal and tuberculous affections, a stock bacterin was employed.

Bacteria.	Infections.
1. <i>Micrococcus albus</i>	18
2. <i>Micrococcus aureus</i>	38
3. <i>Bacillus coli communis</i>	6
4. <i>Bacterium ferrugineum</i>	1
5. <i>Bacillus fluorescens</i>	2
6. <i>Micrococcus gonorrhææ</i>	19
7. <i>Bacillus proteus vulgaris</i>	2
8. <i>Bacterium punctatum</i>	1
9. <i>Bacillus pyocyaneus</i>	3
10. <i>Streptococcus pyogenes</i>	10
11. <i>Bacillus tuberculosis</i>	19
	119

In all, fifty distinct diseases were treated. Both by desire and virtue of circumstances, the time-honored measures of proved worth were never abolished from treatment during the periods of bacterial inoculations, because I have always considered bacterin therapy as an accessory, not as a specific agent. It was never intended to be, nor will it ever be, a "cure-all," but should be regarded as an aid to assist Nature in properly selected cases, to combat infection by fortifying the organism through the production of immunization by antibodies. Moreover, it is the acknowledged duty of the physician to employ all the useful measures at his command in the treatment of his patient and not to experiment with human life in order to determine the efficacy of a new remedial agent, although by such a course the problem might be speedily solved. Consequently, in many instances, it has been a difficult matter to decide how much of the cure should be attributed to the usual method of treatment, and how much to bacterial immunization. I have striven throughout to give an unprejudiced opinion, and in dubious cases have refrained from placing the credit to bacterin inoculations. Nevertheless, in certain classes of cases, the results have been consistently so striking that the value of inoculation by dead bacteria is undeniable.

ABSCESSES

A review of the table of cases indicates that in the treatment of abscesses a fruitful field is afforded for bacterin therapy. Two deaths have occurred in my series, but they were both instances in which active

2. Rosenow: Jour. Infect. Dis., April, 1909.

TABLE OF CASES

No.	Case.	Age.	Disease.	Bacteria.	Duration of Disease.	Bacterin.	Number of Inoculations.	Average Dose of Inoculations.	Duration of Treatment.	Remarks.	Result.
1	W. M.	3	ABSCESS Axillary.	B. tuberculosis. M. aureus and albus.	1 yr.	Antigenous.	8	B=100,000,000; T=600,000 mg. C=250,000,000; S=50,000,000.	3 mos.	General tuberculosis. Mixed infection. Pyemia. Pulmonary hemorrhage (12 oz. a day) and malodorous. Still under treatment. Negative for T. B.	Death.
2	A. G.	23	Pulmonary	Streptococcus pyogenes.	12 yrs.	+	3		10 dys.	Expectoration very profuse (12 oz. a day) and malodorous. Still under treatment. Negative for T. B.	Negative.
3	L. C.	4	Multiple subcutaneous	M. aureus.	10 dys.	+	2	200,000,000.	2 wks.	Complicated convalescence following extensive burns.	Cure.
4	J. Z.	22	Multiple subcutaneous	M. aureus.	1 wk.	+	3	400,000,000.	10 dys.	Complicated convalescence following typhoid fever.	Cure.
5	A. B.	24	Pelvic	M. aureus.	3 mos.	+	6	500,000,000.	1 1/2 mos.	Spongy left in abdominal wall at time of operation causing abscess and rectal fistula.	Improvement.
6	R. G.	22	Pelvic	B. pyocyaneus. B. coli communis.	3 wks.	+	1	400,000,000.	2 wks.	High irregular fever. Very septic. Followed laparotomy. Pleuropneumonia. Sputum = T. B.	Death.
7	W. F.	40	Subdiaphragmatic.	B. tuberculosis.	2 wks.	+	12	300,000,000.	5 wks.	Suppuration following operation was very profuse and persistent.	Cure.
8	B. S.	17	Jaw	M. aureus.	1 mo.	+	19	250 to 800,000,000.	19 wks.	Osteomyelitis was associated with the persistent suppuration. Progressive gain in weight occurred.	Cure.
9	R. G.	22	Abdomen	B. coli communis.	3 wks.	+	1	400,000,000.	2 wks.	See Case 6. Very septic from pelvic abscess. No tuberculin treatment.	Death.
10	W. M.	3	Knee	B. pyocyaneus.	1 yr.	+	8	B=100,000,000; T=600,000 mg. C=250,000,000; S=50,000,000.	3 mos.	Pulmonary hemorrhage. See Case 1. General tuberculosis. Mixed infection. Very septic.	Death.
11	H. G. L.	37	Malar.	B. tuberculosis.	1 yr.	+	8	500,000,000.	2 1/2 mos.	Persistent sinus in rt. malar region for months, on which the best surgeons in Philadelphia had operated. Bier treatment.	Cure.
12	U. W.	62	Mouth	M. aureus.	2 wks.	+	3	S=100,000,000; M=200,000,000.	3 wks.	Sinus resulted following initial lymphadenectomy for carcinoma of lip.	Negative.
13	G. C. F.	57	In ano.	Streptococcus pyogenes.	3 mos.	+	6	200,000,000.	6 wks.	Small slightly suppurating sinus in anal region associated at times with induration. Bier treatment.	Negative.
14	A. B.	24	Rectal	M. albus.	3 mos.	+	6	500,000,000.	1 1/2 mos.	Formed from pelvic abscess. See Case 5.	Improvement.
15	A. B.	24	Vesical	M. aureus.	3 mos.	+	6	500,000,000.	1 1/2 mos.	See Cases 5 and 14. Subsequently entirely recovered. Undetermined as to whether fistula is vesical or urethral.	Improvement.
16	M. McK.	17	Thoracic.	M. aureus.	4 wks.	+	3	400,000,000.	3 wks.	Developed pleurisy and pleural effusion following appendectomy. Rib resection. Drainage. Suppuration. Has had two or three operations but discharge has always persisted. Secstrum removed. M. albus disappeared. Bier treatment associated.	Death.
17	A. L.	18	Thoracic.	B. coli communis.	1 yr.	+	17	400 to 800,000,000.	5 mos.	Persistent suppuration from laparotomy wound following gastroenterostomy for inoperable carcinoma. Infection of laparotomy wound following partial resection of stomach.	Marked improvement.
18	M.	45	Gastroenterostomy	M. aureus.	2 wks.	+	1	80,000,000.	5 dys.	Persistent suppuration from laparotomy wound following gastroenterostomy for inoperable carcinoma. Infection of laparotomy wound following partial resection of stomach.	Improvement.
19	M. B.	45	Partial gastrectomy	B. coli communis.	3 wks.	+	2	100,000,000.	2 wks.	Following operation a typical phagedena of abdominal wall occurred. Completely stopped following first inoculation.	Cure.
20	DeG.	50	Erosion of bubo	M. aureus.	1 wk.	+	4	300,000,000.	3 wks.	Following operation a typical phagedena of abdominal wall occurred. Completely stopped following first inoculation.	Improvement.
21	P.	40	Fracture of patella	M. aureus.	5 dys.	+	3	250,000,000.	2 wks.	Patella was wired and infection occurred between fascia and skin. Patient very fat.	Improvement.
22	D.	48	Breast amputation	B. pyocyaneus.	2 wks.	+	4	180,000,000.	2 wks.	Persistent suppuration following operation. Patient had renal complications and very stout.	Death.
23	F.	30	Hypoglossofacial anastomosis	Streptococcus pyogenes.	2 dys.	+	2	75,000,000.	1 wk.	A diffuse cellulitis occurred on the side of the neck operated. Patient's incision was not closed.	Cure.
24	D.	30	Craniotomy	M. aureus.	2 wks.	+	3	200,000,000.	3 wks.	No appreciable effect. Fungus cerebri developed.	Negative.
25	J. L.	35	Craniotomy	M. albus.	3 wks.	+	2	300,000,000.	2 wks.	Mental aberration prevented continuation of treatment. Patient refusing to have the operation needed.	Negative.
26	McE.	40	Craniotomy	M. aureus.	2 wks.	+	8	400,000,000.	7 wks.	Mental symptoms and symptoms of meningitis supervened. Suppuration steadily decreased and temperature fell.	Improvement.
27	M. T.	10	Chronic suppuration.	B. ferrugineum.	9 yrs.	+	6	100,000,000.	5 wks.	Immediately following inoculations, discharge from ear has increased. Still under treatment.	Negative.
28	S. S.	19	Tibia, acute.	M. aureus.	3 dys.	+	3	150,000,000.	10 dys.	A blood culture early showed Micrococcus aureus following operation. Treatment was futile. Patient worse. Still under treatment.	Negative.
29	J.	44	Tibia, chronic	M. aureus. M. albus. B. fluorescens.	6 mos.	+	1	400,000,000.	1 wk.	Following compound fracture of leg, a sinus has persisted for months. No evidence of caries.	Improvement.

Downloaded From: <http://jama.amanetwork.com/> by a University of Arizona Health Sciences Library User on 05/25/2015

TABLE OF CASES—(Continued)

No.	Case.	Age.	Disease.	Bacteria.	Duration of Disease.	Bacterin.	No. of Inoculations.	Average Dose of Inoculations.	Duration of Treatment.	Remarks.	Result.	
72	G.	65	Neck	CARBUNCULOSIS	M. aureus.....	3 wks.	+	2	880,000,000.....	5 dys.....	Delirious for two weeks. Pyelonephritis. Pneumonia. Myocarditis. Heroic stimulation. Urine and lungs cleared up immed. Carbuncle speedily granulated. Small incision for drainage. Bier cup. Duration of neck healed very speedily. Bier cup. Duration of convales. abscess of ischio-rectal space formed. Incision and drainage. Bier cup. Incision and drainage. Bier cup. Stock bacterin prepared from eight different strains of cultures employed. Incision and drainage. Bier cup. Recurrence. Incision and drainage. Bier cup.	Improvement.
73	C. S. Jr.	45	Neck	M. aureus.....	2 wks.	++	2	400,000,000.....	5 dys.....		Cure.	
74	S.	42	Neck	M. aureus.....	2 wks.	++	3	450,000,000.....	2 wks.....		Cure.	
75	McM ..	34	Neck and forearm	M. aureus.....	3 wks.	++	2	400,000,000.....	2 wks.....		Cure.	
76	S. F.	35	Chin.	M. aureus.....	4 wks.	++	3	450,000,000.....	3 wks.....		Cure.	
77	E. C. B.	29	Neck	M. aureus.....	1 wk.	++	4	500,000,000.....	3 wks.....		Cure.	
78	B.	33	Neck	M. aureus.....	2 wks.	++	2	400,000,000.....	2 wks.....		Cure.	
79	W. M.	3	Elbow	TUBERCULOSIS	B. tuberculosis	1 yr.....	+	8	B.=100,000,000; T.=400005 mg.	3 mos.....	Mixed infection. Pyemia. General tuberculosis. Pulmonary hemorrhage. Patient greatly emaciated and grew steadily worse.	Death.
80	A. T.	18	Enteritis	M. aureus and albus	2 mos..	+	2	0.001 mg. "B. E." 0.001; "B. F." 0.001 mg.	2 wks.....		Death.	
81	C. B.	5	Epiphysitis, tibia.	B. tuberculosis.	4 wks..	+	3	B.=100,000,000; T.=0.0005 mg.	3 wks.....	Following third inoculation, which was less than its predecessor, temp. rose considerably and was maintained, child dying of miliary tuberculosis. See Cases 1, 10, 52 and 79. Pyemia.	Death.	
82	W. M.	8	Face.	B. tuberculosis	12 mos.	+	8	B.=100,000,000; T.=0.0005 mg.	3 mos.....	Pulmonary tuberculosis complicated treatment.	Death.	
83	K. D.	45	Hip.	M. aureus and albus	4 mos..	+	7	B.=300,000,000 T.=0.001 mg.	5 mos.....	Child very anemic and waxy in appearance.	Improvement.	
84	T. H., Jr.	5	Sacroiliac.	B. pyocyaneus.	6 mos..	+	9	200,000,000.....	2 mos.....		Improvement.	
85	G.	19	Hip.	B. coli communis.	4 wks..	+	19	0.002 to .005 mg.	9 mos.....	Intermission of five months without treatment. Beck's bismuth injection acted deleteriously and complicated inoculation treatment. Mixed infection. Fresh air and antituberculosis measures.	Improvement.	
86	T. M.	18	Hip.	B. tuberculosis.	3 wks..	+	15	B.=400,000,000; T.=0.0001	3 mos.....		Improvement.	
87	J. C.	4	Hip.	Streptococcus pyogenes	2 wks..	+	21	B.=200,000,000 T.=0.003	4 mos.....	Fresh air and antituberculosis measures.	Improvement.	
88	J. B.	8	Hip.	M. aureus and albus	5 wks..	+	12	B.=200,000,000; T.=0.002	10 wks..	Fresh air and antituberculosis measures.	Improvement.	
89	W. M.	3	Knee	B. tuberculosis.	12 mos.	+	8	B.=100,000,000; T.=0.0005	12 wks..	See Cases 1, 10, 52, 79 and 82. Pyemia.	Death.	
90	J. W.	3	Mastoid.	M. albus and aureus	1 wk..	+	3	0.0001 mg.....	10 dys.....	Temperature dropped to normal quickly. Wound granulated rapidly. General tonic treatment. Wound healed speedily. At same time an epiphysitis of metacarpal bone subsided.	Improvement.	
91	N. A.	7	Mastoid.	B. tuberculosis.	2 wks..	+	4	0.0001 mg.....	4 wks.....	Immobilization of metacarpal phalangeal joints. General antituberculosis measures.	Improvement.	
92	N. A.	7	Epiphysitis, metacarpal	B. tuberculosis.	2 wks..	+	4	0.0001 mg.....	4 wks.....	Ordinary antituberculosis measures. Has gained 10 lbs. in weight. Peritoneum at time of exploratory laparotomy was literally covered with tubercles.	Cure.	
93	L. H.	30	Peritoneum.	B. tuberculosis.	1 wk..	+	56	0.0001 to 6 mg..	10 mos.....	See Case 83. Condition incidental to tuberculosis of hip, where there was a mixed infection.	Death.	
94	K. D.	45	Lungs	B. pyocyaneus	4 mos..	+	7	B.=300,000,000; T.=0.001	4 mos.....	See Cases 1, 10, 52, 79, 82 and 80. Pyemia.	Death.	
95	W. M.	3	Lungs	B. tuberculosis.	12 mos.	+	8	B.=100,000,000; T.=0.0005	3 mos.....	Shoulder resected years before. Anemic. Cardiac dilatation. Developed pelvic abscess. (T. B.) Obligated to suspend inoculations. No result.	Death.	
96	P.	45	Shoulder	M. albus and aureus	6 yrs..	+	21	0.00000002 mg.	8 mos.....	Discharged from hospital. Usual antituberculous treatment. Temperature fell.	Negative. Improved.	
97	J. T.	4	Hip.	B. tuberculosis.	3 wks..	+	2	0.00000005 mg.	2 wks.....	Suppuration decreased markedly. Posas abscess.	Improvement.	
98	R.	38	Spine.	B. tuberculosis.	4 wks..	+	7	0.00000001 mg.	7 wks.....		Improvement.	
99	J.	20	Hip.	M. aureus.	3 wks..	+	3	100,000,000.....	3 wks.....		Improvement.	
100	C.	22	Spine.	B. tuberculosis.	2 wks..	+	4	300,000,000.....	4 wks.....		Improvement.	
101	C. Y.	14	Spine.	Streptococcus pyogenes	6 mos..	+	6	B.=200,000,000; T.=0.001	6 wks.....	Vitality very low. Iliac abscesses.	Improvement.	
102	H. H.	25	Spine.	M. aureus and streptococcus.	1 mo..	+	11	B.=350,000,000; T.=0.001	3 mos.....	Three weeks after injection of Beck's bismuth paste patient developed pulmonary hemorrhages and died. Psoas abscesses. Temperature steadily decreased. Gained weight.	Death.	
103	E. J.	5	Spine.	B. tuberculosis.	4 wks..	+	21	B.=400,000,000; T.=0.002	4 mos.....	Suppuration lessened markedly. Temperature lower.	Improvement.	
104	S. Y.	4	Spine.	M. aureus.	3 wks..	+	5	200,000,000.....	4 wks.....	Suppuration decreased. Temperature lower.	Improvement.	
105	A. M.	4	Spine.	B. tuberculosis.	7 wks..	+	5	200,000,000.....	4 wks.....	Associated with sacro-iliac disease. Anemia.	Improvement.	
106	T. H., Jr.	5	Spine.	B. fluorescens. B. tuberculosis. B. proteus vulgaris. B. coli communis.	6 mos..	++	9	B.=200,000,000; T.=0.0005	2 mos.....		Improvement.	

immunization was clearly contraindicated. In one, Case 1, the patient was in an advanced stage of tuberculosis and the victim of pyemia; in the other, Case 6, the patient was very septic and had the complication of pulmonary tuberculosis with pleuropneumonia, superadded to which the pus from the abscess, together with other organisms, demonstrated the *Bacillus pyocyaneus*, a bacterium which, in my experience, has always been a bad omen and with the bacterin of which in therapy I have never been able to obtain the slightest success. I am unable to explain this fact, save that it is usually an infection superimposed on a previous suppuration, and is independent of the particular bacterium causing the primary infection; that is, it acts as a scavenger of the primary suppuration—a pus parasite, as it were. It would seem, therefore, illogical to expect that anything can be gained by stimulating, through inoculation, the cells of the human organism to the production of antibodies, by the bacterin of a bacterium which in itself may be entirely dependent on the action and products of a pre-existing infection. Especially is this true if we may accept the theory of the specificity of opsonins.

In multiple subcutaneous or superficial abscesses, either acute or chronic, the results have been most brilliant. In two instances, Cases 3 and 4, not a single abscess developed after the first bacterial inoculation

In conjunction with this I desire to call attention to the important fact that the opsonic index may be very variable and unreliable as a control of the dosage and spacing of the bacterial inoculations. For example, in

Chart 2.—Opsonic index. Patient, J. K.; acne vulgaris. The crosses on the line indicate bacterin inoculations, 880,000,000 *M. aureus*. Observe that after an initial rise the opsonic index persistently fell to the negative phase, where it remained, although a pustular acne existing for fifteen years permanently disappeared. The unreliability of the opsonic index could scarcely be better demonstrated.

Chart 1, the coincidence between the clinical symptoms and opsonic index was beautiful and all that could be desired. On the contrary, compare Chart 2 of the opsonic index of Case 65. In this instance, after an initial rise, the index fell into the "negative phase" and remained there in spite of the fact that, after four inoculations, the pustules disappeared permanently from the face of the patient, who had been a sufferer from acne vulgaris for fifteen years. From these and numerous other experiences, I have come to allow the clinical symptomatology to serve always as the paramount guide in the administration of bacterin inoculations. In my early work, and occasionally subsequently, the index was associated for what it might reveal in the nature of confirmation, but never to supersede the clinical symptoms as a factor in treatment.

In order to illustrate how ideally the symptomatology may be utilized in the conduct of bacterin therapy, to the exclusion of the opsonic index, permit me to present Chart 3 of Case 7 of my series.

SINUSES

Here, again, bacterin therapy is of undoubted value, although the results are not so striking, a much longer period being necessary to effect a cure. One cause partially responsible for this is that not infrequently carious bone or a foreign body lies at the bottom of the sinus. We cannot be expected to remove dead or dying tissue. This is the province of the surgeon. Another cause for the tardy improvement can be ascribed to the fact that these sinuses are frequently surrounded by dense fibrous tissue, crowding out granulations; hence the supply of blood and lymph necessary for the healing process. Then, too, the few lymph spaces that do exist have their orifices occluded by coagulated fibrin and necrotic tissue, preventing the lymph system from properly bathing the sinus wall.

Chart 1.—Temperature (zigzag line) and opsonic index (straight diagonal line). Patient, L. C.: burns one-third to one-half body surface; multiple subcutaneous abscesses; a, skin grafts to arm and leg; b, development of scalp abscesses; c, marked loss of weight and strength; d, many subcutaneous abscesses evacuated; e, pus cultured from new abscess demonstrated micrococcus aureus. The small crosses on the diagonal line indicate inoculation with *M. aureus* bacterin, 205,000,000 bacteria. Note the parallelism existing between opsonic index and improvement in the condition and health of the patient. Note also that no new abscesses formed after the first inoculation of bacterin and the rapidity of disappearance of the recurrent pre-existent subcutaneous abscesses.

and the pre-existent ones disappeared in remarkably short time. The clinical course of one of these cases is represented in Chart 1, associated with the temperature curve of which the phase of the opsonic index is plotted and the time of the active immunization noted.

This can be obviated to a degree, theoretically at least, as Wright has observed, by the irrigation of these tracts with a solution consisting of 0.5 per cent. sodium citrate in 5 per cent. salt or 10 per cent. sugar solution, which possesses the power of decalcification or defibrination and promotion of osmosis.

Death occurred in two of the five cases in the series, as will be observed, but it must be borne in mind, in justification to any therapy, that they were in condi-

Even in ambulatory cases it is my custom to see that the patient is provided with and instructed in the use of a clinical thermometer, so that he may take and record his temperature at least twice daily. This may appear an unnecessary matter, and in itself is probably worthless, but it cannot be too strongly emphasized that if we are to be governed by the clinical symptoms, and are desirous of obtaining the greatest success, we must not ignore the slightest aid, for it is only by judgment based

Chart 3.—Temperature. Patient, W. F. Subdiaphragmatic abscess drained per laparotomy; a, patient very comfortable; drainage satisfactory; b, patient complains of pain in hepatic region; c, leucocytes 20,000; d, excessive suppuration; e, suppuration variable, but very profuse; f, patient's appearance improved; feels much stronger; g, patient discharged from ward; h, treated in out-patient department; i, suppuration markedly decreased; k, discharge from wound very slight; l, marked gain in strength and health; m, no discharge; n, sinus healed; 1, pus demonstrates *M. aureus* and albus; 2, inoculations *M. aureus* bacterin $\frac{1}{2}$ c.c. This case illustrates how admirably active immunization can be conducted by resort to the clinical symptomatology as a guide, to the exclusion of the opsonic index.

tions not to be alleviated by any mortal measure. Patient 11 had undergone operation several times by some of Philadelphia's most distinguished surgeons for a persistent sinus in the malar region of the face. Whether cure in this case can be assigned to bacterin therapy, avoidance of interference or the application of Bier's vacuum cup, remains undetermined. The last, acting to promote the circulation of blood and lymph, is not to

on the total symptomatology, important signs as well as trivial details, embracing a consideration of the temperature, pulse, malaise, indisposition, headache, nausea, vomiting and local phenomena, as reactions in appearance and increase of discharge temporarily following inoculations, that success or failure depends.

FISTULAS AND EMPYEMAS

Whatever may have been the results of others in fis-

Chart 4.—Temperature. Patient, B. S.; osteomyelitis with suppurative sinus of jaw. Crosses on line indicate *M. aureus* bacterin inoculations; a, $\frac{1}{2}$ c.c. bacterin; b, $\frac{2}{3}$ c.c. bacterin; c, 1 c.c. bacterin; d, 1 c.c. bacterin; suppuration less; e, $1\frac{1}{4}$ c.c. bacterin; drain removed; f, sinus healed; 1, weight 120; 2, weight 121; 3, weight 117; 4, weight 118; 5, weight 124; 6, weight 125; 7, weight 126; 8, weight 123; 9, weight 122; 10, weight 127. Note the progressive increase in weight as the temperature, at first above normal and hectic, became subnormal, then rose and fluctuated on the normal line.

be underestimated in the treatment of these cases. Case 8 seemed to be of sufficient importance to warrant the preparation of Chart 4, for two reasons: first, to show again the value of control by the clinical symptomatology; second, to illustrate even in the presence of definite and persistent osteomyelitis, how success may be expected in assisting Nature by stimulating the defense of the organism.

tulas, especially in fistula in ano, my success in this class of affections has been practically nil, or only that ascribed to time and Nature, as will be observed in the three cases it has been my opportunity to treat.

It seems questionable whether or not bacterial inoculations are to enter deeply into the treatment of thoracic empyemas. One of my cases, from which the *Bacillus pyocyaneus* was isolated, was uninfluenced and pro-

ceeded to death. The other, which has been under treatment for a year, shows slight but unmistakable improvement.

POSTOPERATIVE WOUND INFECTIONS

Nine cases of this character have been treated by bacterin therapy, and, although the results have been satisfactory so far as cessation of suppuration and wound repair are concerned, it is impossible to place the credit entirely to the inoculation of dead bacteria. Those cases treated with stock bacterins have responded equally with those in which the autogenous preparation was used.

OTITIS MEDIA

In our series there occurs but one case of suppurative middle-ear disease from which repeatedly the *Bacterium ferrugineum* was isolated in pure culture. The patient has been a sufferer for nine years. He has received six inoculations during the past five weeks, and, aside from slight increase in the amount of discharge, there has been no noteworthy effect. Naturally the case has been

was a sufferer from exostoses of the os calcis, demonstrated by the x-ray, causing him extreme pain on walking and incapacitating him from work. It is hoped that by Neisser bacterin the further formation of osteophytes may be stayed and the patient immunized against the gonococcus so that an operation may be undertaken with the prognosis of a reasonable degree of success. After six inoculations the pain on walking has decreased markedly.

PROSTATITIS AND URETHRITIS

One patient with chronic gonorrheal prostatitis has been not only greatly improved, but practically cured. In justice to other therapeutic measures, be it said that urinary antiseptics, prostatic massage and the passing of cold urethral sounds were not omitted from the treatment.

I have had no experience with gonococco-bacterin therapy in the treatment of acute specific urethritis. Nevertheless, in a case of the chronic posterior type complicated by stricture, a cure has been apparently

Chart 5.—Temperature. Patient, S. S.: acute osteomyelitis of tibia, followed by septicemia. The arrow indicates a point at which blood culture showed *M. aureus*; the crosses indicate *M. aureus* bacterin inoculations. Note that the patient, as indicated by the fluctuating temperature, progressively grew more septic, and that bacterin therapy was not only useless, but possibly harmful.

under observation too brief a period, and conclusions should not be drawn from a single case.

OSTEOMYELITIS AND OSTEOPERIOSTITIS

I am fortunate in having both a typical acute and a chronic case of osteomyelitis for comparative study. The patient with the chronic condition has been a sufferer for months, the infection arising from a compound fracture of the leg. The inoculations have been few, but the result brilliant. The acute case, No. 28 of the series, arose from a hematogenous infection. Following operation, the temperature, after a temporary fall, began to fluctuate and a bacterial inoculation was immediately made. Not the slightest effect could be observed, and a blood culture demonstrated the presence of the *Micrococcus aureus*. Subsequently two more inoculations were made, but the result has not only been not beneficial, but possibly actually deleterious. I believe that we have here a beautiful illustration for the contraindication of active immunization in acute bacteremia. So instructive is this case that I have deemed it advantageous to show the temperature curve (Chart 5).

The only case of osteoperiostitis is one of subacute or chronic gonorrheal infection. The patient for a year

effected; all shreds having disappeared from the urine, employing the usual tests. Here, again, dilatation by sounds was associated with the biologic therapy.

CELLULITIS

In the majority of cases of this affection, no improvement could be definitely ascribed to bacterial immunization, although the patients recovered. In one instance, however, Case 34, the effect was so immediate on a rapidly spreading phagedena of the abdominal wall due to the *Bacterium punctatum* that there can be no question as to the efficiency of bacterin.

CYSTITIS

The report of one case³ can never solve any problem, and in this case a complete cure has not yet been secured, although, if the infection is primary and limited to the bladder, a favorable field seems to be afforded. An encouraging report has been received from Dr. Ohlmacher, of Detroit, who has continued the treatment in

3. Since the preparation of this paper, bacterin therapy has been applied to another case of cystitis due to the *B. coli communis*. Although slight improvement was perceptible during an abbreviated course of treatment, circumstances rendered its continuation impossible.

this case by the inoculation of autogenous *Bacillus coli communis* bacterin.

VULVOVAGINITIS

Through the courtesy of Dr. Howard C. Carpenter, I am privileged to report the results of gonococco-bacterin immunization on thirteen children treated at the Philadelphia Hospital. His report is as follows:

Chart 6.—Opsonic index. Patient, J. B. C.; recurrent erysipelas. The circles indicate opsonic readings; the crosses bacterin inoculations; a, inoculation 25,000,000 streptococci; b, 50,000,000 streptococci. Observe in this case that the "high tide of immunity" indicated by the opsonic index has been maintained in the "positive phase" for two and a half months, no inoculation having been given since March 22. It will be noted that this is entirely in accord with the clinical symptoms of the case.

1. The *Micrococcus gonorrhææ* was demonstrated by smears in each case prior to treatment.
2. Three to ten inoculations of a stock preparation were employed.
3. The earliest disappearance of the gonococcus was noted after the second inoculation.
4. Five million dead gonococci was the constant dose.
5. The period of treatment varied from nine to thirty days.
6. In one case in which no local treatment was employed, the result was most brilliant.
7. In every case the period of detention in the hospital was noticeably shortened.

TENOSYNOVITIS AND ARTHRITIS

The rôle played by bacterin therapy is largely conjectural in the treatment of tenosynovitis. The patients, acute and subacute suppurative types, finally recovered after periods of prolonged treatment, but such patients also recover without bacterial inoculations. The number of cases is too limited for conclusions.

My experience with infectious arthritis seemingly justifies the conclusion that the most favorable results can be expected in the subacute and chronic affections, especially those caused by the *Micrococcus gonorrhææ* in which infection either marked improvement or cure has always occurred. Indeed, in these cases the response to treatment frequently has been extremely prompt and the ultimate functional result better than by any other known method. Fifty million gonococci of a stock preparation constituted the average dose. From two to four inoculations were made over periods of three to four weeks. The results in other infections have been essentially negative.

ERYSIPELAS

Two cases of erysipelas have been successfully treated by biologic therapy in my series. One patient had an acute streptococcal infection of the face and neck. Recovery was prompt following one inoculation of 100,000,000 bacteria of the autogenous strain. Precisely the influence of the antigen cannot be determined even in this instance, as this infection is prone to run a short course.

The other case, No. 30, is one of peculiar interest and of sufficient importance for detailed description. I have merely reported the curve of the opsonic indices (Chart 6). This woman for thirteen years has been periodically, notably in the spring, the victim of recurrent attacks of erysipelas, involving both legs, thighs and genitalia. This infection, when her health is run down, is always superadded to a marked chronic hyperplasia of the loose connective tissues of both lower extremities. Beginning late in the winter, active immunization, employing a stock bacterin of the *Streptococcus pyogenes*, was instituted. Five inoculations were administered over a period of six weeks. The patient reports that she feels very well and that this is the first spring for a number of years that she has not been the victim of the attack of erysipelas. In this case, the opsonic index was utilized to advantage, as will be seen by consulting the chart. It would seem indisputable that bacterin therapy has a place in this type of case.

ACNE VULGARIS, FURUNCULOSIS AND CARBUNCULOSIS

Acne affords a fruitful harvest for active immunization. Every case has either been greatly improved or cured, and to the credit of bacterin therapy be it said that three of the cases were the most aggravated and obstinate types of pustular acne that it has been my fortune to observe. The duration of the disease previous to bacterial inoculations varied from two to fifteen years. Autogenous bacterins were always employed from the *Micrococcus aureus* or *albus*, isolated with equal frequency.

Of all the indications for bacterin therapy, furunculosis is probably the most brilliant. In no case have I failed to obtain a cure in more than five weeks; ten days being the shortest period of treatment. My rule has been invariably to culture the pus from the boil and prepare an autogenous bacterin. The *Micrococcus aureus* has always been the infecting organism.

No less satisfactory have been the results of therapy of carbunculosi by bacterial inoculations. Again, the *Micrococcus aureus* was isolated in seven cases. A cure always resulted, the longest period required by treatment being three weeks; the shortest, five days. In one in-

Chart 7.—Opsonic index. Patient, C. S., Jr.; carbunculosi. Circles indicate opsonic readings; crosses indicate *M. aureus* bacterin inoculations, to the left, $\frac{1}{2}$ c.c., to the right 1 c.c. Note the definite rise of the opsonic index from the negative to the positive phase during the treatment of five days' duration.

stance a polyvalent bacterin, prepared from eight different strains of the *Micrococcus aureus*, was used with a result equally successful with any in which the autogenous preparation was employed. The clinical symptomatology has been practically the sole guide in the administration of the inoculations. The curve of the opsonic indices depicted in Chart 7 demonstrates graphically the

rapid rise of the index often observed in the treatment of a case, No. 73, of short duration.

TUBERCULOSIS

The treatment of tuberculosis, in its manifold phases, constitutes my proudest effort. Broadly, therapy in these cases has been directed to the (1) pure and (2) mixed infections. We have heard much of tuberculin therapy, which, of course, is simply bacterin immunization by the tubercle bacillus or opsonotherapy, if you please, when the bacillus emulsion, "B. E.," product is employed.

The cases in my series should also be divided into the chronic and acute classes. With the acute class the results have been anything but encouraging. Here, again, my belief has been sustained that we must not expect success. It is probably true that in incipient tuberculosis we have to deal with a bacteriemia, and it has been my fortune, in the artificial inoculation treatment of such cases, invariably to experience failure. The results of bacterin therapy in my work with the early cases of tuberculosis certainly supports the findings and opinion of Rosenberger. In the subacute and chronic stages of the disease, even in the pulmonary type, I have reason to believe in the statement of Trudeau and others. A review of the cases tabulated under "Tuberculosis" in the table of cases is almost self-explanatory and confirms this belief. Many of these cases, for divers reasons, have not passed to immunizing doses of tuberculin or to the stage of cure. A great number have manifested definite improvement. I desire to call attention to one case in particular, No. 93 in the series. An exploratory laparotomy revealed general tuberculosis of both the visceral and parietal peritoneum. The patient was a very sick young woman and her life was practically despaired of. Tuberculin therapy by "B. E." variety was instituted in August, 1908. After a very tedious regime of treatment continuously since that time, she is at present positively unaffected in any way by inoculations, weekly, of 6 to 10 mg. of tuberculin, and has gained ten to fifteen pounds in weight. Her health seems perfect. I am not unaware of the fact that occasionally such patients recover after a simple laparotomy.

The particular phase of tuberculous disease to which I especially desire to direct attention is the mixed infection so constantly superimposed on and associated with Pott's disease and hip disease—indeed, any bone or joint tuberculosis. It is well known how frequently these complex mixed infections fail to improve and ultimately cause the death of the child. I do not desire to be overoptimistic, but I am very sanguine over my successes with bacterin therapy in this class of cases, and do not hesitate to say that in no instance in which proper surgical treatment and antituberculous measures were associated with bacterial inoculations have I failed to observe improvement in the health of the child. I have made it a practice to culture and reculture the pus from time to time for the preparation of autogenous bacterins. Commonly as many as three different bacterins have been alternated in the immunization process. So soon as the temperature falls to 100, or lower preferably, and remains permanently below that grade, tuberculin inoculations are begun. I have been inclined to believe, judging from the cases under my observation, that better results have attended the process of active immunization when, just as in tuberculin therapy, pure and simple, the treatment has been started with relatively small bacterin inoculations, progressively in-

Chart 8.—Temperature. Patient, E. J.: caries, tuberculous spine with psoas abscess and mixed infection. Stars indicate inoculations of tuberculin "B. E." and crosses, inoculations with *M. avium* bacterin; a, bacterin inoculation m. iv.; b, tuberculin inoculation 0.0005 mg.; c, tuberculin inoculation 0.0001 mg.; d, bacterin inoculation m. vi (new); e, bacterin inoculation, m. vi.; f, bacterin inoculation, m. vii.; g, tuberculin inoculation 0.00015 mg.; h, bacterin inoculation m. viii.; i, bacterin inoculation m. ix.; k, bacterin inoculation m. x (new); l, tuberculin inoculation 0.0002 mg.; m, tuberculin inoculation 0.00025 mg.; n, tuberculin inoculation 0.0003 mg.; o, operation; p, pus sterile; q, pus demonstrative *M. avium*; r, suppuration very slight; child has gained greatly in weight; s, cast applied and child sent to Atlantic City for the summer. This case illustrates ideally the gradually progressive, but definite tendency to a subsidence of the infectious process, represented by the restoration of the temperature curve to the normal line. Note the manner in which the bacterin and tuberculin inoculations have been carefully alternated. Observe, that depending on febrile rises at times the tuberculin inoculation was suspended, to be resumed at the next stated time, employing only half of the previous dose, which possibly was instrumental in causing the temperature rise. Note that the inoculations both by tuberculin and bacterin began as comparatively small doses, frequently repeated and increased slowly and cautiously throughout the course of treatment.

creased to the therapeutic limit, rather than by recourse to large dosage, thereby, in the former case, establishing immunity, and, in the latter, avoiding anaphylaxis. Apropos of the beneficial effect to be derived from treatment by this method, permit me to submit Chart 8, illustrative of Case 103 in my series.

So far as I am aware, I am alone in advocating the value and importance of alternating bacterial inoculations and tuberculin therapy in this type of tuberculous mixed infection. The inoculations should be given every four to seven days, the clinical symptoms permitting. The opsonic index is unreliable as a guide. Relative to tuberculin inoculations, I have used Bacillen Emulsion alone or Bouillon Filtrate alone, or a combination of the two, as practiced and advocated by Trudeau. The last, theoretically, seems to have the preference, but my results have been equally satisfactory with each. It all depends on the man behind the syringe.

It must be remembered that in the use of tuberculin we have a more potent instrument for evil than for good, unless it is properly administered. In careful hands, it serves as an invaluable aid in treatment and is Nature's own remedy to stimulate the bodily defenses. When tuberculin is so employed, the patients do better with than without tuberculin; their retention in the hospital is materially shortened and the complications, if they occur, are fewer and less severe.

CONCLUSIONS

From the study of the cases in this tabulated series the following conclusions seem justifiable:

1. The diseases contraindicated for bacterin therapy are the diffuse infections characterized by septicemia, pyemia and grave sapremia.
2. Those in which therapy by this agent is beneficial or curative are the superficial acute, subacute and chronic processes, especially the last two.
3. The acute cases, in which brilliant results can be uniformly expected, are those of acne vulgaris, furunculosis, carbunculosis and subcutaneous abscesses.
4. Subacute and chronic gonorrheal and tuberculous affections are amenable to bacterial immunization, and because of the impossibility and impracticability often of employing an autogenous bacterin the reliable stock preparations should be used.
5. Certain acute gonorrheal infections can be benefited.
6. It is questionable whether tuberculin therapy should ever be employed in very acute tuberculosis. Opinion is divided as to whether or not acute miliary tuberculosis and death supervened as a result of tuberculin therapy in one of our cases, No. 81 of the series.
7. The mixed infections in chronic tuberculous disease afford an important prospective field for *alternating bacterial inoculations and tuberculin therapy*.
8. Autogenous bacterins are always to be preferred over the stock preparations, and success or failure frequently depends on this fact.
9. Although the duration of the period of greatest potency of bacterins is undetermined, the best results have been obtained when the pus has been recultured and a fresh bacterin prepared every two to four weeks.
10. It is believed that the best effects, therapeutically, particularly in chronic cases, occur when the quantity of bacterin is slowly and cautiously increased during successive inoculations, thereby, as has been thoroughly demonstrated in tuberculin therapy, avoiding hypersusceptibility or anaphylaxis.

11. Therapy by both bacterins and tuberculins can be satisfactorily executed by keen observance of the clinical symptomatology. Reliance on the opsonic index as a guide is not only unnecessary, but often actually conducive to erroneous conclusions, owing to its variability.

12. Bacterin therapy, by virtue of its potency to do more harm than good, when unskillfully managed, will or should probably not become a universal therapeutic measure in the hands of the general practitioner, unfamiliar with bacteriology or work in the laboratory. Ignorance and wantonness are incompatible with ambition and energy, and an otherwise meritorious therapeutic agent thus abused will ultimately fall into disrepute.

13. Therapy by the employment of bacterins made from *Bacillus pyocyaneus* has been entirely useless.

14. Bacterins and tuberculins are not "cure-alls," but when intelligently used serve as invaluable aids to Nature in fortifying the bodily defenses, thereby accelerating convalescence, diminishing complications and promoting cure.

116 South Nineteenth Street.

PASTEURIZATION OF MILK*

ROWLAND G. FREEMAN, M.D.

NEW YORK

Public opinion in this country swings violently in one direction or another, occasionally on insufficient premises; and this, I think, applies to the present attitude among many pediatricists in this country concerning the pasteurization of milk.

We can all agree that what we want is a safe, raw milk, since we all wish, in the artificial feeding of infants, to approximate as nearly as is feasible the condition of the food as obtained by the infant at the breast.

The only questions for discussion, then, are: Can we obtain a safe, raw milk for the feeding of infants? And, if not, is milk injured by heating, and to what extent, and by what temperature?

There can be no question but that we can to-day feed all the infants of New York on a much safer raw milk than was possible five or ten years ago; but what is the extent of this element of safety? Take, for example, the best milks at present produced under the certified milk plan. How safe are these milks as a raw food for infants?

The danger of tuberculosis in such milk may be said to be fairly eliminated by well-aired, well-ventilated cow stables, and by the repeated tuberculin test. But tuberculosis is one of the lesser dangers in milk. The diseases concerning the spread of which we have most tangible and incontrovertible proof are typhoid fever, diphtheria, scarlet fever, and epidemic sore throat.

Our sanitary dairies do almost all that is possible to protect us from these diseases, but those of us familiar with this subject know that epidemics may occur from the sale of certified milks. No system of control can protect a milk-supply from a mild walking typhoid, or a typhoid-carrier among the employees, unless regular bacteriologic examinations are made of the urine and feces of the employees at frequent intervals; and I take it that no such system would at present be advocated.

Diphtheria has apparently been spread by the best of our milk-supplies; what protection have we against this in any raw milk?

* Read before the New York County Medical Society, Oct. 25, 1909.