interfered with. I have always used, however, a loop of ileum because resection of the ileum is safer than resection of the sigmoid. In all of my cases operative recovery was absolutely smooth, and the new vagina seems to take the place very satisfactorily of the natural organ. I was induced to use the bowel for this purpose because a very extended study of the literature of artificial vaginas had shown that in cases of entire absence all other methods resulted in almost complete failure, although full of promise in many cases when the patient left the operating-table. The favorite method of operating had been by transplantation of pieces of skin, to line the artificial canal. In all of these cases, however, to the reports of which I had access, the ultimate results had been very unsatisfactory owing to cicatricial contraction. By the use of the bowel a normal mucous membrane is provided, surrounded by normal connective and muscular tissue, and with an ample blood-supply. Theoretically and practically, therefore, such a vagina would seem to fulfil to the highest degree all the requirements.

The dangers of operation are simply those inherent to any other deliberate resection and anastomosis of intestine; but the operation is one which should certainly not be undertaken by a tyro in surgery. My first operation took over two hours, the last one about half that time, and I think that in an uncomplicated case the operation should be made within the hour. There is, of course, no hemorrhage to amount to anything, and if the field of operation is carefully protected there should be no shock. The danger, therefore, would be merely the danger of a possible infection from the intestine, and that danger in the hands of an expert is very slight. Nevertheless, the danger should be fully explained to the patient, who would then decide whether the operation would be worth while.

In addition to the four cases of my own, a successful operation has been reported by Mueller² and one by Mori.³ Thus we have six successful cases with no mortality. (So far as I can learn neither Mori nor Mueller refer in any way to my own previous work, although my publications preceded theirs by several years. I understand, however, that such a failure to give credit is in certain foreign medical circles not looked on as either unethical or dishonorable).

115 South Grant Avenue.

TORSION OF THE GREATER OR GASTROCOLIC OMENTUM

COMPLICATED BY ACUTE APPENDICITIS

W. H. LUCKETT, M.D.
NEW YORK

So very little is really known of the mechanical cause of the twisting of the omentum that the report of this case is interesting and may serve to throw some light on the subject.

Less is known of the symptom-complex leading to a correct diagnosis of such torsions. This is demonstrated by the fact that in some sixty odd cases recently collected, extending over a period of twenty-seven years since 1882, when the first report was made by Oberst, only one case was diagnosed as such before the operation.

Corner and Pinches lead one to be satisfied with a diagnosis at the operating table and not a differential one beforehand.

Fuller points out the difficulty of diagnosis and says that it is not at all remarkable or strange for a torsioned omental tumor to escape detection even during a laparotomy, unless the incision happened to be made directly over the tumor.

Richardson suggests an anatomico-pathologic classification according to the point of twisting, and asks for a report of all cases with an explicit statement as to the location of the tumor and the point of torsion. He thinks that if the possibility of torsion of the omentum be kept in mind in all cases of abdominal symptoms pointing to a strangulation a preoperative diagnosis of torsion of the omentum would become more frequent.

Weiner offers the very practical suggestion that if we cut down on a hernia that produced symptoms of strangulation and find only a strand of omentum in the canal we should investigate the intra-abdominal portion of the omentum to make sure that there is no torsion present.

Patient.—H. P. F., foreman by occupation, 32 years old, was admitted to my service at the Harlem Hospital, June 14, 1906.

Previous History.—The patient had an attack similar to the present one two years ago, from which he recovered. He was operated on ten years ago for right inguinal hernia.

Present Illness.—Two days previous to admission he was seized with very severe pains in the right iliac region, which continued unabated up to that time. He had been constipated since the beginning of the attack. There was no vomiting. He was in great pain on admission. His temperature on admission was 103 F., pulse 120, respiration 26.

Physical Examination.—General appearance—a well-developed, conscious male, very obese. Eyes—pupils contracted but reacted naturally; mucous membrane red and dry; lips scarlet; pulse regular in force and frequency; very rapid; respiration shallow and jerky; abdomen rigid and tense, general tenderness, tympanitis, pain more marked on right side. Urine—albumin with casts, hyalin, granular and epithelial. Arteriosclerosis was present. An indistinct mass could be made out to the right of the umbilicus, reaching down into the right iliac region. A cicatrix in the right inguinal region indicated that an operation for hernia had been performed. The wound had apparently healed by primary union, and there was no signs of recurrence. A diagnosis of appendicitis was made. A high leucocytosis and differential blood-count, with the mass and other symptoms suggested abscess.

Operation.—A long incision was made through the deep fascia on the outer edge of the right rectus muscle over the tumor. A quantity of bloody fluid escaped and a large mass of congested and partially strangulated omentum immediately presented itself in the wound, and was found to be adherent beneath the hernial cicatrix by the small thin band of adhesion at the lower end of the specimen seen in the illustration. This and several other small bands of newly formed inflammatory adhesions were easily broken up; as they were so small they were not ligated. The omentum was delivered through the wound and excised near the transverse colon. The appendix, which was found to be very markedly congested, was removed. There were, however, no adhesions between the omentum and the appendix. The abdomen was closed without drainage, and the patient left the operating room in good condition. June 18, four days after the operation, the wound was dressed and was found dry and clean. June 30 the patient was up and well.

Examination of Specimens.—The specimen of the appendix was small; there were no adhesions, though it was markedly congested, but probably not enough to account for the high temperature, pulse and leucocyte count, and surely not for the general congestion and the fluid in the abdomen.

The specimen of the omentum (see illustration) was 17 inches long, weight 9¾ ounces, and had 27 distinct complete twists or turns, the reverse to the thread of a right-hand screw. When partly untwisted it was found that the small band of adhesion to the peritoneal side of the hernia cicatrix

2. Mueller, A.: Formation of Vagina from Loop of Small Intestine, *München med. Wchnschr.*, Dec. 21, 1908; abstr. in *THE JOURNAL A. M. A.*, Jan. 29, 1910, liv, 419.

3. Mori M.: *Zentralbl. f. Gynäk.*, Jan. 1, 1910; abstr. in *THE JOURNAL A. M. A.*, Feb. 12, 1910, liv, 577.

probably came off from the lower right-hand corner of the omentum. The twists or turns as counted commenced near the attachment of the omentum to the transverse colon and ran all the way down the entire length of the omentum to the small band of adhesion to the hernial cicatrix.

With a few more twists the small band of adhesion would have parted and I should have reported this case as a pure intra-abdominal torsion without adhesions or the presence of a hernia. The result of the hernia operation ten years ago was perfect, and excepting the small spot to which the band was attached the peritoneum beneath the hernia scar was perfectly smooth, as demonstrated by palpation within the abdomen. Evidently a small tip of the omentum had either been ligated with

Specimen showing torsion of omentum and small band of adhesion at one end.

the hernial sac, or the omentum had subsequently become adherent at this point, thus affording a diagonal line from the upper left corner of the omentum at the transverse colon to the lower right hand corner of the omentum, around which the free lower left irregular triangle could and probably did revolve. If the adhesion played a rôle in the torsion it does not seem possible for such adhesion at the lower right border of the omentum to have been present for ten years without giving rise to an earlier twisting. Surely the torsion could not have been present for ten years, for there would have been adhesions between the twists that would have prevented the untwisting, which was fairly easily accomplished. If the adhesion was present for ten years without torsion, then some other force was necessary for its production. Was it the acute attack of appendicitis? The omentum has a habit of enfolding the appendix during certain inflammations. Was it making violent efforts to get to the appendix, or was the twisting a result of a purely exterior force such as the peristaltic wave of the intestines particularly imparted by the ileum, over which the omentum lay, just before it entered the cecum? The ileum at this point being more or less fixed, the propulsive effort is constantly in the same direction from the left to the right, and the direction of the twists in my specimen of omentum are the same as would be expected if this was the case. Now, the twists once started and the resulting inflammation in the omentum commenced, the increasing length of the inflamed veins would themselves tend to wrap around a line between two more or less fixed points. The swelling, edema and extravasation of the blood-cells would for the same reason also augment the twisting force.

Now let us go a step further. Nearly every case of reported torsion of the omentum was accompanied by some form of inflammation, either within or without the abdomen, as in a hernia. Was the inflammation the cause or the result of the torsion? The structure of the omentum has for its chief framework its vessels, long thin arteries that come off for the most part from the gastro-epiploic arteries at the great curvature of the stomach, and run downward to the lower border of the omentum, and then turn upward in the posterior layer to anastomose with the arteries of the colon. The veins are about the same. Enlarged full veins have been characteristic of the specimens of torsion of the omentum. Now

the veins increase in size and length more rapidly in inflammation than the arteries, and would have a natural tendency to wrap around the latter and carry the loose mesothelium, and may perhaps be the chief factor in the production of torsion of the omentum; and I think also that we can add to this the force of the pulsations of the omental arteries.

I have suggested the foregoing, for I have long felt that the torsion of the omentum was not due wholly to exterior forces, and I believe that we must look for a cause within the omentum itself that would produce or is capable of producing torsion, because in pure hernial torsion of the omentum where the omentum is in contact with only the smooth wall of the sac it seems as if the possibility of an exterior force is negative.

112 West One Hundred and Nineteenth Street.

A DUODENAL TUBE

SECOND COMMUNICATION

M. GROSS, M.D.
NEW YORK

In my first communication on this subject¹ my principal object was to show that a method had been found to pass over the pyloric ring and to reach the upper part of the small intestine. The present communication serves to present the method in greater detail.

The stomach-tube devised by me in 1894 requires only a few special alterations to answer the purpose of obtaining the duodenal contents.²

Apparatus.—This is shown in Figure 1.—A perforated round metal ball, twice the size of a pea and enveloped by the tube, serves as the end-piece. Attached to the end-piece is a thin rubber tube, not easily collapsible, about 0.2 cm. in diameter 125 cm. in length and graded in divisions of 10 cm. To this is attached a glass receptacle, which is connected by a length of

Fig. 1.—Duodenal tube

tubing with a mouth-piece to be used by the operator if he aspirates himself, as I prefer doing; otherwise this part can be replaced by an aspirating bulb.

Anatomy.—Figure 2 speaks for itself, but a few facts concerning the position and size of the duodenum will be in order. The duodenum is the shortest and widest portion of the small intestine and is situated deeper than all the other parts of the intestinal canal. It is not quite 20 cm. long, the ascending portion measuring 4

1. Gross, M.: New York Med. Jour., Jan. 8, 1910.

2. Einhorn (Med. Rec., Jan. 15, 1910) uses a similar tube arranged on the same principle, which he calls a duodenal pump. He, however, employs a somewhat different method of inserting it.