various factors which I will not go into, to me it is fairly evident that this man is suffering from an ordinary form of hysteria, and I am not inclined to think that it is other than that. Perhaps you will say that that does not explain all of his symptoms. Even so, I cannot account for this condition except as coming under the general classification of ordinary hysteria.

DR. HAROLD N. MOYER, Chicago: This appears to be a new find, or is at least a very rare symptom-grouping. With such exceptional findings a certain degree of modesty in suggesting a diagnosis is becoming. There are certain salient symptoms which are not explainable under any theory that I know of regarding the association of symptoms: First, the sweating; second, the presence of six million red cells; third, a marked eosinophilia; and fourth, a persistently elevated temperature. These make a symptom-grouping that I have never seen associated. The case presents a new combination of symptoms, and possibly a new disorder.

DR. GEORGE A. MOLEEN, Denver: I have been very much interested in this case, but from the start it appeared to me on a functional basis. Someone has said that nothing could be present in hysteria but what could be produced by suggestion. I would like to ask if suggestion has been used in this case as a means of differential diagnosis. It has come within the experience of men who have used hypnotic suggestion that it is quite easy to produce either localized or general sweating to influence the temperature and produce other changes of equally remarkable natures. However, that would not account for the blood changes which occur in this case. The absence, however, of organic symptoms, or symptoms of tissue change, and the fact that the functional disarrangement which would produce or lead to the sweating can be reproduced by suggestion, lead me to believe that it is on a functional basis.

DR. F. E. MURPHY, Kansas City, Mo.: May I ask when this estimate of eosinophils was made?

DR. HENRY HERBERT: Last year.

DR. F. E. MURPHY: Fourteen per cent. of eosinophils is certainly a symptom that necessitates some consideration. It would be interesting to know the percentage of eosinophils at this time. Possibly a year ago there was an intestinal parasite or something of that sort giving eosinophils. It would be interesting to know at this particular time how much the symptom observed a year ago is evidenced. There is one feature I observed, and that is the condition of this boy's vascular system. Several of us palpated his blood-vessels, and there are certainly marked changes in them. I am disposed to believe that there is something more than functional trouble in this instance.

DR. HENRY HERBERT, Los Angeles, Cal.: The temperature was taken regularly and always by the mouth for a period of from two to five minutes. Suggestion has not been used in this case. In regard to the eosinophils, that suggestion of reexamining the blood of the patient is a very good one. I am at a loss to say what the diagnosis is. One interesting feature leads me to believe that it is a lesion of the third sympathetic ganglia: first, the intense pain on pressure; then, the accepted rule, or accepted experiment or experience, that irritation of the sympathetic ganglia, especially of the third on one side shows a unilateral sweat, whereas the paralysis of that center shows anhidrosis. Now, we have general sweating on both sides, pain on both sides on pressure over the third sympathetic ganglia. At the same time, over that ganglia in that approximate region, or the region which is supplied by the approximate nerves, there is anhidrosis—no sweating at all. That would indicate a general irritation of the sweating center and a paralysis of those nerves supplying that area—the suprascapular region. If there is any process, fibrosis or chronic inflammation, to furnish the ganglia with some pressure, that would account for that anhidrotic area, and at the same time irritation of the entire sympathetic ganglia. The symptom-grouping, as one of the speakers mentioned, with the eosinophils, speaks against the supposition of a functional disorder or hysteria. I think that symptom is based on a permanent disorder, probably a disorder of the nervous system.

HOW OUR THERAPEUTICS CAN BE IMPROVED*

BOARDMAN REED, M.D.
LOS ANGELES

Tremendous advances have been made in therapeutics during the past two decades, advances which it is needless to enumerate to educated medical men. But there remains yet room for improvement. There are still failures in treatment, many of which could be avoided by a more perfect mastery of all the conditions which confront those charged with the high duty of endeavoring to heal the sick. This can be attained only by raising the standard of the doctorate to a yet higher plane, so as to assure for every practitioner a better knowledge of all that pertains properly to the science and art of healing and the auxiliary medical sciences, together with clinical experience under instruction. Such a training with industry and patience would assure an accurate diagnosis in every case in which such a diagnosis is practicable, followed by the best possible treatment.

Improvement must result from a more thorough study of (1) man and his organs and functions in both health and disease, including especially diagnosis; (2) a more intimate acquaintance with all the reliable remedies, medicinal, mechanical and psychologic from whatever source, a knowledge which shall recognize the fact that the more active ones seem to be capable of exerting either of two opposite actions according to the dose.

A very large proportion of the failures in treatment, whether made by regulars, sectarians, quacks, counter-prescribers or self-dosers, is due to a wrong diagnosis. No method of treatment is likely to prove helpful when prescribed in the wrong case; very frequently then it does harm. Innumerable patients die because the diagnosis is made too late, the medical attendant not having obtained needed counsel in time. Other failures and often much mischief result from continuing too long and in large dose a remedy which may have done good at first. Sometimes the failure is in consequence of too small a dose having been prescribed. I have seen cases of hyperchlorhydria and ulcer of the stomach or duodenum aggravated by 10- or 15-gr. doses of sodium bicarbonate and yet afterward promptly relieved and finally cured with the help of teaspoonful doses of the same drug well diluted and combined, when necessary, with 30 to 40 gr. of bismuth.

On the other hand, while drop doses of the wine of ipecac will relieve some forms of nausea, and severe vomiting often yields to drop doses of Fowler's solution, *à la* Ringer, and a threatened miscarriage is stopped by the tincture of cimicifuga every hour in the same small dose, as I frequently proved in my general-practice days, any one who would try these medicines in the usual doses in such cases would surely aggravate them.

Evidently, then, it would be a gain to our therapeutics if medical students were plainly taught that tissue-disturbing drugs are capable of producing opposite effects according to the dose. This was demonstrated to my own satisfaction twenty-three years ago by a careful study of the elaborate reports of numerous experiments conducted by eminent pharmacologists in England and Germany especially. My conclusions, though questioned, have never been refuted. They were published first in the *Practitioner* (London), and after-

* Read in the Section on Pharmacology and Therapeutics of the American Medical Association, at the Sixty-Second Annual Session, held at Los Angeles, June, 1911.

ward in this country. In a large experience since 1888, they have been amply confirmed. The matter was summed as follows:

1. A relatively small dose always at first stimulates the parts (nerve-centers, nerves, glands, muscles, membranes or other tissues) which the drug specially affects.
2. A sufficiently large dose (pushed far enough) always finally paralyzes the parts which the drug specially affects.
3. A moderate or medium dose may first stimulate and afterward depress, finally paralyzing if repeated often enough.

Lauder Brunton was quoted in the same paper as having stated in his "Pharmacology" that "this opposite action of large and small doses seems to be the basis of truth on which the doctrine of homeopathy was founded," following which he proceeded to criticize the fallacies of Hahnemann.

But after all, the most important point in therapeutics is to have a correct, an exact and also an early diagnosis. Thousands of persons are to-day taking hydrochloric acid and pepsin or some other exciting digestive aid for that vague entity, "stomach trouble," when the proper examination would show hyperchlorhydria or ulcer which could only be made worse by any form of acid. Nearly as many more are being dosed with alkalies when they need a contrary line of medication. Still others, suffering from a ptosis of one or more of the abdominal viscera—and their name is legion, especially among the women—are receiving various lines of treatment from drugs to osteopathy and from Christian Science to a four-weeks' fast with no possible benefit, when most of them would be relieved at once by having their displaced organs replaced and held up thereafter by either an operation, or, as I have seen in hundreds of cases, by a properly fitted abdominal supporter.

Instances could be multiplied indefinitely of failures in cases in which the diagnosis was in doubt, and yet, no consultation was had for economic or other reasons. Being no longer an aspirant for practice, I can afford to speak plainly and condemn strongly any guessing or prolonged dilly-dallying about the nature of an obscure illness whenever a consultation would probably clear up the doubt. When in such a case the patient or family refuse a consultation, the attendant would be justified in retiring from it. When, on the other hand, the family ask for a consultation and the medical attendant objects, he could have no cause for complaint if asked to make way for a broader-minded man.

Probably in no other feature is the usual current therapeutics more open to improvement than in the treatment of constipation. No affection is more common among civilized people and none is more generally treated ineffectively. The resort to purgative drugs for sluggish bowels is well-nigh universal. Patients rarely seek medical advice for the trouble early but dose themselves with laxatives till it has become chronic and aggravated. Even then, they will not often accept any curative treatment which involves time and expense, such as a course of massage, electricity or mechanical vibration (especially intra-rectal), some one of which, or a combination of them, nearly always proves effective in skillful hands, with the indispensable laxative diet and regulated exercises, often even in chronic advanced cases. Doctors should warn such patients emphatically that neither purgatives nor the milder laxatives ever cure constipation, and that, when continued very long, they may render it incurable by any method.

The diet in most chronic affections is a factor of the very greatest consequence. A more thorough study of dietetics would immensely improve the therapeutics of the average practitioner of whatever school or system.

The modern exact methods in the diagnosis and treatment of gastro-enteric affections should receive more attention in our medical colleges. If every practitioner were required to master them, or not having done so, would seek help from specialists in that line, not only in his frankly stubborn abdominal cases, but also in countless other obscure affections, he would very often be helped sooner to find the key which would solve the difficulty and obviate a harmful prolonged dependence on hypnotics or other inappropriate drugs.

While iron and arsenic will generally cure anemia after the cause has been removed, quinin and arsenic nearly always cure true malaria, and mercury and the iodids have rarely failed to rid the system in time of syphilis, to say nothing of salvarsan, it would be well if everybody, including practitioners of all sorts and the laity, could be impressed with the fact that many of the most powerful remedies are capable of doing harm in large or long-continued doses and that their safe and useful employment is often limited to the tiding over of emergencies or crises or the relief of intolerable pain, and that the cure can usually be accomplished only by a removal of the cause. In a very large proportion of chronic cases at least, the cause is some violation of health laws which demands a radical change in the patient's habits. Much of the present ineffective therapeutics results from attempts to cure people of maladies while they are continuing the bad habits which caused them.

No one can set bounds to the possibilities which may develop from the many new lines of treatment recently introduced, the antitoxins, the internal secretions from various glands, etc. The results already achieved with them afford great hopes for the future of therapeutics.

After all, however, our best and most permanent cures will not result from drugs or other special lines of treatment alone, but from supplementing these by hygienic measures, by inducing our patients to live regular and temperate lives, eating and drinking simply and never to excess, chewing all food thoroughly, taking sufficient exercise and sleep, spending as much time as possible out in the open air and sunshine and, through abundant ventilation night and day, breathing pure air whenever practicable.

South Granada Avenue.

CARCINOMA OF THE CLITORIS *

CHARLES D. LOCKWOOD, M.D.
PASADENA, CAL.

Tumors of the clitoris are of infrequent occurrence, and malignant tumors are rare. The histopathologic study of these tumors has been neglected, and many of the reported cases are not accompanied by a pathologic report. I wish to report a case.

Patient.—Mrs. K., aged 70, mother of two children, healthy, except for her present complaint. For six years she has suffered from painful micturition and irritation about the vulva. For the past year the pain on urination has been almost intolerable.

* Read in the Section on Obstetrics and Gynecology of the American Medical Association, at the Sixty-Second Annual Session, held at Los Angeles, June, 1911.

* Because of lack of space the review of the literature is here omitted. It is given complete in the Transactions of the Section.