Both the cases here referred to were highly suggestive of tuberculosis in their external appearance, but both patients have made recoveries almost too prompt and complete to be consistent with that diagnosis. One of the cases had a distinct history of injury; in the other, there was none. One case became worse clinically during the first month of fixation; the other improved rapidly from the start. In one the temperature was consistent with a tuberculous infection during the height of the symptoms; no record of the temperature in the other case is available. My own theory (not yet wholly abandoned) was that we were dealing with a mild tuberculous focus which had reached the stage of healing and sclerosis before the accident broke down the wall of cicatrization and called attention to the foot.

Empire Building.

SPOROTRICHOSIS OF THE EYE *

WILLIAM H. WILDER, M.D.

AND

CLIFFORD P. McCULLOUGH, S.B.

CHICAGO

In 1898 Schenck¹ published a report of his studies of an organism that he had discovered in the pus from refractory subcutaneous abscesses of the arm. The patient had scratched the index-finger on a nail, causing a small abscess which was opened. In three weeks an ulcer developed on the back of the hand between the second and third metacarpophalangeal joints. The inflammation traveled up the arm and several abscesses formed.

From two different foci of the disease Schenck obtained and cultivated an organism the characteristics of which were similar to those of many of the yeasts and fungi. Dr. Erwin F. Smith of the U. S. Department of Agriculture tentatively assigned it to the genus *Sporotrichum*.

The pathogenicity of the organism was demonstrated by subcutaneous injections in dogs. Nodules developed containing yellowish-red gelatinous material, and cultures from this and the walls of the abscess gave the characteristic growth of the organism. Inoculation of a white mouse caused death, and the organism was recovered from the local lesion and also from the lungs and liver.

In 1900 Hektoen and Perkins² studied and reported a similar case of refractory subcutaneous abscesses from which an organism was isolated, identical in essential details with the one described by Schenck, which they called *Sporothrix schenckii*. In their case, in a boy of 5, the lesion developed on the index-finger after a blow with a hammer. The denuded area did not heal, the finger became swollen, and in a short time a well-defined, undermined ulcer the size of a dime developed, the floor of which was covered with grayish pus. The lymph-vessels

of the arm enlarged and numerous abscesses formed which healed with difficulty after opening.

Glycerin-agar tubes inoculated with material from the abscesses all developed pure cultures of the organism, which showed after a few days in an incubator at 37 C. (98.6 F.) grayish-white, elevated irregular colonies about the size of a pin-head. Hektoen obtained growth on agar, blood-serum, gelatin, potato, bouillon and vegetable infusions.

The first extra-American case was reported by deBeurmann and Raymond in 1903; since then many cases have been reported, the lesions affecting the subcutaneous tissues, lungs, muscles, tendons and also the mucous membranes, including the conjunctiva, pharynx and larynx. DeBeurmann and Gougerot, with their associates, have called attention to various morphologic and cultural differences in the sporothrix, many of which have since proved to be inconstant.

The following case observed by us, of infection of the eye with sporothrix, is, we think, the second indubitable one of its kind in this country, the other having been reported by Gifford in 1910.

REPORT OF AUTHORS' CASE

History.—C. P. M., a student, had been working in the laboratory with cultures of various strains of sporothrix, and on several occasions small capillary pipets containing emulsion of the organism were broken at a distance of 8 or 10 inches from the face.

Fig. 1.—Ulceration on lower lid.

Fig. 2.—Enlarged follicle and small ulceration on upper lid.

On the evening of July 10, 1913, he noticed a soreness of both eyes, together with photophobia and a sensation as if a foreign body was under the lids. The following morning the lids were slightly swollen, the pain was increased and the surrounding lymph-glands were quite tender on pressure.

July 11, the general condition was good; there was no fever. Leukocyte-count was 9,000. The pain, swelling of the eyelids and photophobia had increased. The conjunctiva of the eyelids of both eyes was reddened and so swollen that the fornix rolled out in a mass when the lower lids were everted. In addition there were present on the palpebral conjunctiva and also on that of the fornices several grayish-yellow, slightly elevated spots varying in size from 0.5 to 3 mm. in diameter from some of which the covering epithelium had been cast off so that they seemed like small ulcers. Numerous follicles presented in other portions of the conjunctiva. Secretion was rather scanty, and an examination of a smear made from gentle scraping over the spots and the conjunctiva, stained with alkaline methyl blue, showed no organisms, but numerous pus-cells. The next day scrapings were taken from the shallow ulcers and aerobic and anaerobic cultures were made. Potassium iodid, 10 grains

* From the Memorial Institute for Infectious Diseases.

1. Schenck: Refractory Subcutaneous Abscesses Caused by a Fungus Possibly Related to the Sporotrichia, Bull. Johns Hopkins Hosp., 1898-1899, p. 286.

2. Hektoen, Ludvig, and Perkins, C. F.: Refractory Subcutaneous Abscesses Caused by Sporothrix Schenckii—A New Pathogenic Fungus, Jour. Exper. Med., Oct. 1, 1900, p. 77.

three times daily, was given. The eyes were washed out several times daily with a solution of oxycyanid of mercury, 1:4,000.

July 16, the condition had not improved; there was swelling of the preauricular glands, which were painful on pressure. Because of discomfort from irrigation, the strength of the solution of oxycyanid was reduced to 1:6,000. No growth was obtained in the tubes, which were placed at room temperature (from 18 to 20 C., or 64.4 to 68 F.).

July 17, the condition of the lids was no better; more yellowish spots were seen forming in the conjunctiva. The patient seemed quite sick and had a temperature of 100.5 F. The preauricular, anterior cervical, submaxillary and postauricular glands were somewhat swollen and painful on pressure. The small ulcerated spots on the conjunctiva were treated with tincture of iodine.

July 18, the general condition was worse; the patient had headache and malaise; the temperature was 101; the leukocyte-count was 18,000.

Sporothrix was growing in the tubes, and spores were found in the scrapings from the conjunctiva.

July 19, the general condition was worse and the patient went to bed. Temperature was 102; leukocytes 14,600.

During the night a sudden pain occurred in the left knee on the internal side of the upper end of the tibia; in the morning the limb was very sore, and painful on pressure or motion, but without swelling or redness. On application of the ice-bag after twelve hours this pain disappeared. The swelling of the conjunctiva was somewhat less, and no new spots had appeared. The iodid was increased to 20 grains three times daily.

July 20, the temperature was 99 in the morning, and rose gradually to 101.4 in the afternoon. The ulcers in the conjunctiva were healing.

July 21, in the morning there was pain in the left elbow and wrist and the lower end of the right femur, which was very sharp especially on motion and pressure. The temperature was 100.6 in the afternoon.

July 22, the pains were still persistent. The temperature was 101 in the afternoon. The conjunctiva was much improved; the ulcers had healed.

July 23, in the morning all the pain had disappeared except in the left wrist. During the night there had been severe pains in the right knee around the head of the tibia, and also in the right ankle. The temperature was 100.3 in the afternoon, the leukocyte-count 11,200.

July 24, all pain had disappeared. The temperature was 99.4. The conjunctiva was still red and the fornices considerably swollen. From this time the temperature remained normal and also the leukocyte-count. The swelling of the glands disappeared, but the subconjunctival tissue of the fornices remained swollen for some days. Gradually the follicles and conjunctival swelling disappeared and in about two months the lids presented a normal appearance.

Microscopic Examination and Cultures.—July 12, microscopic examination of pus from an ulcerated nodule in the conjunctiva showed no organism. Aerobic and anaerobic cultures were made on blood-agar, Loeffler's serum and aerobic cultures on 2 per cent. dextrose agar, and incubated at 37 C.

July 16, there was no perceptible growth; the tubes were placed at room temperature (from 18 to 20 C.).

July 17, there were numerous small, white, rather indistinct colonies on the dextrose-agar, composed of long branching filaments; there were no spores.

July 18, the colonies were typical of sporothrix, each being distinct, with a center rising in ridge formation like the peak of a mountain. Microscopically there was an abundance of long filaments and round or oval spores; the latter were not only in the filaments but also free. The organisms stained with the ordinary dyes and retained Gram stain.

At this time Gram-positive, oval bodies were seen in smears of pus from the eye; these resembled sporothrix but were found only singly or in pairs, no definite clumps being observed.

A few days later colonies also appeared on the other aerobic cultures, but the anaerobic cultures remained sterile.

Agglutinative and Opsonic Tests.—At no time during the infection or convalescence was eosinophilia detected.

Blood-serum was collected on different days, heated for thirty minutes at 56 C. (132.8 F.), and then placed on ice. Serum from a normal person and from the patient before infection, treated in the same manner, served as a means of control.

From a culture of sporothrix isolated from the eye, a suspension of spores was made, and the agglutinative reaction of the serum determined microscopically. The suspension of spores was made in sterile salt solution, and spontaneous clumping did not occur in hanging drops incubated for one hour at 37 C. Increasing dilutions of each specimen of serum was made and mixed with equal parts of spore suspension. Hanging drops of each of these dilutions were then incubated for one hour at 37 C., and examined microscopically to determine the highest dilution in which agglutination occurred. These tests gave the results recorded in Table 1.

TABLE 1.—AGGLUTINATION OF SPOROTRICHAL SPORES BY PATIENT'S SERUM (NORMAL=0)

Date	Agglutination at Dilution of:
6/20	0
7/ 3	0
7/18	1:16
7/22	1:16
7/23	1:16
7/24	1:16
7/25	1:24
7/29	1:32
8/ 2	1:32
8/ 6	1:64
8/10	1:64
8/14	1:64
8/18	1:64
8/26	1:32

The same samples of serum were used to determine the point of opsonic extinction. Increasing dilutions of serum were made by means of salt solution. Human leukocytes were obtained by collecting blood-agar from a normal person in sterile sodium citrate solution, washed and suspended in sterile salt solution. A suspension of spores from the organism isolated from the eye was made in sterile salt solution. To each dilution of the various serums was added an equal quantity of the suspension of spores and the suspension of leukocytes, the mixture incubated at 37 C. for thirty minutes, and smears made and stained with carbol-thionin. The number of leukocytes engaged in phagocytosis was now determined in each case, 100 white cells being counted. A serum-free mixture of leukocytes, suspension and salt solution treated in the same manner served as a means of control, the dilution of serum in which no more cells engaged in phagocytosis than in the serum-free mixture being called the point of opsonic extinction (Table 2).

TABLE 2.—OPSONIFICATION OF SPORES OF SPOROTHRIX BY PATIENT'S SERUM (NORMAL) = 1:12

Date	Opsonic Extinction at Dilution of:
6/20	1:12
7/ 3	1:12
7/18	1:48
7/22	1:24
7/23	1:96
7/24	1:192
7/25	1:192
7/29	1:96
8/ 2	1:96
8/ 6	1:96
8/10	1:192
8/14	1:192
8/18	1:192
8/26	1:96

As the patient had worked with five strains of sporothrix, experiments were made to determine if possible which strain caused the eye infection. Four of these strains were obtained from Dr. D. J. Davis.

1. A culture which had been obtained from Dr. Gougerot and supposedly typical of the sporothrix called "sporothrix de Beurmanni."

2. A culture isolated by Dr. Heekten in 1898 from a case of typical sporotrichosis.

3. A culture isolated by Dr. D. J. Davis from a case of sporotrichosis.

4. A culture isolated by Dr. K. F. Meyers from lesions on horses suffering with lymphangitis.

5. A strain isolated by Dr. Benjamin F. Davis from subcutaneous sporotrichosis on the hand and arm of a woman.

Comparison of the organism isolated from the eye in this case, with the other strains by ordinary cultural methods, in smears, and in hanging drop preparations, showed no differences, and immunologic tests gave the same reactions with the various strains, as is shown in the summary (Table 3).

TABLE 3.—AGGLUTINATION TESTS WITH THE PATIENT'S HEATED SERUM AND THE DIFFERENT STRAINS OF SPOROTHRIX

Date	Strain of Sporothrix					Our Case
	A	B	C	D	E	F
7/3	0	0	0	0	0	0
7/18	1:16	1:16	1:12	1:16	1:16	1:16
8/6	1:64	1:64	1:64	1:48	1:64	1:64
8/10	1:32	1:64	1:64	1:64	1:32	1:64
8/26	1:32	1:32	1:24	1:32	1:32	1:32

These results are analogous to those obtained by Dr. D. J. Davis³ with the serum of immunized animals, and indicate that there is no essential difference in the agglutination of various strains of sporothrix. Experiments were then made with the same serum to determine also the point of opsonic extinction of the various strains.

TABLE 4.—OPSONIFICATION OF VARIOUS STRAINS OF SPOROTHRIX WITH PATIENT'S SERUM

Date	Point of Opsonic Extinction with Strains					Our Case
	A	B	C	D	E	F
7/3	1:12	1:12	1:12	1:12	1:12	1:12
7/18	1:48	1:24	1:48	1:48	1:24	1:48
8/6	1:96	1:96	1:96	1:96	1:48	1:96
8/10	1:96	1:192	1:192	1:192	1:192	1:192

Tests were also made with both heated and unheated serums of normal persons, of a woman with sporotrichosis of the right hand and arm and of rabbits immunized with suspension of sporothrix spores. In all of these it was impossible to differentiate in any degree the various strains of the organism by means of opsonic extinction.

PREVIOUSLY REPORTED CASES

The number of reported cases of infection of the eye or its adnexa with sporothrix is not large, and a brief review of the reports of them is instructive in this connection. The first case of sporotrichosis of the eye was reported by Danlos and Blanc⁴ in 1907.

CASE 1.—A man, aged 63, had a granulating ulcer on the lower lid near the outer angle. It began as a small white spot which suppurated and formed an ulcer that gradually increased in size. The lower lid was congested and swollen, especially at the outer angle, where the conjunctiva was reddened and edematous. The lesion, 1.5 cm. in diameter, was covered with red and firm granulations, and at several points a little pus exuded. The preauricular glands were normal, but a slightly enlarged gland was noticed under the maxilla. For several years previous to this ulcer a swelling had existed on the right wrist. Cultures from the lid gave typical growths of sporothrix. Local treatment with iodine was of no avail; after four months the ulcer was excised.

CASE 2.—The first undoubted primary case of sporotrichosis of the eye was reported in 1908 by Morax and Carlotti.⁵ It occurred in a man aged 70 who had a lesion on the left upper eyelid for several weeks. The entire left upper lid and the skin for several millimeters over the eyebrow was red and swollen; a curved line of small yellowish projections

which had appeared like small intracutaneous or subcutaneous abscesses extended down from the inner end of the eyebrow to the outer commissure of the eye. In addition a number of shallow ulcerations, covered by an abundant yellowish exudate, were present along the free border of the lid. These lesions were not painful and the lid was only slightly tender on pressure. No other part of the eye seemed to be affected. The large lymphatic trunk extending from the external commissure to the preauricular gland was swollen and hard, and the overlying skin inflamed. At one point a soft yellowish enlargement indicated a small abscess. In a general manner the trunk took an S-shaped curve through the enlarged preauricular, angulomaxillary and submaxillary glands. The preauricular gland was tender and movable and about the size of an almond; the other glands were not so large. This swelling began one week after the onset of the lesion. Microscopic examination of the pus from the abscesses on the lid was negative, but cultures gave growths of sporothrix three days after inoculation. The patient was placed on 2 gm. of potassium iodid a day; in twenty days the ulcers were healed and in a little more than two months all evidence of the trouble had disappeared.

CASE 3.—In the case reported in 1909 by Thibierge and Gastinet,⁶ a man aged 30 developed a small granuloma on the left external palpebral commissure; later a generalization of these lesions appeared over the entire body. By extension from the original lesion a crescentic ulceration involved most of the lower lid as well as the bridge of the nose. The surface was covered more or less with crusts which on removal disclosed a deeply ulcerated area. The free border of the lid was infiltrated, and swollen, with some degree of ectropion, but not ulcerated. On the upper lid were two small reddish-brown nodules. The palpebral conjunctiva was reddish throughout. Cultures from the exudate and scrapings of the ulcers gave growths of sporothrix. Complete recovery followed treatment with potassium iodid.

CASE 4.—Burnier⁷ described in 1909 a case in a boy, aged 16, with several subcutaneous nodules on the hand, some of which ulcerated, leaving sinuses discharging pus. At the time of ulceration, reddening and a slight swelling of the eyelids were noticed. There were three hard and tender nodules, split-pea in size near the external canthus of the right lower lid; also four or five smaller nodules in the inferior cul-de-sac of the eye. The conjunctiva was not ulcerated, nor were the glands involved. Local cauterization with silver nitrate and galvanic needle followed by healing in about two months; the nodules on the hand resisted treatment. Inoculation of tubes with pus obtained from incision of the nodules demonstrated sporothrix. After one month's treatment with potassium iodid the nodules disappeared.

CASE 5.—A generalized sporotrichosis in a man aged 65, was reported in 1909 by deBeurmann and Gougerot.⁸ One morning the patient noticed that one of his eyes was congested, but without pain or tenderness on pressure. This condition persisted for two months, when the periorbital region became tender; photophobia developed and the cornea became dim and later ulcerated. The ulcerating keratitis was followed by hypopyon, anterior staphyloma, perforation of the cornea, escape of the vitreous and loss of the eye, all in a few days. No cultures seem to have been made from the eyes, but sporothrix was recovered from other sources. This seems to have been a metastatic panophthalmitis from general sporothrix infection.

CASE 6.—Morax⁹ in 1909 reported sporotrichosis of the eye in a woman aged 38. Small yellowish-white nodules appeared in rapid succession in the right semilunar fold, ulcerating later and producing pedunculated lesions. A similar lesion appeared

3. Davis, D. J.: Interagglutination Experiments with Various Strains of Sporothrix, Jour. Infect. Dis., 1913, xii, 140.

4. Danlos and Blanc: Un cas de sporotrichose palpebrale, Bull. et mém. Soc. méd. d. hôp. de Paris, 1907, xxiv, 1450.

5. Morax and Carlotti: La sporotrichose palpébrale, Ann. d'ocul., 1908, cxxxix, 418.

6. Thibierge and Gastinet: Trois cas de sporotrichose dermo-hypodermique dont un avec lésions du pharynx, du larynx et du tibia, Bull. et mém. Soc. méd. d. hôp. de Paris, 1909, xxvi, 537.

7. Burnier: Un cas de sporotrichose gommeuse hypodermique ulcéreuse disséminée, Ann. d'ocul., 1909, cxli, 344.

8. De Beurmann and Gougerot: Sporotrichose cachectisante mortelle, Bull. et mém. Soc. méd. d. hôp. de Paris, 1909, xxvi, 1046.

9. Morax: La sporotrichose de l'appareil visuel, Ann. d'ocul., 1909, cxli, 321.

in the lower retrotarsal fold and another in the upper. There was considerable induration in both lids with an increase in the secretion. The caruncular region was injected, while the bulbar conjunctiva and the cornea appeared normal. The cervical and preauricular glands became enlarged on the second day of the disease, and in a few days were quite tender on pressure. Under surgical treatment and local application of silver the symptoms became worse. As cultures gave growths of sporothrix, treatment with potassium iodid was instituted, and healing took place. The serum of the patient caused agglutination of sporothrix on the thirty-second day of the disease, clumping occurring almost at once in dilutions of 1:10, and in from thirty to sixty minutes in dilutions of 1:200. The blood-count was 65 per cent. polynuclears, 18.5 per cent. mononuclears, 13.5 per cent. lymphocytes and 3 per cent. eosinophils.

CASE 7.—A. Fava¹⁰ recorded in 1909 an infection of his own eye which is remarkable in that the mode of infection is known. While injecting an emulsion of sporothrix into a rabbit, droplets accidentally infected the eye. Half an hour later he washed his eyes with a 1:4,000 solution of oxy-cyanid of mercury. No symptoms appeared until eleven days later, when he noticed a sensation as of a foreign body in the outer corner of the lower lid. On examination a small yellow spot was seen on the lower tarsal conjunctiva, and the eye was congested and slightly tender on pressure. The following day the tenderness increased and there was a slight edema of the lids; the preauricular and angulomaxillary glands were enlarged and tender. Several other nodules also appeared on the lower tarsus, one 3 mm. in diameter and several smaller ones on the upper. Three days later a small nodule appeared on the outer surface of the upper lid. All the nodules ulcerated and developed granulations like those following a broken chalazion. The lids were swollen and the whole conjunctiva congested, and the secretion increased. Cultures from the ulcerated nodules gave sporothrix, and microscopic examination of the secretion demonstrated an abundance of eosinophils and some sporotrichal forms. On the fifteenth day of the disease there were 6,000 leukocytes with 8 per cent. of eosinophils. Recovery occurred in about two and one-half months of treatment with potassium iodid, from 2 to 8 gm. a day.

CASE 8.—Another case of generalized sporotrichosis of the skin and subcutaneous tissues in which there were symptoms of eye involvement is reported by Jeanselme and Chevalier.¹¹ Seven weeks after the appearance of the nodules on the body the conjunctiva of the right eye became congested. The right pupil was obstructed by an exudate and could not be dilated with atropin. A few days later the iris became discolored and vision greatly decreased. Simultaneously there appeared an ulcerated nodule in the mouth. No cultures were made from the eyes, but sporothrix was obtained from lesions elsewhere.

CASE 9.—Morax and Cruchaudau¹² describe a case in a woman aged 22, who first noticed a slight redness of the left eye accompanied by some sense of discomfort and increased lacrimation. The lower lid was swollen and on the tarsal conjunctiva was a yellow ulcerating nodule, from which sporothrix was isolated in culture. A few days previous to the inoculations the serum agglutinated sporothrix at a dilution of 1:50. Tender and enlarged preauricular glands accompanied the eye lesion. After one month's treatment with potassium iodid all the symptoms disappeared.

CASE 10.—A case in which the eyelids were affected by direct extension is reported by Velter.¹³ A man aged 52 had a small abscess on the right lower lid which had existed for three weeks, causing considerable pain. The lid was swollen, its skin smooth and red; incision released a large amount of

pus. The surrounding tissues were indurated. The eyeball appeared normal, but the bone at the outer and inferior wall of the orbit was eroded. Cultures from the lid showed sporothrix. A similar nodule appeared on the upper lid, but at no time was there any involvement of the conjunctiva or marked glandular enlargement. Potassium iodid gave good results.

CASE 11.—Jeanselme and Poulard¹⁴ describe the case of a man aged 46, who had an amputation of the leg for a tumor of the bone, followed several years later by a generalized eruption of small red and hard nodules on the face and body. One of these nodules occurred on the outer surface of the left lower eyelid just below the margin. The anterior part of the eyeball was injected, and the pupil narrowed, irregular and obstructed by an exudate, but there was no infiltration of the cornea. Posterior synechia and discoloration of the iris were present, and there was pain, photophobia and decreased vision. The generalized eruption proved to be sporotrichosis; the eye lesion was regarded as sporotrichal because it also was healed, under the use of potassium iodid.

CASE 12.—The only case reported in the United States is by Gifford.¹⁵ A woman, aged 18, complained that the left eye was congested and slightly uncomfortable, the condition having lasted for about two months without any marked change. When seen the right eye appeared normal. The left eye was inflamed and watery, the tarsal conjunctiva and retrotarsal folds being slightly congested. On the ocular conjunctiva a somewhat crescentic thickening extended from a little above the equator at the inner side of the cornea to a point $\frac{1}{4}$ inch above the horizontal meridian at its outer side. This thickening was $\frac{1}{4}$ inch broad and smooth at its widest part but nodulated with lumps from $\frac{1}{16}$ to $\frac{3}{32}$ inch in diameter, at the extremities. One of these nodules was yellow and contained a semifluid substance, from which sporothrix was isolated by culture. Treatment by potassium iodid seemed to be effective, but the patient disappeared from observation before complete recovery.

CASE 13.—Morax¹⁶ reported the first case of infection of the lacrimal sac by sporothrix. A man, aged 40, complained of watering and a feeling of weakness in his left eye, followed in a short time by swelling and inflammation of the lacrimal region and the appearance of pus at the level of the internal commissure. Three weeks after the onset, suppurative dacryocystitis with prelacrimal abscesses developed. The lacrimal sac was swollen and on gentle pressure pus exuded from both superior and inferior puncta. The skin over the region was somewhat infiltrated. After incision and withdrawal of yellow purulent material, a sound could be passed into the nose.

CASE 14.—In the case of Chaillous,¹⁷ the patient complained of matting of the eyelids after sleeping, this condition lasting for eighteen days before examination. The right upper lid was swollen, reddened and edematous. On the tarsal conjunctiva were numerous yellow nodules, from pin-head to split-pea in size, irregular in form and slightly ulcerated. The bulbar conjunctiva was less congested. A preauricular and a superficial cervical gland became quite tender and enlarged. Cultures from the ulcerated nodules showed sporothrix.

CASE 15.—LaPersonne¹⁸ describes a case in a woman aged 64 who, during the course of numerous subcutaneous and periosteal lesions, developed a violent iridocyclitis of the left eye with deposits on the posterior surface of the cornea and a nodule on the iris. The anterior chamber of the eye became involved, accompanied by pain in the ciliary region. The eye softened and vision was almost lost. Punctures of the anterior chamber liberated a yellowish albuminous fluid from which no diagnosis could be made. Two months later another similar lesion appeared and perforated the coat of the eye-

14. Jeanselme and Poulard: Sporotrichose de l'iris, *Ann. d'ocul.*, 1910, cxliv, 65.

15. Gifford: Sporotrichosis of the Eyeball and Eyelids, *Ophth. Rec.*, 1910, xix, 573.

16. Morax: Sporotrichose primitive du sac lacrymal, *Ann. d'ocul.*, 1911, cxlv, 49.

17. Chaillous: Sporotrichose conjunctivale primitive, *Ann. d'ocul.*, 1911, cxlv, 47.

18. LaPersonne: Sporotrichose oculaire, *Presse méd.*, 1912, xx, 93.

10. Fava: Un cas de sporotrichose conjunctivale et palpébrale primitive, *Ann. d'ocul.*, 1909, cxli, 338.

11. Jeanselme and Chevalier: Sporotrichose à foyers multiples, *Bull. et mém. Soc. méd. d. hôp. de Paris*, 1910, xxix, 784.

12. Morax and Cruchaudau: Sporotrichose conjunctivale primitive, *Ann. d'ocul.*, 1910, cxliv, 69.

13. Velter: Un cas de sporotrichose orbito-palpébrale primitive, *Ann. d'ocul.*, 1910, cxliv, 65.

ball. A fungus was demonstrated among the debris of the iris and exudate, and a few days afterward sporothrix was obtained in culture. The patient bore potassium iodid poorly, and finally the eye was enucleated because of the pain.

CASE 16.—The last case described is by Morax¹⁹ in a man aged 37, who had nodular lesions on the skin of the thighs and a nodule on the outer surface of the lower lid, below the internal palpebral ligament. This nodule was violet-red, firm and easily movable without pain. It was about 0.7 cm. in diameter and projected 2 mm. above the surface, with the center slightly depressed like a lacrimal fistula. The conjunctiva appeared normal and there was no cutaneous reaction or congestion of the surrounding tissues. Exploration by means of a probe showed that there was no connection of the lesion with the eye. Cultures from this nodule gave sporothrix. Cure followed treatment with potassium iodid.

In experiments by Fava²⁰ and others, following the injection of a culture of sporothrix into various parts of the eye of a rabbit, lesions were produced resembling those of sporothrix infection of the human eye. The incubation period generally lasted from twelve to eighteen days. The organism was recovered from the eye lesion.

CLINICAL CONSIDERATION

From a study of the cases that have been recorded and of the one we have observed, we should say that infection of the conjunctiva by sporothrix causes a marked congestion of the membrane, particularly the palpebral portion, the fornix and the semilunar folds. The bulbar portion is not so frequently infected.

Numerous follicular prominences appear in the palpebral conjunctiva and in the fornix.

Small yellowish nodules, varying in size and shape, develop rapidly in the conjunctiva, and these may ulcerate. When opened, the contents of the nodules do not escape rapidly as from small abscesses, but seem to be of a gummy consistence. These little nodules developed so rapidly in the case we observed that on the second day new ones presented that had not been seen the day before.

Secretion is rather scanty and hardly sufficient to stick the lids together at night, but lacrimation is rather abundant. The eyelids are somewhat edematous and thickened, and palpation shows a well-marked induration of the subcutaneous tissue. Enlargement and tenderness of neighboring lymph-glands is also present.

Subjectively, there is a sensation of a foreign substance under the lids, and so much discomfort that use of the eyes is almost impossible. The discomfort comes on rapidly after the infection. The two cases of laboratory infection (that of Fava and the present one) and the absence of a history of trauma in the other cases, seem to show that the sporothrix is able to penetrate the normal conjunctiva. It also appears from the reports of cases that infection of the eye may be secondary to a generalized sporotrichosis. Probably in most of these cases the infection is ectogenous, but the suspicion that it may be endogenous is aroused by the case of LaPersonne, in which, after a violent iridocyclitis and perforation of the eyeball, sporothrix was obtained from the contents of the bulb.

That general symptoms may arise from a primary lesion of the eye is indicated in our case by the fever, leukocytosis and pains in the bones of the extremities.

DIAGNOSIS

Some of the clinical features of this infection are common to other conditions. Lymphadenopathy would be present with chancre of the conjunctiva, but in the initial lesion of syphilis it is very unusual to have such multiple erosions or ulcerations, and scrapings from such an ulcer would probably show the characteristic spirochete.

Tuberculosis of the conjunctiva would probably not be so rapid in its course, and it would be a week or more before the caseous tuberculous nodule would break down and form the ulcer, whereas in sporotrichosis the little ulcers develop in a few days.

Parinaud's conjunctivitis presents more points of similarity, and it is possible, as mentioned by Morax, that cases of sporotrichosis may have been mistaken for Parinaud's conjunctivitis.

In the latter, the vegetations on the conjunctiva are different from the follicles and the yellowish nodules of sporotrichosis. The adenopathy in Parinaud's conjunctivitis points to a severe infection, but all attempts to isolate an organism from the lesions have failed. Recently, however, Verhoeff has observed in such conditions an organism like leptothrix. On the other hand, the diagnosis of sporotrichosis is easy if scrapings from the nodules or ulcers are inoculated on appropriate mediums and left at from 18 to 20 C. for the organisms appear in from three to ten days. The presence of Gram-positive, spore-like bodies in a direct smear from the conjunctiva should suggest sporothrix.

We wish to acknowledge the kindness of Dr. L. Hektoen, director of the Memorial Institute for Infectious Diseases, for valuable suggestions offered in the study of this case.

122 South Michigan Boulevard.

CLINICAL REPORT OF FIVE HUNDRED AND SEVENTY-ONE CASES OF PULMONARY TUBERCULOSIS TREATED WITH TUBERCULIN *

LOUIS SHALET, M.D.

Attending Physician Tuberculosis Clinics, Department of Health
NEW YORK

Early in the spring of 1910, and following my presentation of a short paper on tuberculin at one of the meetings of the medical staff of the Otisville Sanatorium, of which I was a member, Dr. E. S. McSweeney, the physician then in charge, asked me to reorganize the tuberculin class among the patients.

After rather an extensive perusal of much of the literature on the subject, I was struck by the divergent views held by the different investigators on tuberculin therapy in general and the matter of dosage and reaction in particular. I then decided to lay aside all my preconceived notions on tuberculin and begin injections with a dose well below any possible reaction, such as 0.000,000,01 gm., and to double each previous dose at weekly intervals, using Koch's B. E. In the last year of this work, patients in the class were started on Koch's Old Tuberculin exclusively, and when they were standing well from 0.003 to 0.006 gm. at a dose, they were then changed to Koch's B. E., 0.000,1 gm., which was later progressively increased according to the table. The

19. Morax: Sporotrichose primitive des paupières simulant une fistule lacrymale, *Ann. d'ocul.*, 1913, cxlix, 183.

20. Fava: Sporotrichose expérimentale de l'appareil oculaire du lapin, *Ann. d'ocul.*, 1910, cxliv, 77.

* From the Department of Health Sanatorium for tuberculosis at Otisville, New York.