tion, and it is for the purpose of stimulating interest in the subject that this argument is presented. In the writer's experience all syphilitic bone and joint conditions improve more rapidly when the dose of potassium iodid is carried to the point of tolerance than when dependence is placed on a routine dose of moderate size, and it is suggested that this technic be used when medication is adopted.

To illustrate the effect of mixed treatment, the histories of two cases under observation on the service of Dr. H. L. Taylor, at the Hospital for Ruptured and Crippled, are appended with roentgenograms that confirm the diagnosis.

REPORT OF CASES

CASE 1.—The patient, Ida G., aged 8 years, was suffering pain and limped in July, 1914. Night cries were noted at that time. She was treated at two different clinics for tuberculosis of the right hip. The limb was placed in a plaster cast for a year and four months. The plaster cast was then removed for a month. Pain, restlessness at night, restriction of motion and spasm reappeared and another plaster spica was applied at the Hospital for Ruptured and Crippled, Dec. 1, 1915. This treatment was continued until March 21, 1917, when the patient came under my observation. The plaster cast was removed; the hip was found to be stiff and extremely sensitive. The teeth indicated congenital syphilis, and the child was therefore placed on mixed treatment and the spica fixation discontinued. Two weeks later the patient reported for examination. Pain and spasm had disappeared. There was free rotation and considerable flexion without discomfort. From that date there has never been any return of pain. Improvement has continued, and on Oct. 1, 1917, there was flexion of the thigh to 45 degrees inside of a right angle; free rotation; no limp, and the child was extremely active. The roentgen-ray findings were typical of Perthes' disease (Fig. 1).

CASE 2.—The patient, Elsie G., aged 11 years, complained, in February, 1916, of pain in the right leg; the patient limped, and tired easily. She came under observation, May 2, 1917. There was flexion of the thigh to a right angle and full extension, but rotation in extension was limited. Rotation in flexion was blocked and abduction was extremely limited. Pain and spasm occurred on attempt to force motions and there was a marked limp in walking. The teeth indicated congenital syphilis, and the patient was put on mixed treatment without any protection to the hip. A month later there was flexion to 45 degrees inside of a right angle and some rotation with the thigh flexed. There was no pain, and the child was much more active and felt better. July 18, 1917, there was almost normal flexion and free rotation with thigh flexed. There was no pain and no limp. The roentgenograms revealed typical Perthes' disease (Fig. 2).

In Case 2 the symptoms had existed a year and three months and disappeared under suitable medication in ten weeks.

576 Fifth Avenue.

Quadracentennial of Dodoneus.—The *Nederlandsch Tijdschrift voor Geneeskunde* of June 30 was devoted entirely to historical sketches of Rembertus Dodoneus, a great physician and botanist of the sixteenth century. His works include the "Cosmographia," "History of Plants," "Praxis Medica" and the "Cruydt-boeck," in which he contributed much to the materia medica of western and northern Europe which had been neglected by previous writers. He was offered the post of body physician to the king of Spain, as successor to Vesalius, but declined. Later he accepted a similar position with the emperor of Austria until he was called to the chair of clinical medicine at the University of Leyden.

SYSTEMATIZATION OF A SURGICAL SERVICE

EUGENE H. POOL, M.D.

AND

FREDERICK W. BANCROFT, M.D.

Attending Surgeon and Associate Attending Surgeon, respectively,
New York Hospital

NEW YORK

In modern industrial organization, the highest degree of efficiency is attainable only by intense systematization. In hospital management the interde-

Fig. 1.—Movements of nurses during an operation for appendicitis: N-1, head instrument nurse; N-2, sponge nurse; dotted lines indicate steps taken by nurses during the operation; lines without arrows indicate "round trips," those with arrows indicate "one way trips."

pendence of efficiency and systematization should be recognized. Yet, this conception has been slowly grasped, reluctantly accepted, and compromisingly adopted by the medical profession. Although a mechanical and inelastic system is not suggested for the care of the sick, the advantages of a systematic plan of service, so far as this is consistent with the development and application of medical skill, are striking.

Until a few years ago, hospital services were relatively small, diagnostic methods simple, the staffs

Fig. 2.—Steps taken by nurses during an operation for inguinal hernia. Similar in plan to Figure 1.

small, and the turnover slow. Consequently the details were simple, comparatively few, and readily kept in mind. The treatment of the individual patient was, in general, conscientious and satisfactory.

With the large service of the present day, however, the complicated methods of diagnosis, the large staffs and the rapid turnover, systematization has become essential. By means of it, the time and the energy of the staff are conserved; moreover, other things being equal, thoroughness in the study and treatment of each case is better insured, and oversights and mistakes are less in proportion to the degree of systematization.

The essentials for systematization are supervision by a permanent director and specialization in details by other members of the staff. Our service has been developed with these objects in view.

The surgical department of the New York Hospital consists of two independent divisions. They are similar in organization. We are to consider, however, only the second surgical division.

The staff of the division consists of an attending surgeon, senior associate surgeon, junior associate surgeon and assistant surgeon. Gradation is important in that it eliminates questions of authority and pre-rogative.

Fig. 3.—Marked reduction in movements and labor of nurses due to the introduction of simplified methods in the operating room. New and larger tables installed. Diagram was made during an operation for appendicitis.

The attending surgeon is in charge and is responsible continuously. The two associates, while equal in rank so far as hospital appointment is concerned, are graded for work on the division. When the attending surgeon is absent, the senior associate takes his place; the junior associate assumes the work of the senior, and the assistant acts as junior associate. Under this plan, the policy of the division is unvarying, because in the absence of the attending surgeon, the routine and regulations are enforced by the associate, who acts in his stead and not independently. A concerted and unchanging policy is thus possible. But such a plan demands the subordination of the individual to the welfare of the service; associates and assistants must serve the hospital as members of a team rather than as individuals.

Inspection rounds, followed by conference with free discussion, are held once a week, and a follow up or end-result system has been established. This routine furnishes the chief with an accurate knowledge of the work of the division, unifies the ideas of the staff, develops a harmonious spirit among them, and checks the immediate and late results.

The weekly inspection rounds are conducted by the attending surgeon, with the staff and student clerks. Practically every case is dressed, the treatment of the wounds is discussed, and permanent records of the condition of each patient and wound are made on the bedside charts. Following the rounds, there is a conference which is attended by the entire staff, including attending surgeons and interns. The conference embraces:

1. A review of the work of the past week, which includes: (a) operative results; (b) diagnostic errors, and (c) late results.
2. Report of standing staff committees (outpatient department, student clerks, follow up system, operating room technic, clinical records, etc.).

3. Reports on special subjects assigned by the chief to a staff member; or a review of the literature bearing on some unusual case on the service.

We will consider these topics in detail.

Operative Results.—Unfavorable results and unfavorable developments are analyzed: (1) deaths; (2) infections, and (3) other postoperative complications, such as hematomas and pneumonias. An effort is made to determine the cause of each unsatisfactory development. As advised by Codman, the poor results are tabulated into:

1. Errors due to lack of technical knowledge or skill (E-s).
2. Errors due to lack of surgical judgment (E-j).
3. Errors due to lack of diagnostic skill (E-d).
4. The patient's enfeebled condition (P-d).
5. The patient's unconquerable disease (P-d).
6. The patient's refusal of treatment (P-r).
7. The calamities of surgery, or those accidents and complications over which we have no control.

Diagnostic Record of the Week.—Errors of diagnosis (E-d) are referred to the operator for explanation, and a general discussion of the case follows. It is believed that such an analysis and discussion leads to greater care in arriving at a correct and exact diagnosis before operation, emphasizes the causes of diagnostic errors, and suggests means for avoiding them.

To obtain an exact diagnostic record, we employ the following method: The anesthetist, before the beginning of each operation, notes the anteoperative diagnosis on the anesthesia chart. The operator is entitled to record his diagnosis up to the time of making his initial incision; after that it cannot be changed. At the end of the operation, a revised diagnosis is recorded.

In cases in which the diagnosis at the close of the operation is still in doubt, as in certain tumors, and in some cases presenting abdominal symptoms in which no lesion has been found, final decision as to the diag-

Fig. 4.—Similar in plan to Figure 3. By the elimination of unnecessary steps, the work has been greatly simplified. Diagram was made during an operation for inguinal hernia.

nosis is deferred until it is determined by the pathologist's report or by the course of the case.

End-Results.¹—The late results in the cases which have been examined on the previous Sunday morning are analyzed. Those recorded as "fair" and "poor" are discussed, and, if necessary, the original histories of the cases are reviewed. The causes of failures are sought, and measures for avoiding similar failures are considered. The errors or failures are grouped according to the classification described under operative results.

1. The details of our end-result system have been described by Bancroft, F. W.: Bull. Johns Hopkins Hosp., 1916, 27, 201.

A brief comment on the end-result system is appropriate. The conscientious management of a surgical service demands not only a detailed knowledge of the immediate results of operative treatment, but also the ultimate or end-results. The operative policy of a surgical service is directly dependent on the study and analysis of the late results of operations. If it is

Fig. 5.—New supply table with foot pump. Elevation is used when patient is in the Trendelenburg position. Two tables are in use in the operating room, one each for the instrument nurse and the gauze supply nurse.

found that a certain procedure is not giving permanent relief, the members of the staff will be stimulated to devise means that will improve the results. For example: During the two years in which this system has been in use, it has been found that a certain number of patients operated on for chronic appendicitis have not been relieved of their complaint. Certain members of the staff are now making a careful analysis of all cases diagnosed as chronic appendicitis, with the object of determining wherein lies the fault. Such a study can be made effective only by careful analysis of the end-results of a large series of cases in conjunction with the histories, physical examinations, laboratory data, roentgen-ray records, and operative findings.

A follow up system, to be effective, must be so planned that all patients leaving the hospital are followed for at least one year, and selected patients for a longer period.

All patients on leaving the wards are requested to return for observation at the end of three months; if examination at that time indicates that there is no need for earlier observation, they are requested to report again at the end of nine months. We have been able to obtain the late results in approximately 90 per cent. of our cases.

We cannot emphasize too strongly the credit due to Codman of Boston for initiating the follow up system and bringing the profession to an appreciation of its importance; also to Corscaden of the Presbyterian Hospital, New York, for his extremely efficient method of putting the follow up system into effect. We employ his card system with slight modifications.

Attempts have been made to arrange the work in the wards in such a manner that the maximum atten-

tion may be given the patients with the minimum waste of effort. Complete rounds are made daily by one of the staff. Each member of the attending staff has definite days on which he makes morning rounds on all the cases on the division. Patients in the wards are considered as belonging to the division and not to individual members of the staff. In other words, if one of the attending staff refers a case to the service, the patient becomes a ward of the division. Thus, each patient has the advantage of study by the entire staff. No hardship is involved, because no fees are collected for ward patients, except workmen's compensation cases, and the fees for these are collected jointly for the needs of the division.

There is a rule of the division that the operator must examine all patients on the afternoon preceding operation. This, therefore, necessitates the visiting of the hospital every afternoon by one member of the staff.

Three difficulties are often encountered in large surgical services:

1. Delay in ordering and making appropriate laboratory and clinical examinations.
2. Delay in calling the attention of the attending surgeon to very ill patients.
3. Premature dismissal from bed and hospital of patients who have been operated on by the zealous house surgeon who wishes to make room for patients who are to undergo operation.

To overcome these difficulties the following methods have been adopted:

1. An associate has been placed in charge of each ward. It is his duty to visit his ward daily and to see all the patients who are to be operated on, order roentgen-ray and laboratory examinations as indi-

Fig. 6.—Same table as shown in Figure 5, which has been elevated for the Trendelenburg position.

cated, direct and expedite history taking and physical examinations, and note on the chart his diagnosis and suggestions as to treatment. He also supervises the postoperative therapy.

2. A day book is kept on the nurses' desk, with three columns for names: In the first column are noted those patients who have not been operated on;

in the second, patients not doing well or requiring investigation, and in the third column, the names of patients presumably ready for discharge. The attending surgeon can thus learn at a glance the salient facts.

3. Each patient before going home is examined by one of the associates, who writes a discharge note on the chart. This record includes a statement as to the general condition of the patient and the condition of the wound, and a brief summary of the postoperative course. The associate gives the patient a note to the family physician or to the dispensary. In this he describes the operation and convalescence, and suggests treatment. If he considers the patient unable to leave the hospital, he instructs the nurse to keep the patient until further notice. Consequently, it is seldom that patients leave the hospital too soon or in improper condition.

OPERATIVE TECHNIC

An effort has been made to simplify the work in the operating room. The properly trained operating staff should be a well-balanced machine; it is, however, difficult to attain this end. In the average hospital, each member of the attending staff has his individual ideas as to the arrangement of the operating room and of instruments, etc. It often becomes necessary to arrange the room and to plan certain details even of routine operations differently for each man. The lack of uniformity interferes with a fixed policy, leads to confusion, and imposes a greater burden on the helpers. All routine operations should be standardized: the arrangement of the room should be uniform; a given number of instruments should be on the tray, and the same suture materials and instruments should be used. As far as possible, the same incision and procedure should be employed, but unfortunately the vagaries of the human body and the uncertainties of diagnosis require occasional deviations. However, it is possible in most cases to follow a regular system.

The staff, at the conference, decides on the routine types of operation. Let us take inguinal hernia, as an example: The modified Bassini operation with transplantation of the cord has been adopted; all members of the division use this method; all use the same instruments, sutures and needles. If a member of the

staff disapproves the method or of any detail, he must present his criticisms and suggestions at a regular weekly conference.

The chief often assigns the study of a certain type of operation to a member of the staff, who presents his report at a subsequent conference. Discussions at these meetings are often very instructive.

We are making constant efforts to secure simplicity and a safe routine in the operating room. When the effort to systematize this service was begun two years ago, we felt that there was much waste of effort in the operating room work and that certain features of the work were glaringly inefficient. Let us illustrate with one detail: the steps taken by the nurses during an ordinary operation by reason of improper planning and thoughtless arrangement of the necessities for an operation. It was decided to make notes of the steps

taken by all concerned during an operation. To accomplish this, one of us (Bancroft) dictated to a stenographer the movements of the various nurses and physicians during a routine operation performed by the attending surgeon (Pool). Two operations, the repair of an inguinal hernia and an appendectomy for chronic appendicitis, were thus recorded.

We made a ground floor chart of the operating room, drawn to scale with the tables in place for the operations. On this, the movements of each nurse from the beginning to the end of the operations were outlined (Figs. 1 and 2). After analysis of the dictated records and the graphic charts, we found the main faults to be:

1. Lack of system.
2. Too many and too small tables for sterile supplies, extra instruments, etc.
3. Unnecessary energy wasted when the patient is in the Trendelenburg position by nurses stepping down from benches to obtain supplies from tables and stepping up again to reach the operative field or instrument tray.

To correct these faults, we explained to the operating room staff the significance of the charts, and impressed on them the importance of simplifying the work. By interesting the nursing staff, we have been able to cut down the unnecessary steps to an astonishing degree, as is evident by reference to Figures 3 and 4.

Three of the old tables were discarded and two new ones (Figs. 5 and 6) were substituted. These are much

Fig. 7.—Manual signal system for use during operations: 1 *a*, artery clamp; *b*, Allis clamp (intestinal forceps); *c*, artery clamp with wriggling of little finger; *d*, Kocher clamp, as artery clamp with wriggling of thumb; 2 *a*, Towel clip; *b*, Kelly clamp; *c*, as towel clip plus wriggling of thumb; 3, ligature; 4, sponge pads; 5, *a*, scalpel, with cutting motion; *b*, straight needle, with rotary motion; 6, *a*, thumb forceps; *b*, scissors, opening and closing fingers; 7, tissue forceps, one and two teeth; 8, tissue forceps, many teeth; 9, sharp retractors; 10, palm down: *a*, large right angle retractors, thumb and index finger; *b*, medium right angled retractors; use middle finger instead of index finger; *c*, small right angled retractors, use ring finger instead of index finger; 11, palm up: *a*, large string pad, thumb and index finger; *b*, medium string pad, thumb and middle finger; *c*, small string pad, thumb and ring finger; 12, *a*, McBurney, small; *b*, McBurney, large; use both index and middle fingers; 13, abdominal retractors; 14, *a*, aneurysm, rotary motion; *b*, cautery, squeezing motion; 15, palm down: *a*, large sponge stick, sponging motion; *b*, curved needle, rotary motion; *c*, small sponge stick, as 15 *a*, but with palm up; 16, towel, spreading motion.

larger than the old ones, and are provided with a foot pump so that they can be raised or lowered easily. Raising these tables to the proper level eliminates the necessity for the nurse to move from the bench in order to obtain extra supplies during pelvic operations. The stimulus to improve the technic of the routine operations keeps the staff on the alert and counteracts the monotony of the routine cases.

We have adopted a signal system (Fig. 7) for use during operations. Many others, notably Brewer, have adopted such a procedure more or less extensively, and we refer to it merely as a suggestion to those who have not had the idea called to their attention. The habit of signaling becomes automatic, and the appropriate instrument is quickly obtained. The method seems to add to the smoothness of the operation.

COOPERATION WITH DISPENSARY

An effort has been made to bring the dispensary and the hospital service into close association. The fourth man on the service is chief of the dispensary, and his privileges in the wards are dependent on his management of the dispensary. Each of the other three members of the staff is expected to visit the dispensary on a definite day each week; his visit is scheduled for a certain time, and unusual cases are reserved for his inspection and advice.

About once a fortnight, a member of the dispensary staff gives a clinic. Invitation cards are sent to the physicians of the neighborhood, and unusual cases or new methods of therapy are presented. This stimulates interest and prevents initiative being obliterated by routine.

The history of every case sent from the dispensary to the wards accompanies the patient and is attached to the ward history. On discharge, a note is written by the house surgeon summarizing the operation, pathologic findings and convalescence. The dispensary history is then returned to the outpatient department. In this manner, it is possible for the outpatient surgeon to follow his cases.

Ventilation of Shipholds After Fumigation.—Experiments by the officers of the Public Health Service in the fumigation of ships for the killing of rats suspected of being the source of plague have shown that sulphur dioxide and hydrocyanic acid gas are the best fumigants. The latter is most efficient and desirable because it takes less time, and the ship may be put into service sooner after fumigation. To shorten the time still further and render the ventilation of the ship's compartments more complete, experiments were conducted. Grubbs of the Public Health Service describes in *Public Health Reports*, Oct. 19, 1917, the methods employed. In India, according to Heiser, the gas is aspirated from the vessel through the pipes used to fill the holds with the gas. This method was not employed by Grubbs. Electric fans were placed in the vessels and operated by the electric plant belonging to the vessel. It was found that the time of ventilating the ships could be shortened, but there were objections on account of handling the wires connected with the fans. A gasoline-driven air propeller, designed to propel boats and sleds, was then employed. This was found to deliver about 22,700 cubic feet of air a minute. This fan was used for distributing the gas in the hold and also for ventilation. In its use for the latter purpose, it was attached to the hatch combing and the air introduced into the hold through muslin chutes. It was found that this machine in ten minutes would throw into the bottom of the hold an amount of air equal to the cubic capacity of the ship. This fan has been modified so that it will work horizontally, and is arranged on a frame so that it may be carried by two men. It weighs less than 100 pounds. It has been found expeditious and satisfactory.

SURGICAL TREATMENT OF CHRONIC ENTEROCOLONIC DIARRHEA*

SAMUEL C. GANT, M.D., LL.D.

Professor of Diseases of the Colon, Rectum and Anus, New York Post-Graduate Medical School and Hospital

NEW YORK

Chronic diarrhea can be temporarily relieved by rest, strict diet, antiseptic, astringent and antiperistaltic drugs and irrigation through the anus, but often recurs when the patient returns to business, eats normally and discontinues medication.

Chronic inflammation or ulceration of the intestine, however produced, is often difficult to cure except by surgical intervention. By operation and irrigation, I have permanently cured many diarrhea patients who had been previously treated for months or years by nonsurgical measures.

Up-to-date physicians and surgeons realize that removal or direct treatment of bowel lesions is essential to the successful handling of many types of chronic diarrhea; and I hold that the time has arrived for the profession to give "intestinal stasis" a rest and to devote its attention to the operative treatment of otherwise incurable chronic diarrhea.

The family physician must be taught to differentiate between different types of frequent movements that he may intelligently treat them, or refer the patient to the proper specialist according to their cause; for instance, pyorrhea cases to the dentist; ear, nose and throat infections to the laryngologist; psychic and neurogenic disturbances to the neurologist; ptomain poisoning, gastrogenic or enterogenic dyspepsias to the gastro-enterologist; functional and organic disease to the practitioner or internist, and inflammatory, ulcerative or obstructive lesions of the small intestine, colon or rectum to the coloproctologist.

INFLAMMATORY AND ULCERATIVE ENTEROCOLITIS (DIARRHEA)

Usually diseases causing diarrhea are characterized by inflammatory or obstructive intestinal lesions, formerly discussed under the captions of intestinal catarrh, tuberculosis, syphilis and dysentery; but increased knowledge concerning the etiology of diarrhea makes this classification untenable.

The term "dysentery" is often applied to intestinal ailments characterized by the syndrome of abdominal pain, rectal tenesmus and fluid evacuations containing blood and mucus; but the appellation is confusing, since it has no etiologic significance, and dysenteroid manifestations complicate all ulcerative intestinal affections.

Generally chronic diarrhea is incited by colitis. Would it not be wise to forget so-called "dysenteries" and to classify inflammatory and ulcerative intestinal lesions etiologically, an arrangement that would permit their discussion under the headings of catarrhal, tuberculous, syphilitic, gonorrheal, endamebic, bacillary, balantidic, flagellate, coccidic and helminthic colitis and enterocolitis?

Catarrhal Enterocolitis.—This affection is quite common, and may be associated with constipation or diarrhea, when the mucosa is inflamed, excoriated or ulcerated. Catarrhal colitis can be relieved usually

* Read before the Section on Gastro-Enterology and Proctology at the Sixty-Eighth Annual Session of the American Medical Association, New York, June, 1917.