origin. The mercurial stomatitis had evidently given rise to a parotitis with possible mechanical compression of the nerve trunk. The Wassermann was found to be negative and on the suspension of the mercury the paralysis cleared up.

de Jong, S. I. MALARIA AND THE BORDET WASSERMANN REACTION. [Bulletins et mémoires de la Société médicale des hôpitaux de Paris, January 23, 1919.]

The author disputes the conclusion previously reached by Aimé and Lochelongue that malaria causes a positive Wassermann reaction. On the basis of 300 tests, of which two thirds were made in malarial subjects, he maintains that the Bordet Wassermann reaction retains its usual diagnostic value in malarial subjects, unless the serum examined is obtained during an actual malarial paroxysm. At this time the natural hemolytic power of the serum is often absolutely nil, as is the case in any acute affection associated with violent febrile paroxysms, and there is an apparent positive reaction through absence of hemolysis. Furthermore, this false positive reaction during malarial paroxysms is not even constantly observed, for the author obtained some negative reactions with blood obtained at the very beginning or from twelve to thirty hours after a typical paroxysm.

Wimmer, A. NONSPECIFIC MENTAL DISEASE IN SYPHILITICS. [Hospitalst., Sept. 4, 1918.]

The author calls attention to the more usual form of acute syphilitic psychoses and urges caution in referring the illness to syphilis when the disorder resembles the manic depressive psychosis. Twelve cases are described in detail in which mental symptoms resembling hysteria coincided with nervous manifestations. In one woman of 29, the clinical picture was dementia. In other cases there was an attack of mania before the infection with syphilis. Necropsy confirmed the assumption of an old endogenous manic-depressive psychosis with recent syphilitic meningitis. In another paranoid psychosis with hallucinations in a pronounced psychopathic. Even with the four syphilitic reactions positive in the spinal fluid, it does not follow that the psychosis observed is necessarily of syphilitic origin.

Wechsler, I. S. OPHTHALMIC CHANGES IN TABES AND PARESIS. [New York State Journal of Medicine, 1918, Vol. XVIII, p. 312.]

The writer gives the impression that there is no fundamental difference between tabetic neurosyphilis and so-called cerebrospinal or diffuse neurosyphilis. An inflammatory process is behind every form of syphilitic involvement and the spirochete is responsible. Obviously, the inflammatory reaction is in direct proportion to the kind of tissue involved. There is every reason why the meninges should respond more