ing the infection. Some preliminary work in the laboratory on the appendix indicates that it is not a common habitat for hemolytic streptococci.

RÔLE OF HEMOLYTIC STREPTOCOCCI AS PRIMARY AND SECONDARY INVADERS

Summing up these observations on the distribution of hemolytic streptococci, the facts point strongly to the crypts of the tonsils as being the main and in a sense a normal habitat of these organisms in the human body. Perhaps they are as normal to the tonsil crypts as is *Streptococcus viridans* to the buccal mucosa, *Staphylococcus albus* to the skin, or the colon bacillus to the intestine. From here they may be disseminated to adjacent tissues, ready possibly to take advantage of certain conditions, especially diseases of the respiratory tract, and give rise to secondary infections of a most serious nature.

The question arises as to whether or not persons are commonly or usually infected with their own hemolytic streptococci, or are the invading streptococci in infected persons of foreign origin? No doubt the latter statement is often true, as seems clearly proved in the case of sore throat epidemics of milk origin. Furthermore, many facts point to the common dissemination of streptococci by means of droplet infection. The route of infection is not so clear, however, in secondary streptococcus infections following other diseases. In the common terminal infections so often due to this organism there is little doubt that the individual is killed by his own streptococci.²⁶ It has been pointed out that

such streptococci are as virulent as epidemic streptococci for man, especially after growing in the body for a time. It would seem reasonable to assume, as has been assumed by some in connection with the streptococcus empyemas of last year, that individuals are being infected from time to time with their own streptococci, especially following the contagious diseases, and that by this process the streptococci become more and more virulent and aggressive, reaching a point ultimately at which the small doses contained in infected droplets or in other vehicles are sufficient to transmit the disease to a normal person. Thus a secondary invader may become transformed into a dangerous primary infective agent.

26. Richie, D. G., and Goehring, Carl: J. M. Res. 38: 421 (July) 1918.

Nonworkers Should Hibernate.—A correspondent in the *British Medical Journal* (February 26) advocates as a means of food saving a modified hibernation on the part of nonworkers. He says that, roughly, five sixths of food values go to producing and maintaining animal heat, and one sixth to the production of physical and mental energy. Of every 100 calories of heat produced within the body 80.1 per cent. is lost by radiation and conduction; 14.7 by evaporation; 2.6 per cent. in heating food and drink consumed, and 2.6 per cent. in heating inspired air. Lying in bed till late in the day by nonworkers, he says, will clearly reduce the heat loss and output of energy, with consequent saving in food requirement. He therefore says the idle poor, the rich, infirm and insane persons should be compelled to remain in bed, warm and at rest till noon daily.

THE IMPORTANCE OF THE LOUSE PROBLEM *

HARRY PLOTZ, M.D. (BROOKLYN)

Major, M. C., U. S. Army

WASHINGTON, D. C.

The demobilization of armies is usually associated with the introduction of epidemic diseases into civil communities. There is great possibility of louse-borne diseases gaining entrance into the United States unless necessary measures are taken to exclude them. The question as to whether epidemic influenza could have been excluded from this country remains an open one; but it can be said with a certain degree of certainty, that with the application of proper measures, epidemic typhus fever, trench fever and relapsing fever can be excluded. In view of the fact that the louse has spread more disease in the present war than all the other insects combined, its importance as a disease carrier can be appreciated. It is important, therefore, that proper facilities be provided to delouse all returning troops, that the public health authorities provide measures for the delousing of immigrants, and that physicians be on the alert for the possible occurrence of louse-borne diseases, leading to the early isolation of those affected.

Vermin infestation is now prevalent throughout Europe, the annoyance of the "cootie" being well known to all American soldiers. Coincident with this widespread vermin infestation exists the impoverishment of most European peoples. These conditions usually pre-

cede a typhus fever epidemic. This fact is often related in the history of typhus fever. Epidemic typhus fever has already made its appearance in European countries, and with the advent of winter, when the disease is most common, we have much to fear from a general typhus fever epidemic. The Allied troops are coming more and more in contact with the Russian and Balkan peoples, and the spread of this disease from army to army and then from country to country need be anticipated. For these reasons it is believed that due consideration should be given this problem, and proper measures be employed to exclude these diseases from the United States. It is important to point out that endemic typhus fever (Brill's disease) is a disease not uncommon in this country; and with the introduction of vermin, we may fear a greater number of cases and a possible increase in its virulence.

The eradication of vermin infestation calls for strict adherence to a certain definite plan of treatment. It has been said that vermin infestation in European armies started from a few infested individuals, and owing to the close association of the troops, the infestation rapidly spread. The escape of one vermin-infested individual into a community may cause considerable infestation. A vermin-infested man, then, should be regarded in the same light in which one regards a meningococcus carrier: he is a danger to the community until his condition is cleared up.

* From the Division of Sanitation, Medical Department, U. S. Army.

Fig. 1.—Scheme for delousing.

FUNDAMENTAL PRINCIPLES

In order that the best results may be obtained, there are certain fundamental principles regarding the subject of delousing that must be taken into consideration.

1. The clothes louse (*Pediculus humanus-corporis*) transmits typhus fever, trench fever and relapsing fever. These diseases are not transmitted by the bite of the louse but by a virus which is contained in the excreta. This virus gains entrance through the puncture wound made by the louse while sucking, or is scratched in by the infested individual. Therefore, a method which destroys only the lice and eggs without causing the destruction of the virus in the excreta is not sufficient. Delousing, then, should be defined as that process which destroys lice and their eggs and the virus concerned in the transmission of disease.

The clothes then come out dry. If shrinkage of woolen materials result, the disinfector is not properly operated. Shrinkage of woolen materials is due to combined moisture and high temperature; therefore, if the moisture is kept at a minimum and the temperature at a point sufficient to sterilize, this will not occur. In no instance should leather material (such as shoes, puttees and hat) or rubber, celluloid material or money be sterilized. The occurrence of vermin in these materials is rare.

ARMY PLAN

The Army proposes to handle the problem in this manner: All troops will be detained at foreign ports for a period of two weeks, during which time persons with infectious and epidemic diseases will be isolated and universal delousing will be practiced before

Fig. 2.—Plan of delousing plant.

2. Various methods have been recommended for delousing, such as hydrocyanic acid gas, chloropicrin, and various insecticides and heat appliances. It is to be noted that the virus of relapsing fever is a spirochete, while that of typhus fever is a bacillus. As these gases and insecticides are not bactericidal, they cannot be regarded as perfect delousing agents. In all instances in which heat is used, a high enough temperature must be employed to destroy the virus as well as lice and eggs. Steam has been found to be the best delousing agent.

3. All persons coming from a vermin-infested area should be regarded as if they were vermin infested. Examination for vermin in large bodies of men, even in the hands of experts, is of little value. A great deal of time is consumed in the examination, and the final result is of little use.

4. Clothes lice may lay their eggs on the hair of the head and of the axillary and pubic regions. This must be taken into consideration when treating this condition.

Clothes lice should be treated by sterilizing all clothing by exposure to steam under pressure for thirty minutes followed by ten minutes in a vacuum.

embarking. Delousing will be practiced on transports when necessary. All soldiers on transports will be instructed to devote fifteen minutes each day to the search for vermin in their clothing. Universal delousing will be again practiced at debarkation camps in the United States. This will serve as a check on the first delousing.

AN EFFICIENT DELOUSING PLANT

All sick and wounded will be deloused at the debarkation hospital. An efficient method would be to have the delousing room in association with the receiving ward. The following plan is suggested for this purpose. With this method vermin infestation would be prevented from being carried to the wards.

In order that the greatest number of men may be handled in the shortest period of time, it is important to have a rapidly functioning delousing plant. According to the plan here proposed, 260 soldiers and all their equipment can be deloused every hour. According to this plan the man enters with his barrack bag containing all his clothing. The leather materials, rubber, celluloid materials and money are passed in at the locker room. The man receives two numbered tags

corresponding to the number of the locker, and then proceeds to the disrobing room with his barrack bag. All his clothing is placed in the bag, which is tied and numbered with one of the tags, the man retaining the remaining tag. The bag is then placed in the carriage, which is pushed into the steam sterilizer. The sterilizer is 18½ feet long by 5 feet in diameter and is provided with two cars and transfer tracks, so that one car will be loading or unloading while the other is in the sterilizer.

The soldier then proceeds to the hair-cutting room. Before entering, he is inspected for vermin and nits. If vermin and nits are found in the hair of the head, axillary or pubic region, he is passed into the barber shop. If none are found, he passes into the shower room. In the barber shop the hair is cropped with an electric hair-cutting machine. The axillary and pubic hair can be shaved in this room. Following this he enters a shower room, where a bath with liquid soap and warm water is obtained. The soap used is a kerosene soap mixture. A drying room follows, a table being provided for the clean towels and a receptacle to receive the soiled ones.

Provision is made in this plan for a pressing room, as the clothing is wrinkled following steam sterilization. The pressing of garments, however, would somewhat retard the process. The leather material, etc., is reclaimed at the locker room. The floor of this building is made of concrete. Two toilets are provided, one for the dirty men and one for the clean men. All attendants wear the louse-proof suit, which is a one-piece garment covering the shoes and tied about the neck.

With an arrangement such as this, when universal delousing is practiced, there should be no danger of vermin infestation or louse-borne diseases gaining entrance into the United States.

INTESTINAL PARASITES DURING A YEAR AT THE WALTER REED GENERAL HOSPITAL *

M. W. LYON, JR., M.D.
Captain, M. C., U. S. Army
WASHINGTON, D. C.

During the year October, 1917, to October, 1918, there were examined at the laboratory of the Walter Reed General Hospital 477 samples of feces for intestinal parasites. In nearly every instance the samples were submitted because the patient had shown some evident or obscure reason for suspecting the presence of intestinal animal parasites, although there are included a number of samples that were sent for examination for typhoid or dysentery bacilli. The 477 stools came from 163 individuals, seventy-two of whom, or 44 per cent., were found to harbor one or more kinds of parasites, while in ninety-one persons, or 56 per cent., no parasites were found.

With the exception of *Endameba histolytica*, none of these cases of intestinal parasites can be said to possess much medical interest. Their presence is one of greater zoological than medical interest. Next to the ameba infections the hookworm cases are of most interest. Most of these were very light infections,

but they are an indication of how widely scattered this parasite has probably become during the formation of the Army.

The method practiced in looking for protozoan parasites is to place a drop of the feces, either plain or mixed with a little water or salt solution, on a slide and press it out with a cover glass. No attempt is made to concentrate cysts. In searching for ova or embryos a considerable portion of the stool is emulsified with water, usually by pouring water into the container in which it is sent to the laboratory. The liquid material is then poured through two thicknesses of gauze into a urine sedimentation glass. After a distinct sediment has formed in the point of the glass, the brown supernatant water is poured off. More water is then roughly added so that the sediment is thoroughly stirred up, and it is allowed to settle again. This washing is repeated several times until the supernatant water is essentially clear. After the final sedimentation, about 0.25 c.c. of the upper layers of the sediment is spread on a slide without a cover glass and search made with the low powers of the microscope, a mechanical stage being used. The spread occupies most of the slide, and by the use of a wax-penciled border is kept from running over its edges. The chief advantage in the washing and sedimentation is that innumerable small particles, mainly bacteria, are got rid of, and a very clear sediment is left to examine. The washing may appear time consuming, but it takes only a moment to pour off and fill up again, and while the settling is taking place other things can be attended to. Examination of one spread slide is considered sufficient to establish the presence of ova. In a few cases, however, an especially enthusiastic searcher has found an ovum or an embryo in examining a second spread which had not been found on the first.

Hookworm, *Necator americanus*, was found most frequently, occurring in 34.2 per cent. of the parasites and 17.2 per cent. of the persons examined. The number of worms was always small. The largest number recovered from complete stools after hookworm treatment was seventy-one and the smallest two. Its presence often caused a slight eosinophilia. The first case found was in a patient whose blood, on being examined for malaria parasites, showed so many eosinophils that a stool for examination was requested. The farthest north from which the hookworm cases came was Virginia and Tennessee. Hookworm ova were found alone twenty-three times; with *Strongyloides* embryos twice; with *Endameba coli* twice; with *Ascaris* and *Trichuris* twice, and with *E. coli*, *Trichuris* and *Trichomonas* once.

Strongyloides, occurring in 4.3 per cent. of the persons examined and 8.5 per cent. of the parasites, was found alone five times and twice with *Necator*. With two exceptions they were not present in large numbers. In one of these they were the only parasite present, and in the other they were with the second heaviest hookworm infestation, and both were in patients in the tuberculosis ward. In the case occurring with hookworm, successful treatment for the latter failed to have any effect in reducing the number of *Strongyloides* embryos.

The five *Taenia saginata* cases were not diagnosed in the first instance by the laboratory, but having been found elsewhere, were sent into the hospital for treatment.

* From the Laboratory Service of the Walter Reed General Hospital.