Notes.

The Editor desires to point out that the pages of the Journal are open for the inclusion of short notes dealing with analytical practice and kindred matters. Such notes are submitted to the Publication Committee in the usual manner.

DAKAMBALLI STARCH.

A sample of this starch was received from Dr. Vevers of British Guiana through the Wellcome Bureau of Scientific Research. It is prepared by the Indians of British Guiana from the fruit of the tree *Aldina insignis*, and is used

by them as a remedy for dysentery, being administered in the form of a mucilage made by boiling the starch in milk or water.

The starch was of pale brown colour and, on examination under the microscope, was seen to consist entirely of starch granules varying in mean diameter from 11 to 42 micromillimetres. The small granules were mostly circular, a few being truncated and the larger ones roughly ovate. The hilum was at the broader end, and the concentric rings were well marked. The material was tasteless.

On analysis the following results were obtained:—

	Per Cent.
Moisture	19·57
Fat	0·04
Proteins N $\times 6\cdot25$	1·25
Crude fibre.. ..	0·10
Matter soluble in alcohol ..	trace
Matter soluble in cold water	"
Ash	0·21
Starch, by difference	78·83
	<hr/> 100·00 <hr/>

The amount of starch directly estimated by the method of Davis and Daish (*Journal of Agricultural*

Science, 1914, 6, 152) was 76·1 per cent.

No alkaloids or cyanogenetic glucosides could be detected in the sample.

It is well known that mucilaginous substances are sometimes effective remedies in colitis and certain forms of diarrhoea and dysentery, and it is not surprising, therefore, that a mucilage made from clean starch such as this should prove equally useful.

J. A. GOODSON.

WELLCOME CHEMICAL RESEARCH LABORATORIES,
6, KING STREET, SNOW HILL, E.C. 1.

THE USE OF ULTRA VIOLET LIGHT IN ANALYSIS.

It is now fairly well known that the fluorescent effects produced by the action of ultra violet light upon substances have been applied, to a small extent, to work of an analytical nature. In connection with precious stones fluorescence will, in many cases, indicate their place of origin, e.g. will discriminate between Japanese pearls and pearls from other localities. Vegetable and mineral oils can be distinguished, and there is, I believe, some similar application in the cellulose industry. A very large number of substances are fluorescent to some extent, and though some shade of blue seems to be the commonest tint, yet almost every colour can be obtained.

It seems probable that the use of ultra violet light could be extended very considerably, and the results of a few observations made recently are given below with a view to suggesting some of the directions in which research may be profitable.

Cotton is readily distinguished from wool or silk. Various kinds of paper and card differ considerably from one another; this may be due to differences in the sizing, for casein fluoresces more strongly than gelatin or any other proteid material I have tried, and more strongly also than cellulose itself. Even on fabrics, many dyes which are fluorescent can readily be distinguished from other dyes of the same colour which are not. Uranium salts fluoresce brightly in the solid state, and 0.1 per cent. of the nitrate can be detected in solution; a borax bead containing a trace of the oxide is colourless, but distinctly fluorescent. Acetone can be detected in alcohol; 1 per cent. giving a distinct effect. Among the substances which fluoresce most brilliantly are quinine, æsculin, and the dye uranine, and very minute quantities of these substances can be detected by means of ultra violet light. With quinine, one part in a hundred million of water; with uranine, one part in a thousand million; and with æsculin, one part in ten thousand million, show distinct fluorescence under favourable conditions: we have here a sensitiveness comparable with that of the spectroscope.

Suitable means of obtaining ultra violet rays for these experiments are afforded by a powerful source of light in a light-tight box with a small window containing a filter of Chance's ultra violet glass. This latter transmits light between 4000 and 3000 A.U., with a maximum at 3600; the visible spectrum is completely cut out, with the exception of a trace of violet and the extreme red. An ordinary arc lamp will serve as the source of light; but an arc between nichrome or tungsten poles is better. In some respects a mercury vapour lamp is better still, as it gives practically no red rays, which latter are apt to interfere in certain cases. The infra red can be removed by a cell containing copper sulphate or acetate solution, but this diminishes also the intensity of the ultra violet. In some cases it is advantageous to concentrate the rays with a quartz lens.

A. F. KITCHING.

MONKTON COMBE SCHOOL,
NEAR BATH.

Notes from the Reports of Public Analysts.

The Editor would be glad to receive the Annual or other Reports of Public Analysts containing matter of interest to the Society. Notes made from such Reports would be submitted to the Publication Committee.