9. *The CAMBRIAN HORIZONS of COMLEY (SHROPSHIRE) and their BRACHIOPODA, PTEROPODA, GASTEROPODA, etc.* By EDGAR STERLING COBBOLD, F.G.S. (Read April 21st, 1920.)

[PLATES XXI-XXIV.]

CONTENTS.

	Page
I. Introduction	325
II. The Comley Horizons (with Table I)	325
III. Classification, Terminology, and References	326
IV. Bibliography	326
V. The Comley Brachiopoda	328
VI. The Comley Pteropoda, etc.	354
VII. The Comley Gasteropoda	364
VIII. The Comley Ostracoda	367
IX. The Faunas of the Comley Horizons (with Table II) and their Order of Appearance	368
X. General Summary	374

I. INTRODUCTION.

THE objects of this paper are to define the horizons of the Cambrian system at Comley, to describe and illustrate the species of brachiopoda, pteropoda, and gasteropoda, and to discuss the relationships of the faunas one to the other and to those of other Cambrian areas.

The majority of the specimens now described were obtained from the excavations carried out between the years 1907 and 1914, on behalf of the Committee for the Excavation of Critical Sections in Palæozoic Rocks, appointed by the British Association for the Advancement of Science. Some specimens were collected in 1892 for H.M. Geological Survey by Mr. J. Rhodes, and others have been obtained by myself or other observers at various dates since 1890.

II. THE COMLEY HORIZONS.

Details of the sections opened by excavation have been given in the reports of the meetings of the British Association held at Dublin in 1908, Winnipeg in 1909, Sheffield in 1910, Portsmouth in 1911, Dundee in 1912, and Manchester in 1915.

In Table I (facing p. 326) the various horizons from which different assemblages of fossils have been obtained are set out in descending order, with the proposed nomenclature in spaced type, followed by index-letters, thicknesses, general lithological characters, and suggested correlations. The designations used by me in previous communications and in my reports to the British Association are inserted in ordinary type in the first column. Since these

differ considerably from those now proposed, a few words of explanation seem to be necessary.

As the study of the fossils collected from the different beds proceeded, it became apparent that they might prove to be of more than local interest, and it was therefore desirable that the original indefinite or cumbersome terms, based on local names, or lithological variations, should be abandoned in favour of others based upon the fossil contents. It was also evident that, in several instances, beds or groups of beds, to which a local name had been allotted, contained more than one set of fossil forms, and it seemed undesirable to resort to any extended use of such terms as Upper, Middle, Lower, etc.

Thus the 'Lower Comley Sandstone' has been divided into five parts, and the three divisions previously called 'The Grey Limestones,' 'The Black Limestone,' and 'The Shoot-Rough Road Flags,' into two horizons in each case.

III. CLASSIFICATION, TERMINOLOGY, AND REFERENCES.

The classification and terminology used in the descriptions of the brachiopoda follow closely those adopted by Dr. C. D. Walcott. In view of the very full references given by him, it has not been considered necessary to quote more than (1) the work in which any species was first described, (2) the pages, plates, and figures in his monograph, and (3) any references to works later than those to which he had access.

In order to avoid repetition, works enumerated in footnotes are designated by the names of the authors and dates. The full references may be found by referring to the Bibliography.

I have followed the usual practice in including such tubular fossils as *Hyolithellus*, *Helenia*, etc. among the Pteropoda, but am doubtful whether they really belong to that order.

IV. BIBLIOGRAPHY.

- J. BARRANDE. 1867.—'Système Silurien de la Bohême: vol. iii.—Pteropoda.'
E. BILLINGS. 1861.—Geol. Surv. Canada. 'Palaeozoic Fossils' vol. i.
E. BILLINGS. 1872.—Can. Nat. n. ser. vol. vi, pp. 465-80. 'On some Fossils from the Primordial Rocks of Newfoundland.'
C. CALLAWAY. 1874.—Q. J. G. S. vol. xxx, p. 196. 'On the Occurrence of a Tremadoc Area near the Wrekin in South Shropshire.'
C. CALLAWAY. 1877.—*Ibid.* vol. xxxiii, pp. 652-72. 'On a New Area of Upper Cambrian Rocks in South Shropshire.'
C. CALLAWAY. 1878.—*Ibid.* vol. xxxiv, pp. 754-63. 'Quartzites of Shropshire.'
E. S. COBBOLD. 1909.—Rep. Brit. Assoc. 1908, Dublin, pp. 231-42. 'Report on some Excavations in the Cambrian Rocks of Comley (Shropshire).'
E. S. COBBOLD. 1910 a.—*Ibid.* 1909, Winnipeg, pp. 181-86. 'Report on some further Excavations among the Cambrian Rocks of Comley (Shropshire).'
E. S. COBBOLD. 1910 b.—Q. J. G. S. vol. lxxi, pp. 19-51. 'On Trilobites from the Cambrian Rocks of Comley.'
E. S. COBBOLD. 1911 a.—Rep. Brit. Assoc. 1910, Sheffield, pp. 113-22. 'Third Report on Excavations among the Cambrian Rocks of Comley.'
E. S. COBBOLD. 1911 b.—Q. J. G. S. vol. lxxii, pp. 282-311. 'Trilobites from the *Paradoxides* Beds of Comley.'
E. S. COBBOLD. 1912.—Rep. Brit. Assoc. 1911, Portsmouth, pp. 111-15. 'Fourth Report on Excavations among the Cambrian Rocks of Comley.'

TABLE I.—THE COMLEY CAMBRIAN HORIZONS.

	Local or lithological names used by the Author in previous publications.	Proposed names for the faunal horizons.	Index- letters.	Estimated or observed thicknesses in feet.	General lithological characters.	Approximate equivalents in Europe. in America.	
The Comley Sandstone. Upper Division. ↓	Shoot-Rough Wood Shale.		Cx	?	Soft blue-grey shales.	{ Shineton Shales, Shropshire. Tremadoc, North Wales.	
	Shoot-Rough Road Shale.	<i>Orusia</i> Shales.	Ca	? 500	{ Micaceous shale, with thin mudstone or calcareous courses.	{ <i>Orusia-lenticularis</i> Beds, Scandinavia. <i>Orusia-lenticularis</i> Beds, North Wales.	
	Shoot-Rough Road Flags.	{ <i>Billingsella</i> Beds.	Bc	5	{ Fine and coarse grits, with calcareous and shaly beds.	{ <i>Paradoxides-forchammeri</i> Zone, Scandinavia.	
		{ <i>Paradoxides-davidis</i> Flags.	Bb ₅	18	{ Coarse gritty flags, with a calcareous bed at the summit.	{ <i>Paradoxides-davidis</i> Beds, Scandinavia. <i>Paradoxides-davidis</i> Beds, North & South Wales. Upper <i>Davidis</i> Fauna, Nuneaton.	
	Shoot-Rough Road Sandstone.	{ <i>Paradoxides-rugulosus</i> Sandstone.	Bb ₄	?	{ Green and brown sandstone, with a calcareous bed at the summit.	Lower <i>Davidis</i> Fauna, Nuneaton.	
	Unexplored interval.		Bb ₃	?	?	? <i>Hartshillia</i> Fauna, Nuneaton.	
	Hill-House Shale.		Bb ₂	? 300	{ Micaceous flaggy shales, with occa- sional beds of grit.	{ <i>Paradoxides-hicksi</i> Zone, South Wales & Nuneaton.	
	Hill-House Grits.	{ <i>Paradoxides-intermedius</i> Grits.	Bb ₁	5 to 38	{ Coarse conglomeratic grits and breccia of fragments from Ac ₂ & Ac ₁ below.	{ <i>Paradoxides-hicksi</i> Zone, Scandinavia.	
	Non-sequence, with overlap on to all beds from Ab ₅ to Ba ₃ .						
	Hill-House Flags.	<i>Dorypyge-lakei</i> Flags.	Ba ₃	4+	Phosphatic gritty flags.	{ <i>Paradoxides-elandicus</i> Zone, Scandinavia. Purley Shales (<i>pars</i>), Nuneaton.	
	Quarry-Ridge Shales.		Ba ₂	300	{ Fine blue and brown shales, with bands or beds of hard, ringing grit.		
	Quarry-Ridge Grit.	<i>Paradoxides-groomi</i> Grits.	Ba ₁	25	{ Hard, ringing, glauconitic grit, with conglomerate and a thin phosphatic deposit at the base.		
	Unconformity, with erosion down to beds Ab ₄ below.						
The Comley Sandstone. Lower Division. ↑	The Black Limestone.	{ <i>Lapworthella</i> Limestone.	Ad	0.5	{ Calcareous and phosphatic granular material.	{ ? Purley Shales (<i>pars</i>), Nuneaton.	
		{ <i>Protolenus</i> Limestone.	Ac ₅	0.5	{ Black to pale-grey, compact, phosphatic limestone.		
	The Grey Limestones.	{ <i>Strenuella</i> Limestone.	Ac ₄	0.75	{ Grey to red-purple gritty limestone, with phosphatic inclusions.		{ North Attleboro' Beds, Massachusetts; <i>Callavia</i> Beds, Manuel's Brook (Newfoundland).
		{ <i>Microdiscus-bellimarginatus</i> Limestone.	Ac ₃	1.75	{ Grey and pinkish-grey limestone, with much phosphate in places.		
	The <i>Olenellus</i> Limestone.	<i>Olenellus</i> Limestone.	Ac ₂	2.5	{ Red or purple sandy limestone, no- dular.	Camp-Hill Quartzite, Nuneaton.	
	Lower Comley Sandstone.	{ <i>Callavia</i> Sandstone.	Ac ₁	?	{ Bright green sandstone.		
			Ab ₄		{ Green sandstone, with nodular calca- reous beds.		
		{ <i>Holmia</i> Sandstone.	Ab ₃	450	{ Green and reddish sandstone, with calcareous beds.		
		<i>Obolella-groomi</i> Beds.		Ab ₂		{ Muddy sandstone with some flaggy beds.	Hollybush Sandstone, Malvern.
				Ab ₁	27	{ Greenish-grey sandstone and mottled conglomeratic beds below.	Tuttle-Hill Quartzite, Nuneaton (after Lapworth).
The Wrekin Quartzite.			Aa	22	{ Yellow sandy quartzite, graduating down into compact blue quartzite which becomes conglomeratic at the base.	Malvern Quartzite, Malvern Hills. Park-Hill Quartzite, Nuneaton (after Lapworth).	

- E. S. COBBOLD. 1913 *a*.—Rep. Brit. Assoc., 1912, Dundee, pp. 136-43. 'Fifth Report on Excavations among the Cambrian Rocks of Comley.'
- E. S. COBBOLD. 1913 *b*.—Q. J. G. S. vol. lxi, pp. 27-50. 'The Trilobite Fauna of the Comley Breccia-Bed.'
- E. S. COBBOLD. 1914.—Rep. Brit. Assoc. 1913, Birmingham, pp. 486-87. 'Critical Sections of the Cambrian Area called "The Cwms" in the Caradoc-Comley Region of Shropshire.' (Abstract.)
- E. S. COBBOLD. 1916.—Rep. Brit. Assoc. 1915, Manchester, pp. 117-23. 'Sixth Report on Excavations among the Cambrian Rocks of Comley.'
- E. S. COBBOLD. 1919.—Geol. Mag. pp. 149-58. 'Cambrian Hyolithidae from Harts-hill in the Nuneaton District (Warwickshire).'
- T. R. CONRAD. 1840.—Ann. Rep. Geol. Surv. New York, pp. 199-207.
- T. DAVIDSON. 1868.—Monogr. Brit. Foss. Brach. (Pal. Soc. vol. xxii) pt. vii, No. 3. 'Silurian Brachiopoda.'
- T. DAVIDSON. 1871.—*Ibid.* (vol. xxiv) pt. vii, No. 4. 'Silurian Brachiopoda.'
- S. W. FORD. 1873.—Amer. Journ. Sci. ser. 3, vol. vi, pp. 211-15. 'On some New Species of Fossils from the Primordial or Potsdam Group of Rensselaer County (New York).'
- A. W. GRABAU. 1900.—Occ. Pap. Boston Nat. Hist. Soc. vol. iv, pp. 601-94. 'Palæontology of the Cambrian Terranes of the Boston Basin.'
- A. W. GRABAU & H. W. SHIMER. 1909.—'North American Index-Fossils: Invertebrates,' vol. i. Lancaster, Pennsylvania.
- T. T. GROOM. 1902.—Q. J. G. S. vol. lviii, pp. 89-149. 'Cambrian & Associated Beds of the Malvern Hills.'
- J. HALL. 1847.—Geol. Surv. N.Y., Palæontology, vol. i, pt. ii.
- J. HALL. 1879.—*Ibid.* vol. v, pt. ii.
- H. HICKS. 1872.—Q. J. G. S. vol. xxviii, pp. 73-83. 'Undescribed Fossils from the Menevian Group.'
- H. B. HOLL. 1865.—*Ibid.* vol. xxi, pp. 72-102. 'Geological Structure of the Malvern Hills & Adjacent Districts.'
- G. HOLM. 1893.—Sver. Geol. Undersökn. ser. C, No. 112.
- V. C. ILLING. 1916.—Q. J. G. S. vol. lxxi (1915) pp. 386-450. 'The Paradoxidian Fauna of the Stockingford Shales.'
- J. KLER. 1916.—'The Lower Cambrian *Holmia* Fauna at Tömten (Norway).'
- C. LAPWORTH. 1898.—Geol. Mag. pp. 484-86. 'On the Discovery of the *Olenellus* Fauna in the Lower Cambrian Rocks of Great Britain.'
- C. LAPWORTH. 1891.—*Ibid.* pp. 529-36. 'On *Olenellus callavei* & its Geological Relationships.'
- C. LAPWORTH & W. W. WATTS. 1894.—Proc. Geol. Assoc. London, vol. xiii, pp. 297-355. 'The Geology of South Shropshire.'
- C. LAPWORTH & W. W. WATTS. 1910.—*Ibid.* Jubilee volume, pp. 739-69. 'The Geology of Shropshire.'
- G. LINNARSSON. 1876.—Bihang K. Svenska Vetensk.-Akad. Handl. vol. iii, No. 12, pp. 1-34. 'The Brachiopoda of the *Paradoxides* Beds of Sweden.'
- C. A. MATLEY. 1902.—Q. J. G. S. vol. lviii, pp. 135-47. 'On the Cambrian Brachiopoda of the Malvern Hills.'
- C. A. MATLEY. 1911.—*Ibid.* vol. lxxvii, pp. 300-306. 'On some of the Brachiopoda from the *Paradoxides* Beds of Comley.'
- G. F. MATTHEW. 1895.—Trans. N.Y. Acad. Sci. vol. xiv, pp. 101-53. 'The *Pro-tolenus* Fauna.'
- G. F. MATTHEW. 1899 *a*.—Bull. Nat. Hist. Soc. New Brunswick, xviii, pp. 189-97. 'Preliminary Notice of the Etcheminian Fauna of Newfoundland.'
- G. F. MATTHEW. 1899 *b*.—Trans. R. Soc. Canada, ser. 2, vol. v, sect. 4, pp. 39-66. 'Studies in Cambrian Faunas: No. 3, Upper Cambrian Fauna of Mount Stephen.'
- G. F. MATTHEW. 1901.—Bull. Nat. Hist. Soc. New Brunswick, vol. iv, pt. 4, No. 19, pp. 303-304. '*Acrothyra*, a New Genus of Etcheminian Brachiopoda.'
- G. F. MATTHEW. 1902 *a*.—*Ibid.* vol. iv, pt. 5, No. 20, pp. 377-425. 'Additional Notes on the Cambrian of Cape Breton, with Descriptions of New Species.'
- G. F. MATTHEW. 1902 *b*.—Canadian Record of Sci. vol. viii, No. 7, pp. 437-68. 'Ostracoda of the Basal Cambrian Rocks in Cape Breton.'
- J. C. Moberg. 1892.—Geol. Fören. Stockholm Förhandl. vol. xiv, pp. 103-20. 'Om en nyupptäckt Fauna i Block af Kambrisk Sandsten, insamlade af Dr. N. G. Holst.'
- J. C. Moberg. 1910.—Sver. Geol. Undersökn. ser. C, No. 229, Årsbok 4, No. 1. 'Historical-Stratigraphical Review of the Silurian of Sweden. English Translation.'

- O. NOVÁK. 1891.—Abhandl. K. Böhm. Gesellsch. Wissensch. ser. 7, vol. iv, No. 6, Prag. 'Revision der Paläozoischen Hyolithiden Böhmens.'
- F. RAW. MS. quoted by C. D. Walcott, 1910.
- J. W. SALTER. 1859.—Q. J. G. S. vol. xv, pp. 374-81. 'Organic Remains from the Durness Limestone.'
- N. S. SHALER & A. F. FÖRSTER. 1888.—Bull. Comp. Zool. Harvard Coll. vol. xvi, No. 2, pp. 27-41. 'Preliminary Description of North Attleborough Fossils.'
- R. TATE. 1891.—Trans. Roy. Soc. S. Aust. vol. xv, pp. 183-90. 'On Cambrian Fossils of South Australia.'
- K. VON SEEBACH. 1865.—Zeitschr. Deutsch. Geol. Gesellsch. vol. xvii, pp. 338-47. 'Beiträge zur Geologie der Insel Bornholm.'
- G. WAHLENBERG. 1821.—Nova Acta Regiæ Societatis Scientiarum Upsaliensis, vol. viii. 'Petrifacæ Telluris Svecanæ.'
- C. D. WALCOTT. 1884.—U.S. Geol. Surv., Bull. 10. 'On the Cambrian Faunas of North America.'
- C. D. WALCOTT. 1886.—*Ibid.* Bull. 30. 'Second Contribution to the Studies on the Cambrian Faunas of North America.'
- C. D. WALCOTT. 1889.—U.S. Nat. Mus. Proc. vol. xii, pp. 441-46. 'Description of New Genera & Species of Fossils from the Middle Cambrian.'
- C. D. WALCOTT. 1890.—10th Ann. Rep. U.S. Geol. Surv. pp. 515-658. 'The Fauna of the Lower Cambrian or *Olenellus* Zone.'
- C. D. WALCOTT. 1910.—Smithsonian Misc. Coll. vol. liii, No. 6. '*Olenellus* & other Genera of the Mesonacida.'
- C. D. WALCOTT. 1912.—Monogr. U.S. Geol. Surv. vol. li. 'Cambrian Brachiopoda.'
- C. D. WALCOTT. 1913.—Smithsonian Misc. Coll. vol. lvii, No. 11. 'New Lower Cambrian Sub-Fauna.'

V. THE COMLEY BRACHIOPODA.

The species hitherto collected from Comley are placed under the following families and genera :—

Order ATREMATA Beecher.		Page
PATERINIDÆ Schuchert.		
<i>Micromitra</i> (Meek) Walcott.	One species	328
Sub-gen. <i>Paterina</i> (Beecher) Walcott.	Five species	329
<i>Walcottina</i> , gen. nov.	Two species	334
OBOLIDÆ King.		
<i>Obolus</i> Eichwald.	Six species	338
<i>Lingulella</i> Salter	Three species	341
Order NEOTREMATA Beecher.		
OBOLELLIDÆ Walcott & Schuchert.		
<i>Obolella</i> Billings.	Two species	342
ACROTRETIDÆ Schuchert.		
<i>Acrothele</i> Linnarsson.	One species	345
<i>Acrotreta</i> Kutorga.	Six species	346
<i>Acrothyra</i> Matthew.	Two species	348
Order PROTREMATA Beecher.		
BILLINGSSELLIDÆ Schuchert.		
<i>Billingsella</i> Hall & Clarke.	Three species	353
<i>Orusia</i> Walcott.	One species	354

PATERINIDÆ Schuchert.

MICROMITRA (Meek) Walcott, 1912.

MICROMITRA sp. indet. (Pl. XXI, figs. 18 & 19.)

Cf. *Kutorgina cingulata* Billings, var. *pusilla* Linnarsson, 1876, p. 25 & pl. iv, figs. 53-54.

Cf. *Micromitra pusilla* Linnarsson, Walcott, 1912, p. 339 & pl. iii, figs. 2, 2 a-c.

Two specimens [2133, 2134]¹ indicate the presence of a form allied to that described by Linnarsson from the *Paradoxides-forchhammeri* Zone of Scandinavia. They are not sufficiently well-preserved to be identified with certainty.

A ventral valve is semicircular in outline, and has the apex recurved to a point beyond the posterior margin. The shell is corneous, dark, and shows radiating riblets crossed by very fine, impressed, concentric striae, which strengthen in places to concentric rugosities.

Dimensions.—Length = 2 mm.; width at about the half length = 2.5 mm.; height = 1 mm.

Locality and horizon.—Comley, Middle Cambrian; from the *Billingsella* Beds, horizon Bc.

Sub-genus *Paterina*.

MICROMITRA (PATERINA) PHILLIPSI Holl. (Pl. XXI, figs. 1-3.)

Oboella phillipsii Holl, 1865, Q. J. G. S. vol. xxi, p. 102, figs. 10 a-10 c.

Oboella (?) *phillipsi* (Holl), Davidson, 1866, Pal. Soc. Monogr., Brit. Foss. Brach. vol. iii, part 7, No. 1, p. 62 & pl. iv, figs. 17 a-19.

Katorgina cingulata Davidson, 1871, Pal. Soc. Monogr., Brit. Foss. Brach. vol. iii, pt. 7, No. 4, p. 342 & pl. l, fig. 25.

Micromitra (*Paterina*) *phillipsi* (Holl), Walcott, 1912, U.S. Geol. Surv. Monogr. vol. li, p. 351 & pl. iii, fig. 8.

Through the kindness of Prof. Sollas it has been possible to compare the Comley specimens with those in the University Museum, Oxford (Holl & Groom collections). There can be no doubt as to the reference of the Comley specimens to Holl's species. The forms of the tests and their method of preservation are identical, but there are some points in which the Comley shells exhibit features not seen in the specimens from Malvern at Oxford.

- (1) The external surface often, but not always, shows traces of radiating raised lines, as well as the lines of growth; but the two characteristics occupy different parts of the shell, and have not been seen to cross one another as they do in *M. sculptilis* Meek² (fig. 1).
- (2) The line of flexure between the lateral slopes and the false area is well rounded, not a sharp angle.
- (3) The false area is almost vertical; the triangular pseudodeltidium is wide; the angle at its apex (fig. 2, posterior view) is about 120°; it is convex, and extends some distance down towards the posterior margin.
- (4) A well-preserved interior, referred to the ventral valve [2335], with the corresponding internal cast [2334] (fig. 3) exhibits numerous radiating shallow channels, and also under the umbo three comparatively strong ridges, one in an axial position, the others making with it an angle of about 10° on each side.

The shells are widest at about a quarter of the length from the posterior margin.

¹ The numbers in square brackets are those attached to the specimens in my collection, unless otherwise stated.

² C. D. Walcott, 1912, p. 335, & pl. iii, figs. 5 & 5 a-5 c.

The lengths of the Comley specimens range from 4 to 5 millimetres, the ratio of length to width is about 1 to 1·3, and that of length to height about 1 to ·33.

Several of the best-preserved specimens are in the collection made by Mr. Rhodes for H.M. Geological Survey.

Locality and horizon.—Lower Cambrian. The Cwms sub-area of Comley and also west of Comley Quarry; from the *Obolella-groomi* Beds, horizon Ab₁.

MICROMITRA (PATERINA) RHODESI, sp. nov. (Pl. XXI, figs. 4 *a*–4 *c*, 5, 6, & ? 9.)

Type-specimen [RR 1473], Geological Survey collection.

Diagnosis. Ventral valve: outline nearly semicircular, with lateral angles sharp, and inclined to be alate; greatest width at the posterior margin; convexity moderate, anterior and lateral slopes straight; apex marginal, pointed and somewhat erect; false area inclined a little forwards; pseudodeltidium triangular, gently convex, wide, angle made by its bounding sides about 120°, marked by two pairs of radiating angular lines so as to be polygonal in horizontal section; line of flexure from lateral slopes of shell to false area very abrupt and transverse to the axial line; free posterior margin of pseudodeltidium arched upwards; shell-substance corneous and very thick, built up of many laminae; surface marked by irregular concentric ridges which pass round the lateral angles to the false area, and are traceable across the pseudodeltidium.

Remarks.—There are a few radiating ridges on the central part of the test in one specimen [RR 1476]. Several specimens have the surface adventitiously indented in many places.

Dimensions in millimetres:—

	[RR 1473]	[RR 1470]	[RR 1471]
Length	?	6·0	5·5
Width	8·75	9·25	8·0
Height	1·6	1·4	1·4

The species is chiefly distinguished by its abrupt flexure-line, its polygonal pseudodeltidium (fig. 4 *b*), and by being widest at the posterior border.

For comparison with other species, see p. 331.

Locality and horizon.—Comley, Lower Cambrian; from the mottled (conglomeratic) and greenish-grey beds of the Road Quarry in the Cwms sub-area, horizon Ab₁.

Dorsal valve (fig. 9). The exfoliated test [RR 1465] has a very prominent vertical apex, and is regarded as a dorsal valve. It is found on the same piece of rock with an external cast referred to *M. phillipsi*, and may belong to that species; but its general outline is more suggestive of *M. rhodesi*.

MICROMITRA (PATERINA) KINGI, sp. nov. (Pl. XXI, figs. 7 & 8 *a*–8 *b*.)

Type-specimen [King 2], Geological Survey collection.

Diagnosis.—Ventral valve: outline posteriorly somewhat hexagonal, but anteriorly semicircular; length and width equal; widest at about the half-length, posterior border about two-thirds the greatest width, lateral angles well-marked but not sharp; rather strongly convex; false area slightly inclined forwards, pseudodeltidium triangular, convex, of moderate width, angle made by its bounding lines about 90° ; shell-substance corneous, not very thick; exterior marked with concentric ridges, which are irregular and in places present a beaded appearance.

Internally, or on an interior layer, there are numerous radiating striæ.

Remarks.—The concentric ridges appear to be rather coarser in this species than in the two previously noticed, but are only known from the marginal portions of the shell.

Dimensions in millimetres:—

	[King 2 h]	[King 2 g]
Length	5.75	?
Width	6.75	7.5
Width at posterior border	5.0	5.0
Height	2.0	?

The species is distinguished by its short posterior border, by its pseudodeltidium being nearly a right-angled triangle, and by the forward inclination of the false area.

Locality and horizon.—The Cwms sub-area of Comley, Lower Cambrian; from the *Obolella-groomi* Beds, horizon Ab.

These three species of *Micromitra* appear to be closely related. They occur together in the same beds, but in different numerical proportions. *M. kingi* has been found only in the mottled beds, *M. phillipsi* in the same beds and sparingly in the greenish-grey sandstones, *M. rhodesi* sparingly in the mottled beds and fairly abundantly in the greenish-grey sandstones.

They are clearly distinguished one from the other by their outlines and false areas. The surface-characters seem to differ a little in detail.

The divergences may be tabulated as follows:—

	<i>M. kingi.</i>	<i>M. phillipsi.</i>	<i>M. rhodesi.</i>
Apex	(Not known.)	Projecting backwards.	Approximately vertical.
Anterior slope.	Decidedly convex.	Slightly convex.	Straight.
False area ...	Inclined forwards.	Vertical.	Inclined forwards.
Angle at apex of pseudodeltidium }	about 90° .	about 120° .	about 120° .
Greatest width.	At half-length.	At quarter-length.	At posterior margin.
Postero-lateral angles.	Blunt.	Rounded.	Sharp.
Shell-substance.	Moderately thick.	Rather thick.	Very thick.

MICROMITRA (PATERINA) LABRADORICA (Billings). (Pl. XXI, figs. 10 a-13.)

Obolus labradoricus Billings, 1861, Geol. Surv. Canada, Pal. Foss. vol. i, p. 6. fig. 6.

Micromitra (P.) labradorica Billings, Walcott, 1912, p. 347 & pl. ii, figs. 2, 2 a-2 f.

(?) *Kutorgina cingulata* Billings, Callaway, 1878, Q. J. G. S. vol. xxxiv, p. 759.

(?) Do. do. do. Lapworth, 1888, Geol. Mag. p. 485.

(?) Do. do. do. Lapworth, 1891, *Ibid.* p. 532.

(?) *Kutorgina cingulata* Billings, Lapworth & Watts, 1894, Proc. Geol. Assoc. vol. xiii, p. 310.

Kutorgina and *Kutorgina cingulata* Cobbold, Brit. Assoc. Rep. 1909, p. 182; 1910 a, p. 118; 1912, pp. 138, 140, 141.

It seems probable that the specimens from Comley which have previously been referred to *K. cingulata* belong in reality to Billings's second species, now included by Dr. Walcott under the genus *Micromitra*, sub-genus *Paterina*.

The Comley specimens usually have a black outer surface: that is, covered with rounded concentric ridges, spaced, in the anterior part of the valves, at from 4 to 8 to the millimetre. Where exfoliation has taken place the internal laminae of shell are marked by radiating folds, which extend to distances that vary with the degrees of exfoliation.

Two specimens are known in which the dorsal and ventral valves are united. One of these [B 40306] in the British Museum (figs. 10 a-10 c) is now much damaged, and it is impossible to ascertain whether the apex is as much curved down at the back as is represented in the side view, the sketches for which were made before the other specimens were studied. It had the two valves slightly displaced one upon the other. In the other specimen, [730] in the collection for the British Association, the shells are somewhat crushed together, and they have suffered much from weathering.

The dimensions of five dorsal and three ventral valves are as follows, in millimetres:—

Nos. on specimens	Dorsal Valves.					Ventral Valves.		
	[B 40306]	[730]	[2188]	[2179]	[1519]	[B 40306]	[730]	[729]
Length	4.5	about 9	4.5	6.0	4.2	5.0	9.5	7
Width.....	6.5	about 10	7.5	8.5	5.7	6.5	11.0	10
Height	0.7	1.4	1.0	2.5	1.5	1.3	1.6	2

The concentric ridges of the exterior surface can often be traced round the postero-lateral angles of the shell on to the false area. In the dorsal valves the central part of the false area appears to have been widely open, no trace of a pseudochilidium having been detected. In the ventral valves the pseudodeltidium is occasionally seen, and takes the form of a very small, horizontal, semicircular shelf (fig. 13).

Through the kindness of Dr. Walcott, I have had an opportunity of examining several specimens of the species from Bic, Quebec. The only difference that has been detected is that the concentric ridges in the Canadian specimens are rather smaller and more numerous than those on shells from Comley of the same size.

Locality and horizons.—Comley, Lower Cambrian; from the *Callavia* Sandstone, *Olenellus* Limestone, *Microdiscus-bellimarginatus* Limestone, and *Protolenus* Limestone: horizons Ac_1 , Ac_2 , Ac_3 , & Ac_5 .

MICROMITRA (PATERINA) MINOR, sp. nov. (Pl. XXI, figs. 14–16.)

(?) *Iphidea bella* Billings, Lapworth, 1888, Geol. Mag. p. 485.

Iphidea (?) Cobbold, 1911, Rep. Brit. Assoc. (1910, Sheffield), p. 298.

A form of the genus that is much smaller and more convex than *M. (P.) labradorica*, is found associated with it in the *Olenellus* Limestone of Comley.

Type-specimen [755].

Diagnosis.—Ventral valve: outline subquadrate, wider than long, lateral angles slightly rounded; convexity strong, anterior and lateral slopes gently convex; apex fairly sharp, a little incurved, and overhanging the posterior border; shell-substance corneous, exfoliating in layers which mark stages of growth; surface marked with rounded concentric ridges usually very regularly disposed, but sometimes two or more appear to coalesce so as to form one ridge; they are continuous round the lateral angles on to the false area; false area with a wide, open delthyrium (fig. 14); pseudodeltidium either not present at all, or represented only by a minute projection under the apex, which has the appearance of being the posterior margin of the umbonal shield. Internal markings very minute. Three specimens, a little exfoliated at the apex, exhibit a more or less flattened space under the umbo; on the anterior slopes are traces of radiating corrugations, and also two rather stronger ridges that are slightly sinuous and divergent; in one of the specimens [750] (figs. 16 *a*–16 *c*) a slightly-elongated central prominence on the cast may be compared with the ‘apical slit’ mentioned by Dr. Walcott (1912), pp. 334, 335, 345.

Dorsal valve, not identified.

Dimensions.—Four ventral valves have the following measurements, in millimetres.

	[750]	[755]	[858]	[2178]
Length	1.25	2.0	2.1	1.5
Width	1.75	2.7	3.25	2.0
Height	0.5	1.0	1.1	0.6

Comparisons with other species.—*M. (P.) minor* suggests *M. (P.) bella*¹ (olim *Iphidea bella* Billings) at first sight; but the absence of any well-marked pseudodeltidium at once precludes this reference.

¹ C. D. Walcott, 1912, pl. ii, figs. 1 & 1 *a*.

A comparison may be made with *M. (P.) logani* Walcott,¹ which has a wide, open delthyrium and a small 'shelf-like' pseudo-deltidium, but differs in form from the Comley fossil.

M. (P.) stuarti Walcott² has an open delthyrium, but differs in outline and convexity from *M. (P.) minor*.

The Comley form has the same general outline as *M. (P.) undosa* Moberg,³ from drift blocks of Lower Cambrian Sandstone of Öland, but is much smaller, and the false areas cannot be compared, as the Swedish specimens are defective.

MICROMITRA MINOR, var. *GIBBOSA* nov. (Pl. XXI, fig. 17.)

A specimen in the collection made by Mr. Rhodes [RR 1117], and another in the British Association collection [2178] differ from those described above in having the anterior slope much more strongly convex; this gives a gibbous aspect to the shell.

Mr. Rhodes's specimen was obtained from the *Olenellus* Limestone of Comley Quarry; the exact horizon of the other specimen is, however, doubtful.

Localities and horizon.—Comley, Lower Cambrian; from the *Olenellus* Limestone of the Quarry, horizon Ac₂. Also from the same horizon in the Cwms, about 2 miles south of Comley Quarry.

MICROMITRA (?) sp. indet. (Pl. XXI, fig. 26.)

Associated with *M. (P.) phillipsi* and *Walcottina* two valves have been found, which were thought to be dorsal valves of *W. lapworthi* (see below, p. 336), and were figured for description. One of the figures is reproduced in Pl. XXI. There is a strong superficial resemblance to the figures given by Walcott⁴ of the dorsal valve of *Kutorgina granulata* Matthew, but the shells do not agree with the description. The umbo is not 'depressed to the cardinal line,' it stands up almost vertically, and there is no sign of 'a median septum.'

In the specimen figured the posterior portion is imperfect, but suggests a similarity to the false area of the ventral valves of *M. (P.) phillipsi* and its congeners.

Dimensions in millimetres.—Length = 9; width = 12; height = 1 or more.

Locality and horizon.—Comley, Lower Cambrian; from the Cwms sub-area, horizon Ab₁.

WALCOTTINA, gen. nov.

Two species of brachiopoda from the lowest fossiliferous horizon known in the Comley district appear to be intermediate between

¹ 1912, pl. ii, figs. 6, 6 a & 6 b.

² *Ibid.* p. 354 & text-figs. A, A', A'', B, B'.

³ 1892, p. 112 & pl. iii, figs. 10-12.

⁴ 1912, pl. v, figs. 5 b & 5 c.

the genera *Mickwitzia* and *Kutorgina*, to neither of which can they be referred without an extension of one or other of the generic diagnoses given by Dr. Walcott.¹ It is, therefore, necessary to propose a new generic name, and it seems appropriate to dedicate the genus to one who has done so much to aid the study of the Cambrian faunas in general and the brachiopoda in particular.

In common with the two above-named genera, *Walcottina* has the ventral valve considerably elevated, while the dorsal, so far as known, is comparatively flat; the pedicle-opening is a gaping orifice between the valves; and the shell-substance is made up of two or three principal layers.

The characters in which the three genera are divergent, so far as can be determined from the specimens of the new genus that are available, are as follows:—

	<i>Mickwitzia.</i>	<i>Walcottina.</i>	<i>Kutorgina.</i>
(1) Shell-substance ...	phosphatic.	calcareo-corneous.	calcareous.
(2) Exterior sculpture	papillose.	{ concentric striae } { with flat bands. }	{ concentric striae, imbricated. }
(3) Exterior surface of interior layer	{ radiate and con- centric ridgelets. }	radiate corrugations.	{ fine radiating li and puncta. }
(4) Innermost surface	punctate.	smooth.	?
Ventral valve.			
(5) False area	{ obscurely deve- loped. }	{ not marked off from remainder of shell. }	{ distinct, but sh and rudimenta }

No specimens of *Walcottina* have been found that are not more or less in a decomposed condition, and consequently the exact nature of the shell-substance cannot be ascertained. They occur in a somewhat friable conglomerate from which all, or almost all, the calcareous cement has been dissolved away, and with it, presumably, any calcareous material that the shells may have contained. The matter now representing the test, though soft, is often well preserved, its surface-characters can be made out with certainty, and the component layers are usually distinctly observable. They have a corneous aspect, but may have been originally calcareo-corneous. In this respect the specimens approach *Mickwitzia*, but the exterior sculpture and that of the interior layer are more reminiscent of *Kutorgina*.

The false area in *Walcottina* is non-existent, in *Mickwitzia* it 'may or may not be obscurely defined,'² in *Kutorgina* it is quite distinct, judging by the figures given by Walcott.³ In this respect the proposed genus seems to be more primitive than the other genera and to be on a par with *Rustella*.⁴

¹ 1912, pp. 329 & 579.

² C. D. Walcott, 1912, p. 329.

³ *Ibid.* pl. v, figs. 1 k & 1 n.

⁴ C. D. Walcott, *op. supra cit.* p. 327 & pl. i, figs. 1, 1 a-1 e.

Provisional diagnosis.—General outline either transverse or elongate oval. Ventral valve convex, with apex recurved over or beyond the posterior margin; anterior slope convex, with gently-lessening curvature, lateral slopes rather flatter; lateral angles rounded; posterior slope continuous with the lateral, without any defined false area; pedicle-opening indicated by an arching upwards of the posterior margin; shell-substance corneous or calcareo-corneous, consisting of three principal layers, of which the exterior is marked by numerous concentric impressed striæ with flat bands between them, the intermediate layer is marked by impressed lines radiating from the apex, and, towards the margins, by impressions of the concentric bands of the exterior, while the innermost layer is smooth so far as known.

Dorsal valve, not known with certainty.

Type species—*Walcottina lapworthi*, sp. nov.

Horizon—Lower Cambrian of the Comley district, Shropshire.

WALCOTTINA LAPWORTH, sp. nov. (Pl. XXI, figs. 23*a*–23*c*, 24*a*–24*b*, & 25.)

Cobbold, 1916. Rep. Brit. Assoc. 1915 (Manchester), p. 118, 'brachiopod allied to both *Mickwitzia* and *Kutorgina*'.

Type-specimens [King 1*a* & 1*b*] in the Geological Survey collection.

Diagnosis.—Ventral valve: outline transversely oval, lateral angles rounded, posterior margin nearly straight and transverse to the axial line for about three-fifths of the width; convexity well marked; apex somewhat recurved and extending well behind the posterior margin; anterior slope of fairly even curvature, though a little flatter towards the front margin; lateral slopes tending to flatness towards the lateral angles; false area and pseudodeltidium not developed; shell-substance corneous, composed of two principal layers of moderate thickness, and an inner one which appears to be very thin; exterior marked by concentric, impressed lines, spaced fairly regularly at 7 or 8 to the millimetre, the interspaces being of the nature of flat bands, stages of growth indicated by an occasional strengthening of the concentric lines; interior layer of shell marked by many gentle corrugations radiately arranged, and dying away towards the half-length of the shell; third and innermost layer smooth, so far as seen.

A comparison between the external view (fig. 23) and that of the interior layer (fig. 24) shows that the outer layer of shell was thickened round the base of the apical portion, so that the apex in the latter figure is more acuminate than it is in the former.

Dorsal valve, not known.

The type-specimens were collected by Mr. W. Wickham King, and presented by him to the collection which was being made in 1892 by Mr. J. Rhodes.

The species is dedicated to Prof. C. Lapworth, who suggested a reference to *Mickwitzia* in the field at the time when the specimens were found.

Dimensions of type-specimens.—Lengths=5 and 7·5 mm.; widths = 7·1 and 9·25 mm.; heights = 2 and 2·5 mm.

Locality and horizon.—The Cwms sub-area of Comley; from the conglomeratic beds at the base of the *Obolella-groomi* Beds, horizon Ab₁.

WALCOTTINA ELEVATA, sp. nov. (Pl. XXI, figs. 20 *a-e*, & 22.)

Cobbold, 1916. Rep. Brit. Assoc. 1915 (Manchester), p. 118, 'brachiopod allied to both *Mickwitzia* and *Kutorgina*.'

The type-specimens consist of an exfoliated fragment [1914] disclosing about half of a ventral valve and its counterpart [1916]. Another specimen [2282], a sandy internal cast of a complete shell in rotten stone, is used to supplement the description.

Diagnosis.—Ventral valve: general form elevated; outline an elongate oval, slightly flattened at the posterior margin, length rather greater than width; lateral angles well rounded; convexity high, highest point situated at about a quarter of the length from the apex; apex recurved, and projecting but little beyond the posterior margin; shell-substance corneous, and composed of three principal layers, marked as in *W. lapworthi*, except that the radiating corrugations of the inner layer do not extend so far forwards, and that traces of the external concentric bands are seen upon the interior layer near the anterior margin; pedicle-opening represented by an upward arching of the posterior margin (see Pl. XXI, fig. 22).

Dimensions of the type-specimen [1914].—Length=4 mm., width=4 mm., height=3·75 mm.; and of specimen [2282], length=7 mm., width=6·9 mm., height=4·5 mm.

Dorsal valve (?). Associated on the same little piece of rock with the type-specimen are two shells which from their general form are regarded as dorsal valves, but the surfaces are not sufficiently well preserved to prove that the sculpture was similar to that above described for the ventral valve.

The outline is that of a sector of a circle greater than a semi-circle, the two radii forming a notched posterior margin; the apex projects prominently upwards; the anterior and lateral slopes fall steeply from it, then become horizontal or even concave, and finally fall again to the margin; the posterior angles are rounded; the surfaces show concentric lineation, and the dimensions are such as to correspond with those of the type-specimen.

Observations.—The attenuation of the apex of the ventral valve (fig. 20 *a*) [1914] suggests a possible pedicle-tube, especially when compared with figs. 20 *b* & 20 *c*, which represent the exterior of the same individual, but the former is a representation of the interior layer of shell, not an internal cast. The exterior layer of shell was evidently considerably thickened at the apical part of the valve. The arching upwards of the posterior margin shown in fig. 22 [2282] indicates the more probable position of the pedicle-opening.

This species is distinguished from *Walcottina lapworthi* by the

greater elevation of the ventral valve, and by being slightly elongate instead of decidedly transverse.

Locality and horizon.—The Cwns sub-area of Comley, Lower Cambrian; from the *Obolella-groomi* Beds, horizon Ab₁.

OBOLIDÆ King.

OBOLUS Eichwald.

OBOLUS PARVULUS, sp. nov. (Pl. XXII, figs. 13–19.)

Type-specimen [1512 a].

Diagnosis.—General form ovate, bluntly acuminate, apices marginal; dorsal valve, length and width sub-equal; ventral valve longer than wide; convexity moderate, about a sixth of the width; shell, apparently calcareo-corneous, consisting of two principal layers, weathering white or bluish white; exterior with from three to six well-defined lines of growth, numerous faint concentric striae and occasional traces of radiating lines, in the dorsal valve there is an incised mesial line; interior of ventral valve showing the pedicle-groove and, continuous with it (though marked off by a transverse line) the concave visceral area, on each side of which are traces of the main vascular sinuses: interior of dorsal valve showing a narrow but well-marked mesial ridge, which near the middle of the valve gives off outward curving branches on either side, the mesial ridge widens out posteriorly and, in the casts, is lost in the impression of the area.

Dimensions.—Nineteen valves measured from .9 to 2.35 millimetres in length; the average proportion of length to width in ventral valves is 1 to .8, in dorsal valves 1 to .9.

The ventral valve of *O. parvulus* is very like *O. parvus* Walcott¹ from the Lower Cambrian of the Rocky Mountain province.

The dorsal valve of the Comley species differs in being longer than wide, and the mesial depression is much narrower than that shown in Walcott's figure.

The surface-character of *O. parvus*, said to be like that of *Lingulella* (*Lingulepis*) *longinervis* Matthew [(?) a clerical error for *L. (L.) gregwa* Matthew] has not been observed in *O. parvulus*.

Locality and horizon.—Comley, Lower Cambrian; from the *Olenellus* Limestone, Horizon Ac, the *Microdiscus-bellimarginatus* Limestone Ac₂, and the *Protolenus* Limestone Ac₃.

OBOLUS (?) LINNARSSONI, sp. nov. (Pl. XXII, figs. 23–25.)

Obolus (?) sp. indet., Linnarsson, 1876, *Bihang K. Svensk. Akad. Handl.* vol. iii, No. 12, p. 16 & pl. iii, fig. 31.

(?) *Lingulella* sp. indet., Matley, 1911, *Q. J. G. S. vol.* lxxii, p. 301 & pl. xxvi fig. 7.

Obolus sp. indet. *g.*, Walcott, 1912, *Monogr. U.S. Geol. Surv.* vol. li, p. 423.

Linnarsson's description is as follows:—

'The shell is very slightly convex; its circumference resembles a sector of

¹ 1912, p. 408, text-figs. 37 A & 37 B.

a circle; the front is broadly rounded, the slopes forming the beak seem to be nearly straight, and make a somewhat obtuse angle. The surface is not well preserved in any specimen; judging, however, from some small fragments, I should think that a concentric striation is prevailing, but there are also radiating ribs. In the inner shell-layers little more than longitudinal striation is seen: the innermost are perforated by small punctiform pits. The inner layers are polished and glossy, the outermost seems to be opaque.

Except that the Shropshire specimens do not reveal the punctiform pits on the innermost layer and that the radiating ribs are not seen, the above description exactly applies to the two Comley shells, both of which were found associated with *Billingsella lindstrœmi* and *Acrothele coriacea*; Linnarsson's specimens were obtained from 'loose masses' of bituminous limestone belonging to the strata with *Paradoxides forchhammeri*, and associated with the same two species and also '*Orthis exprorecta*.' For some reason that is not explained, Dr. Walcott¹ refers these blocks to the *Paradoxides-œlandicus* Zone.

Neither of the two Comley specimens exhibits the interior surface, and only a very small portion of the exterior is seen in one of them (see fig. 23 *a*). This is near the umbo, and shows concentric striations, but no trace of radiating ribs. In both specimens the shell is thick, and consists of several layers. In one of them are two raised bands (see fig. 23 *b*) on one of the inner layers diverging at an acute angle from the apex, recalling the 'flanges' noted by Dr. Matley² in a fragmentary specimen that he referred with some hesitation to *Lingulella*. A minute shell (fig. 25) possibly represents a young form of the species.

Locality and horizon.—Comley, Middle Cambrian; from the *Billingsella* Beds, horizon Bc.

OBOLUS (?) GIBBOSUS, sp. nov. (Pl. XXII, figs. 20 & 21.)

Type-specimen [2119].

Diagnosis.—Ventral valve: outline a wide oval, from which the apex extends well back beyond the posterior margin, so that the shell is somewhat pear-shaped; convexity very strong and fairly even in curvature, apex curved down to a height above the marginal plane of the valve that is less than half the maximum height of the valve; area more or less concave, inclined at an angle of about 45°; shell thick, marked by a few raised radiating lines, crossed by concentric bands of varying depths of colour, and also by the ragged edges of successive layers of shell-material.

Interior without marks, so far as known from internal casts.

Dorsal valve: outline oval, narrowing slightly posteriorly, strongly convex, curvature flattening a little anteriorly; surface marked by several ragged edges of shell-material; interior with a slight oval prominence under the umbo, on either side of which lie two ill-defined concavities; otherwise devoid of vascular marks.

¹ 1912, p. 424.

² 1911, p. 301 & pl. xxvi, fig. 7.

Dimensions in millimetres.

Ventral valves	[2119]...	Length=3.5.	Width=3.0.	Height=1.1.
	[677]...	Length=2.5.	Width=2.0.	Height=?
Dorsal valves	[2120]...	Length=2.8.	Width=2.6.	Height=0.66.
	[676]...	Length=2.5.	Width=2.25.	Height=?

Locality and horizon.—Comley, Middle Cambrian; from the *Billingsella* Beds, horizon Bc.

OBOLUS (?) cf. SCHMALENSEEI Walcott. (Pl. XXII, figs. 22 *a* & 22 *b*.)

Obolus schmalenseei Walcott, 1912, U.S. Geol. Surv. Monogr. vol. li, p. 412 & pl. xxx, figs. 19, 19 *a*–19 *e*.

Lingulella (?) spp. *pars*, Matley, 1911, Q.J.G.S. vol. lxxvii, p. 301 & pl. xxvi, figs. 8 *a* & 8 *b*.

Additional specimens possessing the sub-pentagonal form described by Dr. Matley have come to light since he wrote his notes on the brachiopoda from the *Paradoxides* Beds of Comley.

In size and general form they are very like *O. schmalenseei*, but they lack the fairly strong internal marks shown for that species in Walcott's figures. In the dorsal valve there is a central prominence and also two ill-defined concavities under the umbo as in *O. gibbosus*.

The specimen figured with Dr. Matley's notes is there called the interior of a brachial valve. From the lateral view there figured it seems more like a ventral.

The dorsal valve now figured [689] has the length and width equal, the convexity is strong, the test consists of several layers, the exterior is marked by the concentric edges of laminae, and the interior by an ill-defined central ridge extending to a quarter or a third of the length of the shell, with ill-defined hollows on each side under the umbo.

Length=3 mm.

Locality and horizon.—Comley, Middle Cambrian; from the *Billingsella* Beds, horizon Bc.

OBOLUS (?) sp. indet. *a*. (Pl. XXIII, fig. 29.)

A single valve [1604], that is referred to this genus on account of its shape, was found with crushed specimens of *Acrotreta* and fragments of *Hyolithellus micans* in fine blue shale, which is mapped as belonging to Horizon Ba₂.

Dimensions.—Length=2.2 mm.; width=2 mm.

OBOLUS (?) sp. indet. *b*. (Pl. XXIII, fig. 28.)

A rather large single valve [162 *b*], also referred to the genus, was found associated with *Acrotreta* in sandy shale mapped as belonging to Horizon Bb₂.

The specimen is remarkable for its strong convexity and the wide flattened margin of the cast, indicating a thick shell.

Dimensions.—Length=3.2 mm.; width=2.7 mm.

LINGULELLA Salter.

LINGULELLA VIRIDIS, sp. nov. (Pl. XXII, figs. 10–12.)

Type-specimen, a dorsal valve [97].

Diagnosis.—General form ovate, a little acuminate posteriorly; apical angle in the dorsal valve about 60° , in the ventral slightly less; the two valves are about equally convex; area, in the dorsal valve, apparently in the same plane as the margin of the valve, in the ventral rising at a gentle angle.

Exterior of both valves marked with from three to six principal lines of growth, showing numerous, very fine, concentric striae between them, also many radiating striae only visible under a strong magnifier.

Interior of dorsal valve (only known from casts) with two gently convex, curved ridges, extending over the middle third of the length of the valve, convex towards the centre-line and tapering to points anteriorly; within these ridges and back towards the apex the surface is roughened by minute projections, outside them it is smooth and polished; faint indications of linear marks are sometimes visible in the roughened area, but no muscle-scars have been detected.

Area gently convex and, in the dorsal valve, separated by a very narrow space from the umbonal portion of the shell; marked by three to six sinuous ridges that are strong as compared with the concentric striae on the body of the shell.

Dimensions.—The type-specimen has a length of 5 mm.; more usually the shells are from 2 to 3 mm. in length. The convexity is about $\cdot 2$ of the length, and the width $\cdot 8$ in dorsal valves and $\cdot 7$ in ventral valves.

Observations.—There is no good interior of the ventral valve. Fig. 10 shows an internal cast from which the shell-substance has been in part cleared away.

In form and external characters the species seems near to *L. nathorsti* Linnarsson; but the internal marks of that species, so far as known, and the details of the area of the dorsal valve are very different.

Locality and horizon.—Comley, Lower Cambrian; from the *Strenuella* Limestone, horizon Ac_1 , and sparingly from the *Protolenus* Limestone, horizon Ac_3 .

LINGULELLA NICHOLSONI Callaway.

Lingulella nicholsoni Callaway, 1874, Q. J. G. S. vol. xxx, p. 196.

Lingulella nicholsoni Callaway, 1877, Q. J. G. S. vol. xxxiii, p. 688 & pl. xxiv, figs. 11, 11 a–11 b.

Lingulella nicholsoni (Callaway), Walcott, 1912, U.S. Geol. Surv. Monogr. vol. li, p. 522 & pl. xxx, figs 3, 3 a–3 f.

Dr. Callaway records¹ the discovery of this species, together with 'the Shineton graptolites,' in shales east of Lawley Hill.

¹ 1879, p. 759.

The area 'east of Lawley Hill' is a continuation of the Comley Cambrian area—the stratigraphical relations to the beds which I have studied are unknown. The horizon is, therefore, lettered Cx.

LINGULELLA cf. FERRUGINEA Salter.

Cf. *Lingulella ferruginea* Salter, 1867, Q. J. G. S. vol. xxiii, p. 340, fig. 1.

Cf. *Lingulella ferruginea* Salter, Walcott, 1912, U.S. Geol. Surv. Monogr. vol. li, p. 496; pl. xxix, figs. 1, 1*a*-1*w*, 2, 2*a*-2*f*; pl. xxx, fig. 1; pl. xxxi, figs. 3, 3*a*-3*c*; and pl. xxxv, figs. 4, 4*a*-4*b*.

Lingulella cf. ferruginea Salter, Matley, 1911, Q. J. G. S. vol. lxxvii, p. 300 & pl. xxvi, figs. 5-6*b*.

No further evidence as to this form has come to light since Dr. Matley's notes were written.

Locality and horizon.—Comley, Middle Cambrian; from the *Paradoxides-rugulosus* Sandstone and from the *P. davidis* Flags, horizons Bb₁ & Bb₂.

LINGULELLA sp. indet.

Lingulella sp. indet., Matley, 1911, Q. J. G. S. vol. lxxvii, p. 301, *pars*.

Dr. Matley calls attention to another 'small oval form' found in the *Paradoxides-rugulosus* Sandstone, horizon Bb₁.

OBOLELLIDÆ Walcott & Schuchert.

OBOLELLA Billings.

OBOLELLA ATLANTICA Walcott.

Obolella atlantica Walcott, 1890, Proc. U.S. Nat. Mus. for 1889, vol. xii, p. 36.

Obolella atlantica Walcott, 1912, U.S. Geol. Surv. Monogr. vol. li, p. 589, text-fig. 61 & pl. lv, figs. 1, 1*a*-1*i*.

Obolella atlantica is represented at Comley by two forms, which are readily distinguished one from the other, but are so closely allied to Walcott's species that they are described here as varieties.

OBOLELLA ATLANTICA, var. COMLEYENSIS nov. (Pl. XXII, figs. 6-9; text-fig. 1.)

This is the small, round, whitish brachiopod so frequently met with in the *Olenellus* Limestone of the Comley Quarry. Usually the specimens are very poorly preserved, but occasionally good internal casts and exteriors are to be found.

The two valves are nearly circular, and have the same degree of convexity. The exterior answers closely to the description and figures given by Walcott. The interior differs a little from them in the development of the vascular marks; the size is uniformly smaller; and the transverse diameter is always a little greater than the length.

Dimensions.—Length = 3.5 to 3.8 mm.; width = 3.7 to 4.0 mm.; height = .7 to .9 mm.

Observations.—In well-preserved casts of the ventral valve (figs. 6 & 9 & text-fig. 1), the cast of the pedicle-tube is usually

visible, and points upwards and backwards at an angle with the plane of the valve-margin; on each side of it obtuse prominences project horizontally backwards from the convex body of the cast; at about a third of the length of the valve from the apex two comma-shaped prominences are situated, having their points directed forwards and nearly meeting at the centre-line; beyond the swollen ends of the commas, ridges extend forwards and outwards, and between them and the centre-line, other ridges, one on each side, may sometimes be detected running in parallel directions; traces of a backward prolongation of the outermost pair of ridges are occasionally observed (see Pl. XXII, fig. 9).

In the cast of the dorsal valve (Pl. XXII, fig. 7 & text-fig. 1) a central depression usually extends from the posterior border to about

Fig. 1.—*Obolella atlantica*, var. *comleyensis* nov. $\times 10$. *Diagrams of internal casts of the ventral and dorsal valves, showing all marks observed in the several specimens.*

half the length of the valve; on each side raised lines diverge forwards, and are traceable for about the same distance; from the back of the convex body of the cast obtuse prominences project horizontally backwards, similar to those of the ventral valve but nearer to the centre-line; beyond these the body of the cast is hollowed out, the hollows being terminated laterally by projecting elongated bosses, more or less parallel to the margin of the valve.

In both valves a wide flattened or even recurved space extends all round the convex body of the cast.

All the internal marks that have been observed in various specimens have been combined in the outline restoration (fig. 1, above).

The shell-substance is calcareous.

Dr. J. Kiær, in his work on the *Holmia* fauna of Tönten, describes and figures a species, *O. rotundata*, that is, at first sight, very like the variety *comleyensis*. The general form is almost exactly the same, and the resemblance between the dorsal valves is particularly striking.

The Tönten shell is rather more pointed posteriorly than the Comley form; it is about 25 per cent. larger in linear dimensions;

the vascular sinuses of the ventral valve are usually, but not invariably, bent out widely so as to follow a curve that is sub-parallel to the margin, instead of being straight; the pedicle-tube appears to be stouter and more triangular.

From Dr. Kiær's figures the shell appears to have been thick, and to have the retroussé margin that is so characteristic of the Comley shells, but possibly this similarity may be due to its preservation in shale.

The variety *comleyensis* is nearer to the American form of *O. atlantica* (that is, *O. rotundata*), but not so near as the variety *transversa*.

Locality and horizon.—Comley, Lower Cambrian; from the *Olenellus* Limestone, horizon Ac₂, and from the *Microdiscus-bellimarginatus* Limestone, horizon Ac₃.

OBOLELLA ATLANTICA, var. *TRANSVERSA* nov. (Pl. XXII, figs. 1-5; text-fig. 2.)

This variety occurs at a higher horizon than *comleyensis*, and always in association with a different group of trilobites. In

Fig. 2.—*Obolella atlantica*, var. *transversa* nov. $\times 5$. *Diagrams of internal casts of the ventral and dorsal valves, showing all marks observed in the several specimens.*

point of size it is nearer to the American form, but differs in having both valves more transverse and also in the details of the internal marks. From the variety *comleyensis* it differs in general shape and size; in having a relatively thinner shell; in the external surface, which is more in accord with Walcott's description; and in the internal features, which are proportionately smaller. Though a larger shell, it appears to be built on more delicate lines.

Dimensions.—The lengths observed vary from 5.3 to 6.5 millimetres, the widths are from an eighth to a tenth greater, the convexity averages about a fifth of the length.

All the internal marks shown on several casts have been combined in text-fig. 2. In order that the internal characters of the

two varieties may be readily compared, the scale of enlargement used in fig. 2 is half of that used in fig. 1 (p. 343).

The exterior is often beautifully preserved, 'the fine radiating striae that are sometimes raised lines'¹ are particularly well marked (Pl. XXII, fig. 5).

Locality and horizon.—Comley, Lower Cambrian, from the *Strenuella* Limestone, horizon Ac₄.

OBOLELLA (?) GROOMII Matley. (Pl. XXI, fig. 27.)

Obolella (?) *groomii* Matley, 1902, Q. J. G. S. vol. lviii, p. 137 & text-figs. 1-2.
Obolella groomii Matley, Walcott, 1912, U.S. Geol. Surv. Monogr. vol. li, p. 595
& text-figs. 52 A-52 B.

Several examples of this species have been found in the mottled and greenish-grey beds at the base of the Lower Comley Sandstone, at the Road Quarry in the Cwms. Fragments are readily identified by the external characters, the edges of successive lines of growth being very sharp, and almost 'undercut' in their sculpture. The shell-substance is well preserved in the sandstone, and appears to be corneous. It is doubtful whether it was ever to any considerable extent composed of carbonate of lime, for the shells are not preserved as empty casts notwithstanding the open, sandy nature of the rock.

As with the Malvern specimens, dorsal and ventral valves cannot be distinguished with any certainty, and no interiors have been met with. In view of these facts the generic reference must remain doubtful.

Locality and horizon.—Comley, the Cwms sub-area; Lower Cambrian, horizon Ab₁.

ACROTRETIDÆ Schuchert.

ACROTHELE Linnarsson.

ACROTHELE CORIACEA Linnarsson. (Pl. XXIII, figs. 19 a-22 b.)

Acrothele coriacea Linnarsson, 1876, Bihang K. Svenska Vetensk.-Akad. Handl. vol. iii, No. 12, p. 21 & pl. iv, figs. 44-48.

Acrothele cf. *granulata* Linnarsson, Matley, 1911, Q. J. G. S. vol. lxvii, p. 302 & pl. xxvi, figs. 9 a-9 b.

Acrothele coriacea Linnarsson, Walcott, 1912, U.S. Geol. Surv. Monogr. vol. li, p. 642 & pl. lvi, figs. 1, 1 a-1 i.

The specimens submitted to Dr. C. A. Matley were not wholly satisfactory; he noticed, however, that the position of the apex of the ventral valve was more in accord with that of *A. coriacea* than with that of *A. (Redlichella) granulata*. Since his notes were written, Dr. C. D. Walcott's work has appeared, with details as to the aspect of the granulations of both species.

A number of additional specimens have now come to light from the *Billingsella* Bed of Caradoc Dingle, and these agree in all respects with the published descriptions of *A. coriacea*. The shells usually split along one of the inner layers, which are bright and glistening. The exterior surface of the ventral valve near the

¹ C. D. Walcott, 1912, p. 589.

margins, and on the greater part of the false area, is covered with rugose concentric ridges, which give place to granulations on the central parts of the shell.

Of the dorsal valve Dr. Walcott (*op. cit.* p. 642) says:—

‘On some shells a slight depression or flattening extends (from the apex) to about the middle of the valve.’

In the Comley specimens this feature is seen in exfoliated dorsal shells (fig. 22 *a*). If, however, the successive laminae are removed, the depression becomes increasingly distinct, and, when the surface of the internal cast is reached, there is a distinct central hollow indicating a well-developed ridge or septum on the interior of the shell reaching nearly as far as the anterior border (fig. 22 *b*).

The characters of the exterior of the two valves are somewhat dissimilar. The dorsal valve seems to be covered with rugose concentric ridges, like those on the margins and false area of the ventral.

Dimensions.—Greatest length observed = 9 mm.; the usual length = 5 to 6 mm.

Locality and horizon.—Comley, Middle Cambrian; from the *Billingsella* Beds, horizon Bc; from the *Paradoxides-davidis* Flags, horizon Bb₂; and from the *Orusia* Shales, horizon Ca.

ACROTRETA Kutorga.

ACROTRETA SAGITTALIS Salter.

Obolella sagittalis Salter, 1866, Rep. Brit. Assoc. (1865) p. 285.

Acrotreta (*Linnaarssonina*) *sagittalis* Salter, Matley, 1911, Q. J. G. S., vol. lxxvii, p. 302 & pl. xxvi, figs. 10 *a*–10 *b*.

Acrotreta sagittalis Salter, Walcott, 1912, U.S. Geol. Surv. Monogr. vol. li, p. 704 & pl. lxxi, figs. 2, 2 *a*–2 *h*, 3, 3 *a*–3 *j*.

A few valves that are referred to this species occur in association with *A. schmalenseei* var. at the two sections on the Shoot-Rough Road, and in the *Billingsella* Bed of Caradoc Dingle, horizon Bc.

ACROTRETA SCHMALENSEEI Walcott, var. MATLEYI nov. (Pl. XXII, figs. 26–30, 32, & (?) 31.)

Acrotreta aff. *socialis* Seebach, Matley, 1911, Q. J. G. S. vol. lxxvii, p. 303 & pl. xxvi, figs. 11 *a*–13 *b*.

Acrotreta schmalenseei Walcott, 1912, U.S. Geol. Surv. Monogr. vol. li, p. 709 & pl. lxx, figs. 1, 1 *a*–1 *s*.

The study of about a hundred additional examples of the Comley form of this species, which have been obtained from the original rock-bed since Dr. Matley wrote his notes, shows that the ventral valve is persistently different in form from that depicted in Dr. Walcott's figures; in the absence of direct comparison with Scandinavian specimens, it seems therefore advisable to refer to them under a varietal name.

In the specimens examined by Matley, the anterior margins of the ventral valves were never satisfactorily cleared of matrix; with the new material available it has been possible to do this in many instances, and to establish the fact that the apices are always

situated over or behind the posterior margin of the opening of the valves, never in front of it, and that the anterior slope is distinctly convex. In all other respects the shells accord with Walcott's figures and description.

Dimensions.—The lengths observed for ventral valves vary from 1 to 3 mm.; the width is 1.25 of the length, and the height is about .5 of the length.

Locality and horizon.—Comley, Middle Cambrian; from the *Billingsella* Beds, horizon Bc, also doubtfully from the *Paradoxides-davidis* Flags, horizon Bb.

ACROTRETA SOCIALIS Seebach (?). (Pl. XXII, fig. 35.)

Acrotreta socialis Seebach, 1865, Zeitschr. Deutsch. Geol. Gesellsch. vol. xvii, p. 341 & pl. viii a, figs. 1-4.

Acrotreta socialis Seebach, Walcott, 1912, U.S. Geol. Surv. Monogr. vol. li, p. 711 & pl. lxxii, figs. 3, 3 a-3 c, 4, 4 a-4 c.

Some small ventral valves, found associated with *Acrothyra comleyensis* and *Dorypyge lakei*, agree with some of the figures of this species given by Dr. Walcott.¹ The anterior and posterior slopes are nearly straight, the apex is well forward from the posterior margin, the foramen is distinctly visible behind it, the height is about a half of the length, the surface is striated with fine concentric lines, the false area is moderately well-defined, and the vertical groove is narrow.

Dimensions.—Length=2 mm., width=2 mm., height=.8 mm.

These shells differ from Walcott's description of Seebach's species in being rather smaller, in having finer concentric striae, and in the narrower vertical groove.

From *A. schmalenseei* they differ very decidedly in the position of the apex and in the anterior slope being almost straight, instead of curved.

Locality and horizon.—Comley, Middle Cambrian; from the *Dorypyge-lakei* Flags, Ba.

ACROTRETA sp. indet. a. (Pl. XXII, fig. 33.)

A few valves have been found in association with *A. schmalenseei* var. *matleyi*, which appear to be distinct from any described species, but are too scarce or indifferently preserved to be definitely named.

They are distinguished by (1) the ventral valve being very elevated (it is almost as high as it is long); (2) the straight anterior slopes; (3) the false area being inclined backwards; (4) the absence of the vertical groove on the false area in the only exterior known.

The exterior is marked by fine concentric lines, set more widely apart than in *Acrotreta schmalenseei*.

Dimensions.—Length=1.5 mm.; width=1.5 mm.; height=1.25 mm.

¹ 1912, pl. lxxiii, figs. 4, 4', 4'', 4 a, 4 a', 4 a''.

Locality and horizon.—Comley, Middle Cambrian; from the *Billingsella* Beds, horizon Bc.

ACROTRETA sp. indet. *b*. (Pl. XXIII, fig. 30.)

A few valves were found in fine blue shale in the bed of Comley Brook; all are much crushed. Fig. 30 [1606] represents one of the best-preserved ventral valves.

Dimensions.—Length=1·2 mm.; width=1·4 mm.; height, greater than 4 mm.

Locality and horizon.—Comley, Middle Cambrian; horizon Ba₂.

ACROTRETA sp. indet. *c*. (Pl. XXII, fig. 34.)

Two dorsal valves [1621 *a*, 1622] that appear to belong to the genus were found in sandy shale east of Hill-House Ridge.

The internal characters are well marked, but there is not sufficient material for specific determination.

Dimensions in millimetres.

1621 *a*. Length=2·4, width=2·5.

1622. Length=1·5, width=1·7.

Locality and horizon.—Comley, Middle Cambrian; horizon Bb₂.

ACROTHYRA Matthew.

Dr. G. F. Matthew¹ proposed the generic name *Acrothyra* for certain *Acrotreta*-like forms having

‘the ventral valve elongate-conical, with the apex either overhanging the cardinal line, or but little in front of it; [and the interior] with a long, narrow, or a quadrate visceral callus, extending forward from the foramen about a third of the length of the valve and widening as it goes.’

Dr. C. D. Walcott,² in his monograph, adopts the separate generic term; but he observes that

‘this genus is intimately related to *Acrotreta*, and, except for the presence in some species of a well-marked visceral area of a type not known in *Acrotreta*, there would be difficulty in assigning more than subgeneric value to it.’

The Comley specimens that are here assigned to Matthew’s genus are very like *Acrotreta*; but they have a strongly-marked visceral area in the ventral valve, and in the dorsal valves the main vascular sinuses are well developed.

ACROTHYRA COMLEYENSIS, sp. nov. (Pl. XXIII, figs. 1 *a*–*S*;
text-fig. 3, p. 349.)

Acrothyra sp., Cobbold, 1916, Rep. Brit. Assoc. (Manchester, 1915) p. 119.

Type-specimen [1710].

Diagnosis.—Ventral valve subcircular, conical; apex pointed, and somewhat in advance of the posterior margin; false area ill-defined, traversed by a broad vertical channel and inclined forwards

¹ Bull. Nat. Hist. New Brunswick, vol. iv (1901) pt. 4, No. 19, p. 303.

² 1912, U.S. Geol. Surv. Monogr. vol. li, p. 716.

at an angle of 70° to 80° ; interior showing a bell-shaped cavity connected posteriorly with the foramen by a short groove in the wall of the cavity; main vascular sinuses deep on each side of the cavity, then disappearing, except in a few instances when they are traceable a long way towards the anterior margin; cardinal scars large; parietal band sinuous, connecting the forward ends of the

Fig. 3.—*Acrothyra comleyensis*, *sp. nov.* $\times$ about 24. *Restoration of the interior of the ventral valve.*

[*f* = Foramen; *p* = Minute pit; *b.c.* = Bell-shaped cavity; *v.s.* = Vascular sinus; *M.* = Cardinal muscle-scar; *p.b.* = Parietal band; *c.m.* = Central muscle-scar (?); *a.l.m.* = Anterior lateral muscle-scar (?).]

deeper parts of the vascular sinuses and passing the centre-line at about the half-length; muscle-scars (?) represented by four slightly-raised prominences almost touching the parietal band; two little pits occur in the wall of the bell-shaped cavity on each side of the foramen and a little in advance of it. Associated dorsal valves subcircular, convex, with apex depressed and marginal; interior with a median ridge extending about two-thirds the length of the valve, sometimes interrupted, sometimes only weaker at the middle of its course, from which point two branches diverge slightly and then pass forwards parallel to the median ridge; main vascular sinuses well developed posteriorly; cardinal muscle-scars large.

Shell-substance in both valves corneous, thick, and formed of many laminae; exterior marked by very closely-set concentric striae; inner layers occasionally showing faint traces of radiating striae.

Dimensions.—Lengths=from 1·9 to 3·25 mm.; proportion of length to width=about 1 to 1·1; of length to height, in ventral valves=about 1 to ·33, in dorsal valves=1 to ·2.

Observations.—The internal characters seen in the casts vary so much in size, in proportion to the diameters, as to suggest that more than one species may be present. These divergences are, however, due to the amount of shell remaining on the cast, and disappear on its removal.

In fig. 2, Pl. XXIII, representing an internal cast of the ventral valve from which all the shell-substance has come away, the bell-shaped cavity extends to about a third of the length of the shell; fig. 3 represents a cast with some of the shell left on it, and here the cavity extends to about a fifth of the shell-length; in fig. 4, from a cast to which much shell adheres, the extension is but to a twelfth, and in this case the visceral cavity has the circular form seen in the young shell. Figs. 1 *a* to 1 *e* are five views of the apex of one ventral valve, from which four successive layers of shell were removed. In fig. 1 *a* the exterior of the shell is seen; in 1 *b*, after the first exfoliation, the visceral cavity is represented by the circular prominence, while in the succeeding figures the cavity assumes its more normal shape. The various forms are evidently the internal casts at successive stages of growth.

Variation.—Occasionally a full-sized shell is found, in which the cavity retains somewhat of the circular shape seen in the young shell (fig. 7, Pl. XXIII).

The relations of *Acrothyra comleyensis* to other species are discussed with those of *A. cf. sera* (below). It may, however, be remarked here that indications of the two little pits near the foramen may be seen in the figures of the following species in Dr. C. D. Walcott's monograph:—

Acrotreta definita Walcott, pl. lxiv, figs. 2 *c*, 2 *c'*.

Acrotreta kutorgai Walcott, pl. lxy, figs. 36, 36', 3 *c*, 3 *c'*.

Acrotreta argenta Walcott, pl. lxvii, figs. 5 *a*, 5 *b*, 5 *b'*.

They are also seen in *A. cf. sera*; see Pl. XXIII, fig. 10, of this paper.

Locality and horizon.—Comley, Middle Cambrian. *A. comleyensis* is found associated with *Dorypyge lakei* Cobbold, at two local horizons, Ba₁ and Ba₃; in the former, fragments of *Paradoxides* have also been found.

ACROTHYRA cf. SERA Matthew. (Pl. XXIII, figs. 9 *a*–18.)

Acrothyra signata sera Matthew, 1902, Bull. Nat. Hist. Soc. New Brunswick, vol. iv, pt. 5, No. 20, pp. 383, 384 & pl. xiii, figs. 3 *a*–3 *e*.

Acrothyra sera Matthew, Walcott, 1912, U.S. Geol. Surv. Monogr. vol. li, p. 718 & pl. lxxx, figs. 4, 4 *a*–4 *c*, 5, 5 *a*–5 *d*, 6, 6 *a*–6 *i*, 7, 7 *a*–7 *b*, 8 (?). Reasons are given for grouping several of Dr. Matthew's species or varieties under one specific name.

A form which is very close to, if not identical with, the Cape Breton species, occurs in considerable quantity in the granular

portion of the Black Limestone of Comley, now called the *Lapworthella* Limestone.

The ventral valve is conical and nearly circular in outline, somewhat flattened posteriorly, and has a very slight indentation at the middle line; the posterior slope may be vertical (fig. 12), or inclined backwards (fig. 11), or a little forwards (fig. 9 *a*); the anterior and lateral slopes are nearly straight; the false area is ill-defined and traversed by a vertical furrow, indistinct towards the margin. The internal marks vary with the amount of exfoliation. Fig. 9 *a* shows the cast of an interior where slight exfoliation has occurred; fig. 9 *b* represents the same specimen after the material of the apical callosity had been removed. It discloses the cast of the elongated visceral cavity and those of the vascular sinuses and cardinal muscle-scars. Fig. 10 shows an interior in which the visceral cavity and the two little pits are well seen. The foramen itself is lost in the broken posterior margin, but the narrowing of the visceral cavity to the notch or groove connecting it with the foramen is quite obvious. The casts of the little pits are rarely or never seen; they are not obvious, as in *A. comleyensis*.

Occasionally the casts of the vascular sinuses are proportionally wider apart than they are in the more typical form represented in fig. 9 *b*.

The dorsal valve is very little wider than long, rather strongly convex, and has the apex bent down almost to the margin. In the interior (figs. 15 & 17) there is a septal ridge which widens at its termination; the vascular sinuses are well developed posteriorly; between them and the septal ridge are two elongated prominences; the cardinal muscle-scars are large.

The shell-substance is corneous and very thick, both in the dorsal valve (see fig. 14 *b*) and in the apical callosity of the ventral. The exterior is marked by concentric striae, fairly closely set and apparently strongest on the dorsal valves; where exfoliation has occurred, radiating striae are clearly seen.

Dimensions.—The lengths vary from 1·5 to 3 mm.; the proportion of length to width is about 1 to 1·1; and that of length to height is 1 to ·4 in ventral, and 1 to ·25 or ·3 in dorsal valves.

Localities and horizon.—Comley, Lower Cambrian, from the *Lapworthella* Limestone of Comley Quarry, horizon Ad; also near Neves' Castle, in the Wrekin district.

COMPARISON BETWEEN *ACROTHYRA COMLEYENSIS* AND *A. cf. SERA*.

	<i>A. comleyensis</i> .	<i>A. cf. sera</i> .
(1) Posterior margin ...	decidedly notched.	almost straight.
(2) Inclination of pos- terior slope	forwards.	{ forwards, vertical or backwards.
(3) Concentric striae ...	weak and very close-set.	comparatively strong.
(4) Radiating striae ...	barely visible.	moderately strong.
(5) Visceral cavity ...	widely bell-shaped.	elongate.
(6) Casts of little pits	very obvious.	not obvious.
(7) Septal ridge of dorsal valve	{ strong posteriorly, weak centrally, reaching far forwards.	{ weak posteriorly, widening centrally.

Fig. 4.—Outline-sketches of the interiors and lateral aspects of five forms of ventral valves of the genus *Acrothyra*, omitting the muscle-scars and vascular sinuses.

[Figs. C, D, & E are copied from Matthew.]

- A. *A. comleyensis* Cobbold.
- B. *A. cf. serri* Matthew, Cobbold.
- C. *A. signata* sera Matthew.
- D. *A. signata* Matthew.
- E. *A. proavia* Matthew.

Many of Dr. Walcott's figures of *Acrothyra sera* (*op. cit.* pl. lxxx) may be matched from the Comley specimens; but the vertical groove on the false area in his figures 4 *a''* & 5'' is much stronger than any noted from Comley.

The description given by Dr. Matthew under *A. signata sera* (*op. cit.* p. 383) applies fairly accurately to the Comley shells. Of the interior he says that it has

‘Near the hinge a thick rectangular callus hollowed at the middle, with a depression that deepens towards the hinge; from this it is divided by a low transverse ridge, behind which is a pit leading to the foramen.’

He makes no mention of the connexion between the depression and the foraminual pit, such as is seen in *A. comleyensis* and in *A. cf. sera*. In his description of *A. signata* (*op. cit.* p. 381), however, he says:—

‘There is a strong narrow callus, one-third the length of the valve, bounded by a raised ridge at the sides and in front; at the front of this callus is an oval pit from which a groove runs backward nearly to the apex of the shell, where it is supposed to connect with the foramen.’

In *A. proavia*, the type of the genus, no mention is made of a connecting groove; but the hollow is represented in Matthew's figures (*op. cit.* text-fig. 2, p. 386, & pl. ii, figs. 4 *b-5 b*) as continuous nearly as far as the apex. Apparently in this species the hollow, the groove, and the foraminual pit are merged into one.

These differences appear to be connected with the elongation of the valve backwards, though which form is the more primitive cannot be determined upon the present evidence.

Text-fig. 4 (p. 352) is introduced so as to show graphically the modifications here mentioned.

BILLINGSSELLIDÆ Schuchert

BILLINGSSELLA Hall & Clarke.

BILLINGSSELLA COBBOLDI Matley.

Billingsella cobboldi Matley, 1911, Q. J. G. S. vol. lxxvii, p. 305 & pl. xxvi, figs. 20-23.

Locality and horizon.—Comley, Middle Cambrian; from the *Paradoxides-davidis* Flags, horizon Bb₂.

BILLINGSSELLA LINDSTRÖMI, var. *salopiensis* Matley. (Pl. XXIII, figs. 24 *a-24 c.*)

Billingsella lindströmi Linnarsson, var. *salopiensis* Matley, 1911, Q. J. G. S. vol. lxxvii, p. 303 & pl. xxvi, figs. 14 *a-17 c.*

Since this form was described by Dr. Matley, a large number of additional specimens have been obtained from the *Billingsella* Bed of Caradoc Dingle, many being more complete than those figured with his notes. From these it appears that the apex of the ventral valve is recurved over the posterior border to a greater extent than is shown in his figure, hiding the convex deltidium, which has not been observed.

An extreme case, probably an old individual, is now figured.

Locality and horizon.—Comley, Middle Cambrian; from the *Billingsella* Beds, horizon Be; also, sparingly, from horizon Bb₃.

BILLINGSSELLA sp. indet. (Pl. XXIII, figs. 23 a–23 c.)

A single ventral valve, represented by the external and internal casts [1879, 1880], was found in the matrix of the Comley Breccia-Bed. The specimens do not preserve any of the finer internal markings, and the external characters are but poorly indicated. These consist of some twenty-five angular (?) radiating ribs, which increase at varying distances by interpolation, and are crossed by concentric lines of growth. The ribs are reproduced on the internal surface.

Dimensions.—Length = 8 mm.; width = 9.5 mm.; height = 1.6 mm.

Locality and horizon.—Comley, Middle Cambrian; from the Comley Breccia-Bed, *Paradoxides-intermedius* Grits, horizon Bb₁.

ORUSIA Walcott.

ORUSIA LENTICULARIS Wahlenberg. (Pl. XXIII, figs. 25–27.)

Orthis (Orusia) cf. *lenticularis* Wahlenberg, Matley, 1911, Q. J. G. S. vol. lxvii, p. 306 & pl. xxvi, figs. 24–26.

Orusia lenticularis Wahlenberg, Walcott, 1912, U.S. Geol. Surv. Monogr. vol. li, p. 765 & pl. xeviii, figs. 1, 1 a–1 p, 2, 2 a–2 k, 3, 3 a & 3 b, 6, 6 a–6 c.

Since Dr. Matley wrote his notes additional specimens have been found at Comley, among which almost every variation figured by Dr. Walcott can be matched. There are also internal casts showing the fissures which represent the crural plates of the dorsal valve and the dental plates of the ventral mentioned,¹ but not figured, by Dr. Walcott; the former are comparatively near together and sometimes converge, the latter are wide apart and divergent.

Locality and horizon.—Comley, Upper Cambrian; from the *Orusia* Shales, horizon Ca.

VI. THE COMLEY PTEROPODA, ETC.

HYOLITHUS Eichwald.

HYOLITHUS CRASSUS, sp. nov. (Pl. XXIV, fig. 28.)

Type-specimen [2603].

Diagnosis.—Shell straight (so far as seen), apparently subquadrate in section, ratio of diameters about 1 to .7, lateral angles well rounded, rate of taper uncertain; shell-substance of two layers—the inner thick, smooth, and calcareous, the outer thinner and marked with numerous, irregular, and very coarse wrinkles; aperture transverse on the ventral face, projecting forward on the dorsal, and sinuous at the sides.

Dimensions.—Apertural diameter, 5 or more millimetres. The length may have been from 40 to 50 mm.

¹ 1912, p. 767.

The wrinkles of the outer surface recall those of *H. corrugatus* Kiær¹; they are, however, more irregular and less coarse than those shown in Dr. Kiær's figure. He regards his species as belonging to the sub-genus *Orthotheca*.

The resemblance in the shell-structure of this species to that of *H. sculptilis*, sp. nov. (p. 356) is very close, although the character of the sculpture differs radically.

Locality and horizon.—Comley, Lower Cambrian; from the *Protolenus* Limestone, Ac.

HYOLITHUS STRETTONENSIS, sp. nov. (Pl. XXIV, figs. 30 & 31.)

Hyolithus cf. *primævus* Groom, Cobbold, 1916, Rep. Brit. Assoc. (Manchester, 1915) p. 118.

Cf. *Hyolithus primævus* Groom, 1902, Q. J. G. S. vol. lviii, p. 116.

Type-specimen [1903].

Diagnosis.—Shell straight, tapering uniformly at rates which vary between 1 in 4·5 (apical angle 13°) and 1 in 7 (apical angle 8°); section a segment rather greater than a semicircle, ratio of diameters about 1 to ·66, dorsal side very slightly convex, ventral semicircular, lateral angles rounded; aperture in two planes, projection of dorsal lip about ·2 of the width; sculpture consisting of faint closely-set striae of growth.

Dimensions.—Width of aperture = 6 to 7 mm.; total length = 30 to 40 mm.

HYOLITHUS STRETTONENSIS, var. *BREVIS* nov. (Pl. XXIV, fig. 32.)

From the same general horizon a variety, or allied species, is found, which has a rate of taper varying from 1 in 2·5 to 1 in 3·5 (apical angles 23° to 18°), and a rather smaller proportional projection of the dorsal lip.

Dimensions.—Width=about 5·5 mm.; length=16 mm.

The variation in the apical angle is similar to that of *H. americanus* Billings,² which differs from *H. strettonensis* in having a greater proportional projection of the dorsal lip and a triangular section.

H. communis Billings³ also has a greater projection of dorsal lip than *H. strettonensis*; it is more elongate, and its section is flattened both dorsally and ventrally.

A curious confusion appears to have crept in as to the sub-generic reference of *H. communis*. In the original description Billings distinctly refers to the projecting [dorsal] lip; he, however, describes an operculum under the same reference, which seems to belong to an *Orthotheca*. Walcott (1886, p. 136) adopts Billings's description, and then observes that Ford's species *H. emmonsii* seems to be a variety. From the original description (Ford, 1873, p. 214) this species is evidently an *Orthotheca*, and is recognized

¹ 1916, p. 25 & pl. iii, figs. 10 a-10 b.

² 1872, p. 215, text-figs. 2 a & 2 b

³ 1872, p. 214.

as such by A. W. Grabau (1900, p. 655). G. Holm (1893, p. 30) places *Hyalolithus communis* under *Orthotheca*, and other writers seem to have followed him.

H. strettonensis resembles *H. primævus* Groom¹ from Malvern, and may indeed be identical with it. Dr. T. T. Groom is, however, uncertain whether his species is an *Orthotheca* or a *Hyalolithus* sensu stricto; and, until specimens elucidating this point are forthcoming, it seems advisable to describe the Comley form as a distinct species, lest a similar confusion to that mentioned above should arise.

Locality and horizon.—The Cwms, Church Stretton (a continuation of the Comley Cambrian in a south-westerly direction). Lower Cambrian. From the greenish-grey sandstone, horizon Ab₁ of the Road Quarry.

The variety *brevis* was found in beds of the same horizon at the north-eastern end of Caradoc Hill.

HYALITHUS (?) *SCULPTILIS*, sp. nov. (Pl. XXIV, figs. 25–27.)

Type-specimens [2583 & 2587].

Diagnosis.—Shell straight, or slightly tapering, sub-oval in section, with one side rather less convex than the other, ratio of diameters 1 to .7; shell-substance of two layers—the inner calcareous, thick, and smooth internally, the outer thin and covered with short, parallel, transverse wrinkles arranged in longitudinal rows that are separated by lines of plain shell; occasionally two or more rows will coalesce, but as a rule they remain distinct. The wrinkles may be straight or crescentic, and sometimes those of one row are set at a sharp angle with those of the contiguous row. In one fragment, probably from the neighbourhood of the aperture, the rows are traversed and replaced by continuous ridges or lines of growth.

Estimated dimensions.—Length = 30–40 mm.; apertural diameter = about 4 mm.

Small fragments with the characteristic sculpture are plentiful in the *Protolenus* Limestone; in only one instance is the sectional form seen, and in no case is there anything like a complete shell; on this account the generic reference is given with reserve. At first sight the fragments suggest a reference to *Conularia*; there are, however, no re-entering suture-like lines such as are depicted in Tate's figures of his species *H. conularoides*.²

Locality and horizon.—Comley, Lower Cambrian; from the *Protolenus* Limestone, Ac₅.

HYALITHUS sp. indet. Operculum (Pl. XXIV, fig. 29).

An operculum, which cannot with certainty be referred to any accompanying shell occurs in considerable numbers in the *Protolenus* Limestone.

¹ 1902, p. 110 & text-figs. 17–24.

² 1891, p. 186 & pl. ii, figs. 1–1a.

It is somewhat elliptical in general outline: the conical dorsal portion has a slight elevation, is marked by numerous fairly-strong concentric and also radiating striae; the ventral (upturned) portion is set nearly at right angles to the dorsal, and rises above the general plane of the plate to a height of about a third of the dorsi-ventral diameter. The angle at the apex of the conical portion is about 115° .

Dimensions.—Dorsi-ventral diameter = 7.5 mm.; transverse diameter = 9 mm.

Locality and horizon.—Comley, Lower Cambrian; from the *Protolenus* Limestone, horizon Ac_5 .

Sub-genus *Orthotheca*.

HYOLITHUS (ORTHO THECA) BAYONET Matthew. (Pl. XXIV, figs. 12–16.)

Orthotheca bayonet Matthew, 1899, Bull. Nat. Hist. Soc. New Brunswick, xviii, p. 193 & pl. iii, figs. 1 a–1 e.

Orthotheca bayonet Matthew, 1899, Trans. Roy. Soc. Canada, ser. 2, vol. v, sect. 4, p. 114 & pl. vii, figs. 1 a–1 e.

This species is of frequent occurrence at one horizon at Comley, and is represented a little lower in the series by a variety or mutation.

The more typical forms agree in all respects with the two descriptions given by Matthew, except that the rate of taper is about 1 in 6, and not 1 in 5 or 1 in 4 as given for the Newfoundland specimens, and that the ventral faces sometimes exhibit a few longitudinal striae.

The declination of the orifice from the dorsal side to the ventral angle is well shown (fig. 15), as is also the concave outline of the dorsal lip (fig. 13).

Locality and horizon.—Comley, Lower Cambrian; from the *Protolenus* Limestone, Ac_5 .

A variety which may be named *STRIATUS* (fig. 11) has more definite and regularly-arranged longitudinal striae, and the growth-lines indicate that the declination of the aperture towards the ventral angle was slighter.

Locality and horizon.—Comley, Lower Cambrian; from the *Strenuella* Limestone, Ac_4 .

HYOLITHUS (ORTHO THECA) COMPRESSUS, sp. nov. (Pl. XXIV, figs. 17 a & 17 b.)

Type and only specimen [225, 226].

Diagnosis.—Shell, a curved elongated pyramid, rate of taper changing from 1 in 4 near the apex to 1 in 3 towards the aperture (equivalent angles 14° & 19°). Dorsal and ventral sides indistinguishable. Section, an elongated ellipse, ratio of axes 1 to .33, lateral angles well rounded. Shell-substance thin, bluish grey, phosphatic (?). Surface: marked by transverse lines of growth,

spaced 4 to the millimetre. Aperture at right angles to the axis of the shell.

There appear to be no nearly-allied forms with which this species may be compared. The apical portion, if found detached, would be very similar to *Torerella*.

Locality and horizon.—Comley, Lower Cambrian; horizon Ac_2 (the *Olenellus* Limestone) of Comley Quarry.

HYOLITHUS (ORTHOTHECA) DE GEERI Holm (?).

Hyolithus (Orthotheca) de geeri Holm, 1893, Sver. Geol. Undersökn. ser. C, No. 112, p. 54 & pl. i, figs. 25-27.

Hyolithus Cobbold, 1912, Rep. Brit. Assoc. (Portsmouth, 1911) p. 114.

Hyolithus (Orthotheca) de geeri Holm, Cobbold, 1919, Geol. Mag. p. 150 & pl. iv, figs. 1-6.

Some of the casts mentioned in the British Association Report agree in cross-section and apical angle with Holm's species, but are not sufficiently well preserved for accurate determination. They are smaller than the type from Scandinavia.

Locality and horizon.—Comley, Lower Cambrian; from beds which are mapped as belonging to horizon Ab_2 .

HYOLITHUS (ORTHOTHECA) sp. indet. a. (Pl. XXIV, fig. 34.)

An oval operculum, which appears to belong to the sub-genus *Orthotheca*, cannot be assigned to any associated shell. It is almost flat, the nucleus rising a very little above the plane of the margin; the exterior is marked by numerous concentric striae and two lines on each side radiating from the nucleus (which is sub-central) towards the lateral much rounded angles.

Dimensions.—Dorsi-ventral diameter = about 3 mm.; transverse diameter = about 6 mm.

This operculum recalls that assigned by me with reserve to *H. (O.) de geeri* Holm, from the *Hyolithus* Limestone of Harts-hill, Nuneaton.¹ The principal difference is that the angle at the apex of the dorsal portion of the Comley specimen is about 130° instead of 90° .

HYOLITHUS (ORTHOTHECA) ? sp. indet. b. (Pl. XXIV, fig. 35.)

A very different operculum from the same horizon as the last is referred somewhat doubtfully to the sub-genus *Orthotheca*. It is sub-triangular, has the nucleus situated near the ventral angle and the conical dorsal projection is very narrow, the angle at the apex being only about 45° . On each side of its boundary-lines two projections on the interior appear to indicate muscle attachments. The exterior has faint striae of growth near the margin. The margin appears to be in one plane, and the plate when in place would be (on the whole) concave.

Locality and horizon for both opercula.—Comley, Lower Cambrian; from the *Protolenus* Limestone, horizon Ac_3 .

¹ Cobbold, 1919, p. 150 & pl. iv, figs. 7-9.

LAPWORTHHELLA, gen. nov.

Diagnosis.—Small, tapering, annulated shells of circular or sub-polygonal section, consisting of two layers, the inner one of medium thickness, the outer very thin and chitinous. Annulations formed by pronounced raised ridges encircle the tube, and divide it into sub-equal parts.

Type-species—*Lapworthella nigra*, sp. nov.

Localities and horizon.—Lower Cambrian, near the summit. Comley and Neve's Castle, both in Shropshire.

The genus appears to be intermediate between *Hyolithus* (*Orthotheca*) and *Salterella*, with certain resemblances to *Tentaculites*.

From *Orthotheca* it is distinguished by the character of the encircling ridges, which are quite different from striae of growth.

The resemblance to *Tentaculites* consists in the form and disposition of the same encircling ridges.

The relations to *Salterella* are much closer. In size, curvature of shell, and frequent circularity of aperture the two genera have much in common. The differences consist in the structure of the shell and the form of the annulating ridges.

The generic characters of *Salterella* are thus given by Billings:

'small, slender, elongate-conical tubes, consisting of several hollow cones placed one within another, the last forming the chamber of habitation of the animal. The cross-section of the tube is circular or sub-triangular, and they are either straight or gently curved, the surface is concentrically or longitudinally striated.' (1861. vol. i, p. 17.)

In his description of the type-species *Salterella rugosa*, Billings states that the annulations are formed by the projecting edges of the successive cones, and Dr. Walcott's figures of the species¹ show this structure very clearly.

Other species referred to *Salterella* are (1) *S. pulchella* Billings,² (2) *S. curvata* Shaler & Føerste,³ and (3) *Serpulites maccullochi* Salter.⁴

In the first the structure of the tube has a tripartite arrangement; in *S. curvata* it is stated to be thick, but no indication is given of the shell-structure; in *Serpulites maccullochi* the tube appears to be very thick, but its structure is not described. Billings suggests (*op. jam cit.*), however, that *Salterella rugosa* 'must be closely allied to *S. maccullochi*,' and his generic reference for that species has been generally adopted.

In the absence of any detailed study of the shell-structure of the two last-named species, it may be left an open question whether they should continue to be referred to the genus *Salterella*.

¹ 1890, pl. lxxix, fig. 4.

² 1861, p. 18. Walcott's figures of *S. pulchella* (1890, pl. lxxix, figs. 5, 5 a-5 e) suggest that more than one species may be represented.

³ Shaler & Føerste, 1888, p. 34.

⁴ 1859, p. 381.

Prof. Charles Lapworth was the first to point out that the *Olenellus* and *Paradoxides* Zones were in contact in the Shropshire area, and he has also shown very great interest in the explorations conducted at Comley for the British Association. It seems, therefore, appropriate to dedicate a genus that marks, at least locally, the dividing-line between the Lower and Middle Cambrian to his honour.

LAPWORTHHELLA NIGRA, sp. nov. (Pl. XXIV, figs. 1-6.)

Salterella (perhaps 2 spp.), Cobbold, 1909, Rep. Brit. Assoc. (Dublin, 1908) p. 235.
Salterella Cobbold, 1910, Q. J. G. S. vol. lxvi, p. 20.

Type-specimens [270, 267]; also [RR. 1221], Geological Survey collection.

Diagnosis.—Shell: a small, pyramidal or conical, curved tube, divided into from twelve to twenty annulations; rate of taper, variable, from 1 in 1·4 to 1 in 4·3 (equivalent apical angles 41° to 13°); aperture transverse, varying from sub-circular to sub-rhomboidal; apex sharp; shell-substance of two layers: (1) inner layer moderately thick, calcareous and annulated on the exterior by prominent ridges, which are at right angles to the curved axis of the tube, pass all round it, and are set at nearly uniform distances apart; (2) outer layer very thin, of black, chitinous (?) material, which passes over the transverse ridges giving them a sharp edge; interior of shell smooth, or slightly marked by annular lines of thickening corresponding to the transverse ridges. No septa or operculum have been observed.

Dimensions.—Lengths = 2 to 3 mm.; diameter of aperture = 2 to 1·5 mm.

The species is very variable: not only is there a considerable range in the apical angle, but the sectional form is not constant. Some specimens appear to be truly circular, others are more or less semicircular, and others again are sub-trapezoidal. There is also much variation in the spacing of the encircling ridges.

Localities and horizon.—Comley and Neve's Castle; from the *Lapworthella* Limestone, horizon Ad.

SALTERELLA Billings.

SALTERELLA ($\frac{?}{2}$) BELLA, sp. nov. (Pl. XXIV, fig. 33.)

Type-specimen [523].

Diagnosis.—Shell a small, tapering, curved tube with elliptical aperture, axes in the proportion of 1 to ·6, rate of taper 1 in 1·5, equivalent apical angle 40° ; shell-substance calcareous, thick, and strong; exterior marked with numerous, delicate, impressed striae of growth and from five to eight irregular constrictions; no radial striae; apex not preserved.

Dimensions.—Height = 1·25 mm.; longer diameter of aperture = 1 mm.; length = 1·5 mm.

SALTERELLA (?) *STRIATA*, sp. nov. (Pl. XXIV, fig. 45.)

Type-specimen [139].

Diagnosis.—Shell a small, tapering, curved tube with elliptical aperture; axes in the proportion of 1 to '6, rate of taper 1 in 1·8, equivalent apical angle 32°; shell-substance calcareous, thick, and strong; exterior marked with eight or more constrictions (which are reproduced on the interior) and numerous fine radiating striæ; no lines of growth seen.

Dimensions.—Height=2 mm.; diameter=1·7 mm.; length=3 mm.

Observations.—Owing to the absence of any obvious cone-within-cone structure such as characterizes the type-species *S. rugosa*, the reference of these two species to Billings's genus is made with reserve. The thick calcareous shell and bent conical form indicate a relationship with *S. curvata* and *Serpulites maccullochi* (see p. 359). On the other hand, *Salterella striata* has points of resemblance with *Helcionella* and *Scenella*.

Locality and horizon.—Comley, Lower Cambrian. Both species are found in the *Protolenus* Limestone, horizon Ac₅.

HYOLITHELLUS Billings.

HYOLITHELLUS MICANS Billings. (Pl. XXIV, figs. 19–21.)

- Hyolithellus micans* Billings, 1872, 'Canadian Naturalist' n. ser. vol. vi, p. 215.
Hyolithellus micans Billings, Walcott, 1886, U.S. Geol. Surv. Bull. 30, p. 142 & pl. xiv, figs. 2, 2a–2c.
Hyolithellus micans Billings, Shaler & Færste, 1888, Bull. Mus. Comp. Zool. Harvard, vol. xvi, p. 34 & pl. ii, fig. 23.
Hyolithellus micans Billings, Walcott, 1890, 10th Ann. Rep. U.S. Geol. Surv. p. 624 & pl. lxxix, figs. 1, 1a–1c.
Hyolithellus cf. *micans* Billings, Lapworth, 1891, Geol. Mag. p. 532.
Hyolithellus micans Billings, Matthew, 1899, Bull. Nat. Hist. Soc. New Brunswick, xviii, p. 192 & pl. ii, figs. 1a–1d.
Hyolithellus micans Billings, Matthew, 1899, Trans. Roy. Soc. Canada, vol. v, sect. 4, p. 109 & pl. vi, figs. 1a–1d.
Hyolithellus micans Billings, Grabau, Occ. Pap. Boston Nat. Hist. Soc. vol. iv, pt. 3, p. 658 & pl. xxxii, fig. 12.

Short lengths of tubes with black shining surfaces are of frequent occurrence at Comley, and, although they assume many aspects, there are no well-defined characters to separate them one from the other or from Billings's species. They occur at many horizons, ranging from Ab₁ in the Lower to Bb₁ in the Middle Cambrian. In section they are circular, the surface may be smooth, or annulated with lines or bands of growth; they are most frequently straight, but sometimes sinuous (fig. 20) or even strongly curved (fig. 19) and they show very slight traces of taper. The diameters vary from '1 to 2·5 or even 3 millimetres.

In thin sections the tube is seen to consist of one thin layer of chitinous material, which must have been comparatively strong and resistant, for the specimens rarely show any sign of collapse or crushing.

Locality and horizon.—Comley, Lower and Middle Cambrian; from horizons Ac₁, Ac₂, Ac₃, Ac₄, Ac₅, Ad, Ba₁, Ba₂, Ba₃,

Bb₁; that is, from below the *Olenellus* Limestone to the representative of the *Paradoxides-tessini* Zone of Scandinavia.

HYOLITHELLUS MICANS, var. *PALLIDUS* nov. (Pl. XXIV, fig. 23.)

A form, which may be separated as a distinct variety, occurs at a comparatively low horizon in the Lower Cambrian. It has a truly circular section, a rather stronger rate of taper, and, so far as known, is always straight. The surface is grey and dull, not glistening; it shows slight traces of striae of growth. The diameter varies from .3 to 1.2 millimetres.

Locality and horizon.—Comley, Lower Cambrian; from the *Holmia* horizon Ab₃; also doubtfully from horizon Ab₂.

HYOLITHELLUS MICANS, var. *ROBUSTUS* nov. (Pl. XXIV, fig. 22.)

This variety is one of those forms that is easily distinguished when seen and handled, but the special characteristics are very difficult to describe in words.

It is separated from the more typical forms of *H. micans* by its usually larger diameter (3 to 5 millimetres), its more decided rate of taper (about 1 in 30), and in that it only occurs associated with a *Paradoxides* fauna.

The outer skin is thin (?), phosphatic, and usually grey; in addition, it frequently has a thick lining of calcite which, in one case, where the tube is 3 mm. in diameter, is as much as .4 mm. thick, and simulates an inner shell. Whether this lining represents part of the original organism, or is a mineral deposit subsequent to its death cannot at present be determined.

Locality and horizon.—Comley, Middle Cambrian; from horizons Ba₁ of the *Paradoxides-ølundicus* Zone and Bb₁ at the base of the *Paradoxides-tessini* Zone.

HYOLITHELLUS (?) *SINUOSUS*, sp. nov. (Pl. XXIV, fig. 24.)

Type-specimen [1912].

This specific name is proposed for small, circular, phosphatic tubes which are curved in more than one plane, have a dull grey surface, and are not associated with any that can be referred to *H. micans*. Only two or three fragmentary specimens have been as yet found, and one of these shows distinct lines of growth, evenly spaced along the tube.

Locality and horizon.—The Cwms, near Comley, Lower Cambrian; from the greenish-grey sandstone, *Obolella-groomi* Beds, horizon Ab₁.

SALOPIELLA, gen. nov.

Diagnosis.—Small tubes with elliptical section and sub-parallel margins; shell-structure calcareous (?), thin, and formed of a number of closely-set, step-like annulations, which encircle the tube and are reproduced on the interior.

Genotype.—*S. obliqua*, sp. nov.

Locality and horizon.—Comley (Shropshire), Lower Cambrian.

These tubes agree with Holm's genus *Torella* in their elliptical section and sub-parallel margins; but there is no trace of the phosphatic shell of that genus, and the wall of the tube is very different.

The external aspect recalls that of *Salterella*, but the elongate form of the tubes and the absence of any sign of the cone-within-cone structure of the shell preclude any reference to that genus.

SALOPIELLA OBLIQUA, sp. nov. (Pl. XXIV, figs. 18 a–18 c.)

Type-specimen [242].

Diagnosis.—Shell tubular, thin, slightly curved, and having a very slight taper; section elliptical, ratio of diameters 1 to .55. Shell-substance calcareous (?); shell-structure consisting of a number of angular step-like corrugations which are spaced about .12 to the millimetre, pass obliquely round the tube, and are reproduced on the interior. One slope of these corrugations is nearly parallel to the axis of the tube, the other is very steeply inclined to it.

Dimensions.—Larger diameter=about .6 mm.; greatest length seen = 3.5 mm.

Observations.—The type-material consists of an internal cast in calcite [212] and a partly internal, partly external cast in sandy limestone [231]. The shell appears to have been calcareous, for in the second specimen it has entirely disappeared, with no trace of any phosphatic material.

Locality and horizon.—Comley, Lower Cambrian; from the *Strenella* Limestone, horizon Ae₄.

HELENIA Walcott, 1890.

HELENIA CANCELLATA, sp. nov. (Pl. XXIV, figs. 7 a–10.)

Helenia Cobbold, 1911, Q. J. G. S. vol. lxxvii, p. 297.

Type-specimens [245, 246].

Diagnosis.—Shell: a flattened curved tube, tapering very gradually to a blunt, closed end; section, a narrow oval; shell-substance thin, phosphatic; exterior beautifully sculptured by two sets of impressed lines running obliquely across the shell and leaving minute, diamond-shaped projections between them.

Dimensions.—Diameter=about .5 mm.; greatest length seen = 6 mm.

Observations.—Usually there is a longitudinal depression close to the concave margin—it does not reach the closed end, is not seen at all in one specimen, and is probably due to the collapse of the tube.

The type-species of the genus, *H. bella* Walcott,¹ is described as 'probably open at both ends'; it has but one set of diagonal lines.

H. granulata Matthew² is described as having a minutely

¹ 1889, p. 39; and 1890, p. 610 & pl. lxxviii, figs. 4, 4 a–4 b.

² 1899 a, p. 192; 1899 b, p. 107 & pl. vi, figs. 7 a–7 e.

granular surface; it is larger, and tapers more rapidly than *H. cancellata*.

Locality and horizons.—Comley, Lower Cambrian; from the *Olenellus* Limestone, horizon Ac₂, the *Microdiscus-bellimarginatus* Limestone, horizon Ac₃, the *Protolenus* Limestone, horizon Ac₄, and the *Lapworthella* Limestone, horizon Ad.

VII. THE COMLEY GASTEROPODA.

SCENELLA Billings:

SCENELLA ELEVATA, sp. nov. (Pl. XXIV, fig. 36.)

Type and only specimen [RR 1211, Geological Survey, Rhodes collection, 1892].

Diagnosis.—Shell patelliform with apex towards the anterior (?) end, depressed slightly below the highest point; aperture not seen, probably an ellipse; posterior and lateral slopes very slightly convex and inclined, anterior slope approaching the vertical; shell-substance calcareous, exterior with closely-set, impressed radiating lines, scarcely visible on the upper half of the shell, but strong towards the margins with the interspaces rounded; these are crossed by delicate concentric lines of growth and six or seven stronger lines, which near the apex tend to form constrictions, but in the remainder of the shell mark angular changes of curvature.

Dimensions.—Length=about 3.5 mm.; width=3 mm.; height=1.66 mm.

The species is referred to Billings's genus, on account of the absence of pronounced concentric rugosities, though in the depression of the apex it resembles *Helcionella*. It may be contrasted with Dr. Kiær's two species, *Scenella antiqua* and *Sc. depressa*,¹ from Tönten.

Locality and horizon.—Comley, Lower Cambrian; from the *Olenellus* Limestone, horizon Ac₂.

HELCIONELLA Grabau & Shimer.

HELCIONELLA CINGULATA, sp. nov. (Pl. XXIV, fig. 37.)

Type-specimen [827].

Diagnosis.—Shell patelliform, with elevated apex; aperture not known, but, as indicated by the cross-section, an elongated ellipse with axes in the ratio of 5 to 2; apex slightly curved over towards the anterior (?) end; shell-substance calcareous, with black, phosphatic, outer pellicle; exterior marked by concentric rounded fillets, set at even distances apart and separated by flat or slightly concave spaces about three times as wide as the fillets themselves.

Dimensions.—The only known specimen is fragmentary; it is 2 mm. in height and 4 in length.

¹ 1916, pp. 17, 18 & pl. ii, figs. 1, 1 a-1 c, 2 a-2 c.

Observations.—Similar concentric fillets are shown in Shaler & Færste's figure¹ of their species *Stenotheca curvirostra*, which is different in general form and has longitudinal striae.

Locality and horizon.—Comley, Lower Cambrian; from the *Protolenus* Limestone, horizon Ac₃.

HELICONELLA OBLONGA, sp. nov. (Pl. XXIV, figs. 38 & 39.)

Stenotheca Cobbold, 1911, Q. J. G. S. vol. lxvii, p. 297.

Type-specimen [1316].

Diagnosis.—Shell patelliform, with apex strongly recurved; aperture oblong, with straight parallel sides and rounded ends; apex moderately sharp, bent down over the anterior (?) end to a height of about .75 of the maximum. and at a point a little short of the anterior border; shell-substance thick and calcareous; exterior marked by very regular, step-like, concentric corrugations, retaining throughout the form of the aperture, also by numerous radiating thread-like raised lines spaced about five to the millimetre and increasing by interpolation; interior smooth, except for the reproduction of the external corrugations.

Dimensions of type-specimen.—Length=14 mm.; width=9.5 mm.; greatest height=4 mm.; height of apex=about 3 mm.

Observations.—This species occurs fairly plentifully in the matrix of the Middle Cambrian conglomerate of Comley Quarry, and is readily identified, even when poorly exposed, by the regularity of sculpture and the form of aperture. These features also serve to separate it from *H. rugosa* (Hall) and from other Comley forms of the genus.

The regularity of the step-like corrugations recall those shown in J. W. Salter's figure² of *Stenotheca cornucopia* from the Middle Cambrian of South Wales. That species is preserved in shale, and appears to have had a thin shell.

Locality and horizon.—Comley, Middle Cambrian, horizon Ba₃; from the Conglomerate of Comley Quarry, and from the same bed at Robin's Tump.

HELICONELLA RUGOSA Hall, var. *COMLEYENSIS* nov. (Pl. XXIV, fig. 40.)

Metoptoma rugosa Hall, 1847, Pal. N. York, vol. i, p. 306 & pl. lxxxiii, figs. 6a-6c.

Stenotheca rugosa Hall, Billings, 1872, Canad. Nat. n. ser. vol. iii, p. 479.

Stenotheca rugosa Hall, Walcott, 1886, Bull. U.S. Geol. Surv. 30, p. 128 & pl. xii, figs. 1, 1 a-1 e.

Stenotheca rugosa, Hall, Walcott, 1890, 10th Ann. Rep. U.S. Geol. Surv. p. 617 & pl. lxxiv, figs. 1, 1 a-1 e.

Type for variety [1897].

This variety differs from the more typical forms of the species in the character of the encircling rugosities. The aperture is a wide ellipse, ratio of axes 1 to .9; the concentric rugosities of

¹ Bull. Mus. Comp. Zool. Harvard, vol. xvi, No. 3, p. 30 & pl. i, fig. 8.

² 1872, Q. J. G. S. vol. xxviii, p. 180 & pl. vii, figs. 12-13.

growth are very irregular, varying in strength from one part of their course to another and running one into another, so that there are no well-defined step-like annulations; the radiating striae are, on the other hand, quite regular: they increase by interpolation, and are spaced about seven to the millimetre.

Dimensions of largest specimens.—Length=17 mm.; width=15·5 mm.; greatest height=7 mm.; height of apex=4 mm.

Locality and horizon.—Comley, Lower Cambrian; from the *Olenellus* Limestone, horizon Ac₂, and from the *Callaria* Sandstone, horizon Ac₁.

LATOUCHELLA, gen. nov.

Diagnosis.—Shell small, calcareous, coiled or incipiently coiled so as to be isometrical about the median plane; aperture elongate, entire, not notched; whorls free, not in close or overlapping contact.

Type-species—*L. costata*, sp. nov.

Horizon.—Lower Cambrian.

The incipient coiling and elongate aperture are reminiscent of *Platyceras*, from which these shells differ in having the apex on the median plane.

In side view they approach some forms of *Helcionella*, particularly *H. paupera* Billings, but they are without encircling annulations or rugosities.

The absence of a notch in the apertural margin seems to preclude a reference to any of the Heteropod-like shells, such as *Pelagiella* Matthew, *Watsonella* Grabau, etc.

The external resemblance of *L. costata* to an immature ammonite is remarkable; there are, however, no indications of septa.

LATOUCHELLA COSTATA, sp. nov. (Pl. XXIV, figs. 41 & 42.)

Type-specimen in the collection of the Rev. W. M. D. La Touche, of Wistanstow, Craven Arms.

Diagnosis.—Shell consisting of about one and a quarter whorls; aperture oval (ratio of diameters about 1 to ·5) and slightly campanulate; sides provided with six to eight or more elongate prominent ribs, swollen towards the convex margin of the shell, which they do not cross, and narrowing to disappear in the opposite direction, these are reproduced in the interior: whorls standing clear of one another in the internal cast, probably just touching one another in the complete shell. Dorsum narrowly rounded, but not keeled.

Dimensions of the larger specimen.—Apertural diameters = 4×2 mm.; total length = 4·5 mm.; height = 2·5 mm.

Locality and horizon.—Comley, Lower Cambrian; from the *Olenellus* Limestone, horizon Ac₃, and from the *Protolenus* Limestone, horizon Ac₅.

LATOUCHELLA (?) STRIATA, sp. nov. (Pl. XXIV, figs. 43 & 44.)

Type-specimen [2655], other specimens [2659, 2661, 2662].

Diagnosis.—Shell coiled so as to make about three-quarters of a complete whorl; aperture an elongate egg-shaped oval, narrowing towards the dorsum; surface showing numerous, fine, impressed radiating striae which strengthen as they approach the aperture, and of which some four or six are still more strongly marked along the dorsum. Dorsum narrowly rounded, but not keeled; no concentric lines of growth visible in the specimens.

Dimensions of largest specimen.—Apertural diameters=about 6×2 mm.; total length=about 6 mm.; height=5 mm.

This species is proportionately higher and narrower than *L. costata*, and the whorls are distinctly separated one from the other.

Localities and horizon.—Comley, Lower Cambrian, from the *Protolenus* Limestone, horizon Ac_3 ; also one specimen from the *Hyolithus* Limestone of Woodlands Quarry, Hartshill, Nuneaton [Birmingham University Collection].

VIII. THE COMLEY OSTRACODA.

LEPERDITIA Rouault.

LEPERDITIA (?) LENTIFORMIS, sp. nov. (Pl. XXIV, figs. 46 *a*–46 *c*.)

Type-specimen [Birmingham University Collection]. Three other specimens known.

Diagnosis.—General outline bean-shaped; anterior end shortly truncate; dorsal border gently convex for about two-thirds of the total length, then bending more rapidly and passing to the posterior rounded end in a line that is almost straight; ventral border one long fairly-regular curve from the truncate anterior to the posterior end; surfaces of both valves regularly convex, except near the anterior end, where there is a short ill-defined depression passing obliquely towards the ventral border; test thin, corneous, having the surface marked by a network of impressed lines, the interspaces being convex, and more or less elongate and parallel with the border.

Dimensions.—Length=11.5 mm.; depth=7 mm.; thickness of the two valves together=4 mm.

The general aspect of the species recalls that of *L. (?) dermatoides* Walcott,¹ which chiefly differs in having a very thin membranous and punctate test; that species has also a minute scar towards the posterior end, and a rather differently curved border.

In the type-specimen the two valves are in their natural position.

Locality and horizon.—Comley, Lower Cambrian; from the *Olenellus* Limestone, horizon Ac_2 .

¹ 1890, p. 626 & pl. lxxx, figs. 1–1 *a*.

LEPERDITIA (?) *DERMATOIDES* Walcott (?).

Leperditia (?) *dermatoides* Walcott, 1890, 10th Rep. U.S. Geol. Surv. p. 626 & pl. lxx, figs. 1-1 a.

A single imperfect specimen [RR 1285] in the Geological Survey collection differs from *L.* (?) *lentiformis* in having a punctate test; it is also somewhat more convex and elongate. In these respects it simulates Walcott's species.

Locality and horizon.—Comley, Lower Cambrian; from the *Strenuella* Limestone, horizon Ac₁.

ESCASONA Matthew, 1902.

ESCASONA (?) spp. indet. (Pl. XXIV, figs. 47 & 48.)

Two forms suggest by their shape and convexity a reference to this genus, notwithstanding the fact that there is no sign of the ocular tubercle.

The two specimens figured [RR 1332 & RR 1335] are in the Geological Survey collection, and are too imperfect for closer identification. They probably represent two species.

Locality and horizon.—Comley, Lower Cambrian; from the *Olenellus* Limestone, horizon Ac₁.

BRADORIA Matthew.

BRADORIA (?) sp. indet. (Pl. XXI, figs. 28 a-28 c.)

A single specimen [14536] in the British Museum collection appears to belong to this genus. It is an internal cast, with some of the test adhering. In shape it is not unlike the figures of *Fordilla trogensis* Barrande, given by Dr. C. D. Walcott.¹ It has, however, a distinct marginal flat border, ventrally and at the two ends. There is no trace of an ocular tubercle near the hinge-line, but there is a tubercle low down towards the ventral anterior border. The body of the valve is marked with faint concentric lines, both inside and outside.

Dimensions.—Length=2.5 mm.; width=1.75 mm.; depth=.7 mm.

Locality and horizon.—Comley, Lower Cambrian; from the *Olenellus* Limestone, horizon Ac₁.

IX. THE FAUNAS OF THE COMLEY HORIZONS AND THEIR
ORDER OF APPEARANCE.

[Table II (p. 376) contains a list of all the fossils described or noted in this paper, together with the trilobites that have been recorded in previous communications; and also the horizons at which they have been found.]

The brachiopoda confirm and amplify the principal faunal divisions sketched out by me in 1911 (*b*, p. 296), and in my reports to the British Association. In some instances they lend considerable assistance in the correlation of the beds with those of other

¹ 1886, p. 125; and 1890, p. 615 & pl. lxxiii, figs. 2, 2 a-2 c.

areas; the pteropoda and gasteropoda, on the other hand, are, unfortunately, almost all new to science, but they serve to accentuate the divergences between the local faunas. The ostracoda, though well preserved, are very difficult of extraction without damage. Their general aspect is reminiscent of those described by Dr. G. F. Matthew (1902 & 1903) from the *Protolenus* and Etcheminian faunas and, so far as they go, these fossils help to confirm the opinion that the place of those two faunas is rather high up in the Lower Cambrian Series.

LOWER CAMBRIAN.

Horizon Aa. The Wrekin Quartzite.

Only one fragment of a corneous brachiopod has been found.

Horizon Ab₁. *Obolella* (?) - *groomi* Beds.

Micromitra (*Paterina*) *phillipsi*.
Micromitra (*Paterina*) *rhodesi*.
Micromitra (*Paterina*) *kingi*.
Walcottina lapworthi.
Walcottina elevata.

Obolella (?) *groomi*.
Hyolithus stretonensis.
Hyolithellus sinuosus.
Certain ovoid and elongate bodies
that have been calcareous, and show
only vague traces of organic origin.¹

Micromitra (*P.*) *phillipsi* and *Obolella* (?) *groomi* are found in the Malvern Quartzite and Hollybush Sandstone, while *Hyolithus stretonensis* may possibly be identical with *H. primævus* Groom of the same group.

On physical evidence Dr. C. Callaway correlated the sandstones of Comley with the Malvern Hollybush Sandstone; the fauna as enumerated above tends to prove that his correlation was, at least in part, correct.

Horizon Ab₂.

Hyolithus de geeri (?), *Hyolithellus micans* var. *pallidus*, and tracks on the surfaces of the flaggy beds.

Horizon Ab₃. *Holmia* Sandstone.

Holmia (?) sp. indet., *Hyolithellus micans* var. *pallidus*.

Horizon Ab₄.

Small and obscure ostracoda, gen. et sp. indet.

Horizon Ac₁. *Callavia* Sandstone.

Micromitra (*Paterina*) *labradorica*.
Obolella atlantica var. *comleyensis*.
Hyolithus sp. indet.
Hyolithellus micans.
Helcionella rugosa var. *comleyensis*.

Microdiscus atleboroensis.
Callavia callavi.
Wanneria (?) sp. nov.
Callavia cobboldi Raw MS. (?).

The relation of this fauna with that of other horizons is discussed below (p. 370).

Horizon Ac₂. *Olenellus* Limestone.

The *Olenellus* Limestone was the first horizon in the Comley area to yield any recognizable Lower Cambrian fossils, and naturally becomes a stratigraphical and palæontological datum to which other horizons may be referred. The complete fauna, so far as at present known, is tabulated below (p. 370).

¹ Cobbold, 1914, p. 486; and 1916, p. 118.

THE FAUNA OF THE *OLENELLUS* LIMESTONE OF COMLEY.¹

<i>Micromitra</i> (<i>Paterina</i>) <i>labradorica</i> .	<i>Bradoria</i> (?) sp. indet.
<i>Micromitra</i> (<i>Paterina</i>) <i>minor</i> .	<i>Microdiscus</i> <i>annio</i> .
<i>Micromitra</i> (<i>Paterina</i>) <i>minor</i> var. <i>gibba</i> .	<i>Microdiscus</i> <i>attleborensis</i> .
<i>Obolus</i> <i>parrulus</i> .	<i>Micmacca</i> (?) <i>ellipsocephaloides</i> .
<i>Obolella</i> <i>atlantica</i> var. <i>comleyensis</i> .	<i>Micmacca</i> (?) <i>ellipsocephaloides</i> var. <i>senior</i> .
<i>Hyalolithus</i> (<i>Orthotheca</i>) <i>compressus</i> .	<i>Micmacca</i> (?) <i>ellipsocephaloides</i> var. <i>spinosa</i> .
<i>Hyalithellus</i> <i>micans</i> .	<i>Micmacca</i> (?) <i>ellipsocephaloides</i> var. <i>strenuelloides</i> .
<i>Helenia</i> <i>cancellata</i> .	<i>Micmacca</i> (?) <i>parrula</i> .
<i>Scenella</i> <i>elevata</i> .	<i>Callaria</i> <i>callarei</i> .
<i>Helcionella</i> <i>rugosa</i> var. <i>comleyensis</i> .	<i>Callaria</i> cf. <i>callarei</i> .
<i>Latouchella</i> <i>costata</i> .	<i>Callaria</i> <i>cartlandi</i> .
<i>Lepeditia</i> (?) <i>dermatoides</i> (?)	
<i>Lepeditia</i> (?) <i>lenticularis</i> .	
<i>Escasona</i> (?) spp. indet.	

There are obvious connexions between this fauna and that immediately below. *Micromitra labradorica*, *Obolella atlantica* var. *comleyensis*, *Helcionella rugosa* var. *comleyensis*, *Microdiscus attleborensis*, and *Callaria callarei* have already appeared in Ac_1 , from which, however, the five forms of *Micmacca*, as well as the less frequent other fossils, are all absent, and one or more species of the Mesonacidae, not appearing in the *Olenellus* Limestone, are present.

On comparing the Ac_2 fauna with those above it, there is a similar connexion observable, and to a considerable extent through the same species, but *Micmacca* is absent in the superior horizons, as also is *Microdiscus attleborensis*, their places being taken in horizon Ac_3 by *Strenuella pustulata* and *Microdiscus bellimarginatus*.

The faunas of the two next higher horizons, Ac_4 and Ac_5 , are connected with the datum-fauna Ac_2 by *Micromitra labradorica*, *Obolus parrulus*, *Helenia cancellata*, *Latouchella costata*, and also by numerous fragments referred with reserve to *Callaria callarei*. Notwithstanding this connexion, the faunas of the *Strenuella* and the *Protolenus* Limestones (Ac_4 & Ac_5 , see p. 371) are very distinct from those below and also one from the other. In Ac_4 occur three species of *Strenuella*, *Microdiscus lobatus*, *M. speciosus*, *Lingulella viridis*, *Obolella atlantica* var. *transversa*, *Hyalolithus* (*Orthotheca*) *bayonet* var. *striata*, and *Salopiella obliqua*, none of which have been seen before; and in Ac_5 occur two species of *Protolenus*, two of *Mohicana*, *Microdiscus comleyensis*, *Hyalolithus crassus*, *H. sculptilis*, *H. (O.) bayonet*, *Salterella* (?) *bella*, *S. (?) striata*, *Latouchella striata*, and *Helcionella cingulata*, all of which are confined to the horizon. *Strenuella* has entirely disappeared, as also has *Obolella atlantica*.

NOTE.—In the foregoing paragraphs all mention of *Hyalithellus micans* has been omitted, in order to avoid repetition. Forms referred to this species

¹ In Cobbold, 1911 b, p. 298, this fauna was tabulated under the title 'Helena Group.' A specimen has now been found of the pygidium then assigned to *Microdiscus helena* Walcott, attached, with three thoracic segments, to a cephalon then referred to *Ptychoparia* (?) *attleborensis* Shaler & Færste. The name *Helena* is, therefore, now abandoned.

occur in all the Lower Cambrian horizons Ac_1 to Ad , and also in the Middle Cambrian horizons Ba_1 , Ba_2 , Ba_3 , and Bb_1 : that is, throughout the whole of the *Protolenus-Callavia* Fauna and of the *Paradoxides-elandicus* Zone, and in the *Paradoxides-tessini* Zone.

The fragments assigned to *Callavia*, together with *Helenia cancellata* and the connexions between the individual faunas of horizons Ac_1 to Ac_3 , serve to link them into one group, which has been called the *Protolenus-Callavia* Fauna¹ of Comley.

Of the species recorded, the following are found or represented in the Lower Cambrian of the Atlantic Province of North America, a considerable proportion being included in Dr. Matthew's *Protolenus* and *Etcheminian* Faunas.

Species of the *Protolenus-Callavia* Fauna of Comley (Horizons Ac_1 to Ac_3) which are represented in the Atlantic Province of North America:—

<i>Micromitra (Paterina) labradorica.</i>	<i>Microdiscus attleboroensis.</i>
<i>Obolella atlantica</i> var. <i>comleyensis.</i>	<i>Microdiscus bellinarginatus.</i>
<i>Obolella atlantica</i> var. <i>transversa.</i>	<i>Microdiscus lobatus.</i>
<i>Hyalolithus (Orthotheca) bayonet.</i>	<i>Microdiscus speciosus.</i>
<i>Hyalithellus micans.</i>	<i>Strenuella platycephala.</i>
<i>Helenia cancellata.</i>	<i>Strenuella salopiensis.</i>
<i>Helcionella rugosa</i> var. <i>comleyensis.</i>	<i>Protolenus latouchei.</i>
<i>Leperditia (?) dermatoides.</i>	<i>Micmacca (?) ellipsocephaloides</i> var.
<i>Escasona (?)</i> sp.	<i>spinosa.</i>
<i>Bradoria (?)</i> sp.	<i>Callavia callavei.</i>
<i>Microdiscus annio.</i>	<i>Wanneria (?)</i> sp.

Horizon Ac_1 . The *Strenuella* Limestone of Comley.²

<i>Obolella atlantica</i> var. <i>transversa.</i>	<i>Microdiscus lobatus.</i>
<i>Lingulella viridis.</i>	<i>Microdiscus speciosus.</i>
<i>Hyalolithus (Orthotheca) bayonet</i> var.	<i>Strenuella parva.</i>
<i>striatus.</i>	<i>Strenuella platycephala.</i>
<i>Hyalithellus micans.</i>	<i>Strenuella salopiensis.</i>
<i>Salopiella obliqua.</i>	<i>Callavia cf. callavei</i> (fragments).

The relations between this fauna and those of contiguous horizons are discussed above (p. 370).

Horizon Ac_3 . *Protolenus* Limestone.

<i>Micromitra (Paterina) labradorica.</i>	<i>Latouchella costata.</i>
<i>Obolus parvulus.</i>	<i>Latouchella striata.</i>
<i>Lingulella viridis.</i>	<i>Microdiscus comleyensis.</i>
<i>Hyalolithus crassus.</i>	<i>Microdiscus lobatus.</i>
<i>Hyalolithus sculptilis.</i>	<i>Microdiscus speciosus.</i>
<i>Hyalolithus (Orthotheca) bayonet.</i>	<i>Protolenus latouchei.</i>
<i>Salterella (?) bella.</i>	<i>Protolenus morphus.</i>
<i>Salterella (?) striata.</i>	<i>Mohicana clavata.</i>
<i>Hyalithellus micans.</i>	<i>Mohicana lata.</i>
<i>Helenia cancellata.</i>	<i>Callavia cf. callavei</i> (fragments).
<i>Helcionella cingulata.</i>	

For discussion of the relationship of this fauna to those below, see above (p. 370).

¹ Cobbold, 1911 b, p. 297.

² The faunas of horizons Ac_1 and Ac_3 were combined under the one heading 'Lobatus Group' in Cobbold, 1911 b, p. 297.

Horizon Ad. *Lapworthella* Limestone.

The fauna of the *Lapworthella* Limestone, Ad, is as follows:—

Species.	Remarks.
<i>Acrothya</i> cf. <i>sera</i>	Confined to this horizon, but represented by a near ally, <i>A. comleyensis</i> , in the <i>Paradoxides</i> Beds.
<i>Lapworthella nigra</i>	Confined to this horizon.
<i>Hyolithellus micans</i>	Ubiquitous, both above and below.
<i>Hyolithellus micans</i> var. <i>robustus</i> .	Now found for the first time, and passing up into the <i>Paradoxides</i> Beds.
<i>Helenia cancellata</i>	Found in the Lower Cambrian, from Ac ₂ , but not above this horizon.
Certain trilobitic (?) fragments, with characteristic tuberculate surface.	Found also in the <i>Protolenus</i> Limestone, but not in the <i>Paradoxides</i> Beds.

The fauna seems to indicate the incoming of a change from Lower to Middle Cambrian. On balance it must be classed with the Lower Cambrian, and the physical evidence points in the same direction.¹

NOTE.—A collection made by Mr. J. Rhodes in 1892, at Neve's Castle in the Wrekin district, for the Geological Survey, contains a number of specimens of a black rock having the lithological characteristics of the *Lapworthella* Limestone, and yielding *Acrothya* cf. *sera*, *Lapworthella nigra*, and *Hyolithellus micans* in similar abundance. The *Paradoxides* and *Olenellus* faunas are known to be in juxtaposition in this area; but the section at the junction is not open, and it is consequently impossible to say whether the *Lapworthella* Limestone of Neve's Castle stands in the same relation to the other beds as it does at Comley.

MIDDLE CAMBRIAN.

Horizon Ba₁. *Paradoxides-groomi* Grits.

<i>Acrothya comleyensis</i> .	<i>Conocoryphe emarginata</i> var.
<i>Hyolithellus micans</i> .	<i>longifrons</i> .
<i>Hyolithellus micans</i> var. <i>robustus</i> .	<i>Dorypyge lakei</i> .
<i>Helcionella oblonga</i> .	<i>Paradoxides groomi</i> .
	<i>Paradoxides</i> spp. indet., 3 species.

The deposits grouped in this horizon consist of three lithological divisions:—(A) An impersistent layer of phosphatic gritty material which adheres to the surface of the *Lapworthella* Limestone, and is rarely more than 1 inch thick; (2) a conglomeratic deposit containing numerous blocks of pre-existing limestone and clots of contemporaneous calcareous material; (3) a coarse grit, with a few similar calcareous clots and blocks of pre-existing limestone.

Horizon Ba₂.

<i>Obolus</i> sp. indet. <i>a</i> .	<i>Hyolithellus micans</i> .
<i>Acrotreta</i> sp. indet. <i>b</i> .	

The paucity of this fauna is attributed to the crushing of the shale-beds and the absence of good exposures.

¹ Cobbold, 1916, pp. 119, 120.

Horizon Ba₃. *Dorypyge-lakei* Flags.

Acrotreta socialis.
Acrothyra comleyensis.

Hyolithellus micans.
Dorypyge lakei.

The three horizons Ba₁, Ba₂, & Ba₃ may be correlated¹ with the *Paradoxides-elandicus* Zone of Scandinavia, in which *A. socialis* is found typically, though it is also recorded from higher zones.

Horizon Bb₁. *Paradoxides-intermedius* Grits.

Billingsella sp. indet.
Hyolithellus micans var. *robustus*.
Microdiscus punctatus.
Conocoryphe æqualis.
Conocoryphe bufo.
Conocoryphe impressa.

Ptychoparia (Liostracus) lata.
Ptychoparia (Liostracus) dubia.
Agraulos cf. quadrangularis.
Strenuella (?) spp. indet.
Dorypyge reticulata.
Paradoxides intermedius.

The relation of this fauna to that of the *Paradoxides-tessini* Beds, of Scandinavia (near the *C. exsulans* sub-zone) and to the *Paradoxides-hicksi* Zone of South Wales has already been discussed.²

Horizon Bb₂.

Obolus sp. indet. b.

Acrotreta sp. indet. c.

Horizon Bb₃.

No exposures available. By position it is inferred that the *Hartshillia* fauna,³ Horizons E₁, E₂, E₃ of the Nuneaton area, if present at Comley, should come here.

Horizon Bb₄. *Paradoxides-rugulosus* Sandstone.

Lingulella ferruginea.
Lingulella sp. indet.

Paradoxides rugulosus.

Paradoxides rugulosus suggests a reference to the Lower *Davidis* Fauna of Nuneaton (Horizons F₁, F₂ of Illing).⁴

Horizon Bb₅. *Paradoxides-davidis* Flags.

Lingulella ferruginea.
Acrotreta sagittalis.
Billingsella cobboldi.
Billingsella lindstræmi var.
Microdiscus cf. punctatus.
Agnostus fallax.

Conocoryphe sp. indet.
Ptychoparia (Liostracus) pulchella.
Ptychoparia (Liostracus) dubia.
Agraulos (?) holocephalus.
Agraulos cf. quadrangularis.
Paradoxides davidis.

This fauna may be compared with those of the *Paradoxides-davidis* horizons of Scandinavia, South Wales, and Nuneaton.

Horizon Bc. *Billingsella* Beds.

Micromitra sp. indet.
Obolus cf. schmalenseei.
Obolus gibbosus.
Obolus (?) tinnarsoni.
Acrothele coriacea.

Acrotreta sagittalis.
Acrotreta schmalenseei var. *matleyi*.
Acrotreta sp. indet. a.
Billingsella lindstræmi var. *salopiensis*.

Eight of these nine species are either identical with, or closely related to, species cited by Dr. C. D. Walcott as occurring in the *Paradoxides-forchammeri* Zone of Scandinavia, with which the Comley *Billingsella* Beds are consequently correlated.

¹ Cobbold, 1913, p. 43.

³ V. C. Illing, 1915, p. 399.

² Cobbold, 1913, pp. 39, 40.

⁴ 1915, p. 400.

UPPER CAMBRIAN.

Horizon Ca. *Orusia* Shales.

Acrothele coriacea.
Orusia lenticularis.

Trilobite-fragments, suggesting one
or more species of the *Olenida*.

Orusia lenticularis occurs plentifully in thin calcareous or mudstone-beds intercalated with shales and, sporadically, in the shale itself. Prof. J. C. Moberg, writing¹ of the *Orusia-lenticularis* Beds of Sweden, says, 'not infrequently we find this brachiopod by itself crowding entire limestone-bands. It occurs very often together with *Parabolina spinulosa*, especially in the shales.' The conclusion that Horizon Ca should be grouped with the Upper Cambrian seems fully borne out.

Horizon Cx.

This designation is used to include the beds which yielded '*Lingulella nicholsoni* and the Shinton graptolites' to Dr. C. Callaway.² His exposure has not been identified. It was situated some 500 yards, more or less, to the north-east of Comley Quarry.

X. GENERAL SUMMARY.

As a result of this study of the Cambrian faunas of Comley, the strata of the Lower Cambrian fall into eleven local divisions, Aa, Ab, Ab₁, Ab₂, Ab₃, Ac, Ac₁, Ac₂, Ac₃, Ac₄, and Ad.

The first defined fauna to invade the area was one comparable with, and in extension of, that found in the quartzite and Hollybush Sandstone of the Malvern Cambrian area. From it the genus *Walcottina*, intermediate between *Mickwitzia* and *Kutorgina*, is now described.

A second incoming fauna is possibly foreshadowed by a few forms in Ab₃, which include a species of *Holmia* or a near ally. For want of brachiopoda and other fossils this fauna cannot be closely defined.

A third faunal group, the *Protolenus-Callavia* Fauna, appears in Horizons Ac₁, Ac₂, Ac₃, Ac₄, and includes a very considerable number of species that find their analogues in the Lower Cambrian of North America. This group is separable into five faunas, not necessarily of equal value, but sufficiently distinct to be worthy of separate record. The three lower faunas in horizons Ac₁, Ac₂, Ac₃ are characterized by an abundance of fragments referable to *Callavia callavei* and other related species, and are separated by the changes that are noted among the smaller trilobites, brachiopods, etc. The two higher faunas in horizons Ac₄, Ac₅, are characterized by plentiful remains of *Strenuella* in Ac₄, and of *Protolenus* in Ac₅. Two genera among the pteropoda and one among the gasteropoda are now described for the first time.

Immediately above the *Protolenus* Limestone the *Lapworthella* Limestone, horizon Ad, occurs: this appears to contain a fauna

¹ 1910, p. 176.

² 1877, p. 659.

somewhat intermediate between those of the Lower and the Middle Cambrian, but still to be classed with the former. As it is also recorded from Neve's Castle in the Wrekin district, where the Lower and the Middle Cambrian are again in juxtaposition, this bed and its fauna seem of interest as a possible index of the dividing-line between the two formations. Further evidence of a transition, if such ever existed in the district, is lost by reason of the unconformity that ushers in the Paradoxidean fauna, which only arrived after a considerable interval.

The Middle Cambrian faunas have already been indicated in previous publications by me; the *Paradoxides-ælandicus* fauna in horizons Ba₁, Ba₂, and Ba₃; the *P.-tessini* fauna in horizon Bb₁, and probably higher zones; the *P.-davidis* fauna in horizons Bb₄ and Bb₅.

The brachiopoda now prove the presence of a fauna of *Paradoxides-forchhammeri* age in horizon Bc, and the Upper Cambrian age of the *Orusia* Shales, horizon Ca, at the summit of the sandstone series of Comley.

Certain points of, perhaps, minor importance result from the detailed study of the brachiopoda. The genus *Acrothyra* appears in the district at the summit of the Lower Cambrian deposits, and its range is extended into the Middle Cambrian. The specimens are in such excellent preservation that it has been possible to determine the forms of the visceral cavity in at least four stages of growth. *Micromitra (Paterina) labradorica* exhibits some features of the growth of the adult shell, and was preceded in time in this district by *M. (P.) phillipsi*. The specimens of *Obolella atlantica* in two varieties show the internal characters distinctly.

I could not have undertaken the study of the Comley brachiopoda without the help of Dr. C. D. Walcott's monograph, which has served as text-book, bibliography, stratigraphical guide, and work of reference, all in one, and I desire to acknowledge the very great assistance which I have received from that comprehensive work. I am also most grateful to Dr. Walcott for the loan and gift of some American specimens.

I am also further indebted to Dr. C. A. Matley for the notes on the brachiopoda of the *Paradoxides* Beds of Comley (1911): they were almost my first lesson in the study of this group of fossils.

At the outset of the excavation work at Comley, Dr. T. T. Groom very kindly examined some of the pteropoda, and his preliminary assistance was of considerable value. Grateful thanks are also due and tendered to Prof. W. J. Sollas, for the opportunity of studying the specimens from Malvern that are in the Oxford Museum. Last, but by no means least, I owe an ever-increasing debt of gratitude to the late Prof. Charles Lapworth for much kind advice as to the scope of this paper and for encouragement throughout its preparation.

TABLE II.—THE FAUNAS OF THE COMLEY HORIZONS.

[illegible]

[illegible]

[illegible]

EXPLANATION OF PLATES XXI-XXIV.

[The numerals in brackets are those attached to the specimens. Those with the letters RR before them refer to specimens collected by Mr. Rhodes for the Geological Survey. The name 'King' before a number indicates that the specimens had been collected by Mr. Wickham King, and presented by him to Mr. Rhodes's collection. A few specimens belong to the British Museum, and all others are from the collection for the British Association, which is in process of being deposited in the Sedgwick Museum, Cambridge.]

PLATE XXI.

Micromitra (Paterina) phillipsi Holl, sp., $\times 4$, see p. 329; horizon Ab₁.

- Fig. 1. Ventral valve, with lateral view [RR 1472].
2. Do. do. with lateral and posterior views [RR 1467].
3. Do. do. internal cast, with lateral view [2334].

Micromitra (Paterina) rhodesi, sp. nov., $\times 4$, see p. 330; horizon Ab₁.

- Fig. 4. Type-specimen, ventral valve; *a*, summit view; *b*, posterior view; *c*, lateral view [RR 1473].
5. Ventral valve, alate form, from a squeeze of an external cast in soft sandstone [2271].
6. Ventral valve with lateral view, shell somewhat flattened and impressed with sand-grains, apex broken, revealing the thickness of shell [RR 1471].

Micromitra (Paterina) kingi, sp. nov., $\times 4$, see p. 330; horizon Ab₁.

- Fig. 7. Type-specimen, ventral valve with lateral view [King 2].
8. Ventral valve; *a*, summit view; *b*, posterior view, showing pseudodeltidium [King 2].

Micromitra sp., $\times 4$, see p. 330; horizon Ab₁.

- Fig. 9. Dorsal valve, with lateral and posterior views, associated with the previous specimens, shell exfoliated [RR 1465].

Micromitra (Paterina) labradorica Billings, $\times 4$, see p. 332;
horizons Ac₁ to Ac₃.

- Fig. 10. The two valves in contact: *a*, ventral; *b*, lateral; *c*, dorsal views [British Museum Collection, B. 40306], Ac₂.
11. Dorsal valve, exfoliated, and showing radiate striae on five successive layers of shell-material [2241], Ac₂.
12. Part of surface of valve in which the rounded concentric bands vary much in width, enlarged to 8 diameters [730], Ac₂.
13. Ventral valve, with lateral and posterior views. The false area is complete, and shows a minute shelf-like pseudodeltidium projecting horizontally under the umbo [2243], Ac₂.

Micromitra (Paterina) minor, sp. nov., $\times 8$, see p. 333; horizon Ac₂.

- Fig. 14. Type-specimen, ventral valve, exterior, with lateral and posterior views [755].
15. Surface of another specimen [RR 1242], $\times 16$.
16. Ventral valve exfoliated at the apex; *a*, summit view; *b*, lateral view; *c*, the exfoliated apex $\times 16$ showing casts of vascular marks and median depression [750].

Micromitra (Paterina) minor, var. *gibbosa* nov., $\times 8$, see p. 334; horizon Ac₂.

- Fig. 17. Ventral valve, summit and lateral views [RR 1117].

Micromitra sp. indet., $\times 4$, see p. 328; horizon Bc.

Fig. 18. Ventral valve, with lateral view [2134].

19. Surface of another specimen [2133] $\times 8$.

Walcottina elevata, gen. et sp. nov., $\times 4$, see p. 337; horizon Ab₁.

Fig. 20. Type-specimen, ventral valve; *a*, internal cast, lateral view [1914];
b, exterior, from a squeeze of the external cast; *c*, lateral view of
same [1916].

21. Dorsal valve [1916], with lateral view.

22. Ventral valve, posterior view of sandy cast [2282].

Walcottina lapworthi, gen. et sp. nov., $\times 4$, see p. 336; horizon Ab₁.

Fig. 23. Type-specimen, ventral valve, exterior; *a*, summit view; *b*, lateral
view; *c*, posterior view [King 1].

24. Ventral valve, exfoliated to the inner layer of shell, with lateral view
[King 1].

25. Surface characters, $\times$ about 12.

Micromitra (?) sp. indet., $\times 4$, see p. 334; horizon Ab₁.

Fig. 26. Interior, with section in outline [2183].

Obolella (?) *groomi* Matley, $\times 5$, see p. 345; horizon Ab₁.

Fig. 27. Ventral (?) valve, exterior, with lateral view [2226].

Bradoria (?) sp., $\times 5$, see p. 368; horizon Ac₂.

Fig. 28. Right valve [14536] in the British Museum (Natural History) Col-
lection; *a*, lateral view; *b*, terminal view; *c*, dorsal view.

PLATE XXII.

Obolella atlantica Walcott, var. *transversa* nov., $\times 5$, see p. 344; horizon Ac₁.

Fig. 1. Type-specimen, dorsal valve, internal cast [810].

2. Ventral valve, internal cast, with lateral outline [807].

3. Do. do. do. [2162].

4. Dorsal valve, exterior [53].

5. Apex of ventral valve, exterior [2189].

Obolella atlantica Walcott, var. *comleyensis* nov., $\times 5$, see p. 342;
horizons Ac₁ to Ac₃.

Fig. 6. Type-specimen, ventral valve, internal cast, with lateral outline
[1288], Ac₃.

7. Dorsal valve, internal cast with portion of shell adhering [785];
also lateral outline, Ac₃.

8. Ventral valve, exterior, with lateral view [2159], Ac₂.

9. Do. do. internal cast, with lateral outline [766], Ac₂.

Lingulella viridis, sp. nov., $\times 6$, see p. 341; horizons Ac₁ & Ac₃.

Fig. 10. Ventral valve, internal cast, with lateral outline [35].

11. Dorsal valve, exterior, with lateral outline [1996].

12. Type-specimen, dorsal valve, internal cast, with lateral view [97].

Obolus parvulus, sp. nov., $\times 6$, see p. 338; horizons Ac₂ to Ac₃.

Fig. 13. Type-specimen, dorsal valve, internal cast, with lateral outline [1512],
Ac₂.

14. Dorsal valve, interior much exfoliated, showing a median ridge
representing the median depression of the exterior [1512], Ac₂.

- Fig. 15. Ventral valve, internal cast, with lateral outline [2200], Ac.₂.
16. Minute ventral valve, internal cast, young and rather wider proportionally than the adult [2198], Ac.₂.
17. Ventral valve, exterior, with lateral outline [760], Ac.₂.
18. Do. do. do. do. [1755], Ac.₂.
19. Dorsal valve, exterior, with lateral outline [1756], Ac.₂.

Obolus (?) *gibbosus*, sp. nov., × 6, see p. 339; horizon Bc.

- Fig. 20. Type-specimen, ventral valve, exterior, with lateral view [2119].
21. Dorsal valve, internal cast, with lateral outline [2120].

Obolus (?) cf. *schmalensei* Walcott, × 6, see p. 340; horizon Bc.

- Fig. 22. Dorsal valve, internal cast; *a*, summit view; *b*, lateral outline [689].

Obolus (?) *linnarssoni*, sp. nov., × 4 and × 6, see p. 338; horizon Bc.

- Fig. 23. Ventral (?) valve, × 4, variously exfoliated, with lateral outline, showing at *a*, portion of exterior; at *bb*, the two raised bands on an inner layer [R 1868].
24. Dorsal (?) valve, × 6, much exfoliated [2385].
25. Minute valve, × 6, doubtfully referred to the species [2399].

Acrotreta schmalensei var. *matleyi*, nov., × 6, see p. 346; horizons
Bc & (?) Bb₃.

- Fig. 26. Type-specimen, ventral valve, exterior, with lateral and posterior views [2066].
27. Ventral valve, exterior, exfoliated and apex removed, with lateral and posterior views [2076].
28. Ventral valve, internal cast, with lateral and posterior views [2109].
29. Dorsal valve, internal cast, with lateral outline [2079].
30. Do. do. exterior somewhat exfoliated, with lateral outline [2078].
31. Dorsal valve, exterior, wider and with surface more strongly marked, doubtfully referred to the species [2103].
32. Surface of a ventral valve, × 12 [2087].

Acrotreta sp. indet. *a*, × 6, p. 347; horizon Bc.

- Fig. 33. Ventral valve, internal cast, with lateral and posterior views [2093].

Acrotreta sp. indet. *c*, × 6, see p. 348; horizon Bb₂.

- Fig. 34. Dorsal valve, internal cast [1621].

Acrotreta socialis Seebach, × 6, see p. 347; horizon Ba.₂.

- Fig. 35. Ventral valve, with lateral and posterior views [1640].

PLATE XXIII.

Acrothyra comleyensis, sp. nov., × 6, see p. 348; horizons Bb₁ & Bb₂.

- Fig. 1. Five drawings of the apical part of a ventral valve [1714], from which successive layers of shell were removed; *a*, exterior; *b*, first exfoliation, showing, so far as the curved line, the internal cast of a young shell; *c*, second exfoliation, showing internal cast of a small shell attaining adult features; *d* and *e*, further exfoliations, showing further stages of growth.
2. Type-specimen, ventral valve, internal cast [1710], from which all the shell has been removed, with lateral and posterior views.

- Fig. 3. Ventral valve, internal cast [1708], with posterior view; some shell remains on the cast.
4. Ventral valve, internal cast [1709], with lateral view; much shell remains on the cast.
  5. Dorsal valve, internal cast, with lateral and posterior views [1707].
  6. Ventral valve, interior [1711].
  7. Do. do. do. [1706], the visceral cavity more rotund than usual.
  8. Ventral valve, internal cast, young individual, with lateral and posterior views. The visceral cavity is somewhat rotund [1716].

Acrothyra cf. *sera* (Matthew) Walcott, × 6, see p. 350; horizon Ac.

- Fig. 9 a. Ventral valve, exfoliated slightly, with lateral and posterior views [1].
- 9 b. The same valve, from which all the shell has been removed, with lateral view.
  10. Ventral valve, interior [2218].
  11. Do. do. lateral view, posterior slope inclined backwards [1800].
  12. Ventral valve, lateral view, posterior slope vertical [1799].
  13. Do. do. exterior of apical portion [2207].
  - 14 a. Dorsal valve, strongly exfoliated, with lateral outline [2207].
  - 14 b. The counterpart, showing many laminae [2208].
  15. Dorsal valve, internal cast, with lateral and posterior views [1801].
  16. Smaller dorsal valve, partial exterior, with lateral view [1801].
  17. Dorsal valve, internal cast, with lateral and posterior views, showing a greater extension of the median septum [1781].
  18. Dorsal valve, exterior, with lateral view, much wider than usual and showing strongly-marked sculpture [18].
[The specimens represented by figs. 17 & 18 are referred doubtfully to the species.]

Acrothele coriacea Linmarsson, × 6, p. 345; horizons Bb., Bc, & Ca.

- Fig. 19. Ventral valve, partly exfoliated. The surface of the shell is seen near the posterior margin where the sculpture is linear, towards the apex it is granular; a, summit; b, posterior view; c, lateral view [2345].
- 19 d. Exfoliated interior of the same shell [2346].
  20. Ventral (?) valve, fragment of external surface [2349].
  21. Dorsal valve, exfoliated, but showing two portions of the exterior, with lateral outline [2352].
  - 22 a. Dorsal valve, exfoliated, showing one of the two apical tubercles and a slight median depression [2379].
  - 22 b. The same specimen, after much shell-substance had been removed.

Billingsella sp. indet., × 3, see p. 354; horizon Bb₁.

- Fig. 23. Ventral valve, exterior from a squeeze of the cast; a, summit view; b, lateral outline [1880].
- 23 c. Internal cast of the same [1879].

Billingsella lindstræmi, var. *salopiensis* Matley, × 3, see p. 353;
horizons Bb₅ & Bc.

- Fig. 24. Ventral valve, exterior from a squeeze of the cast; a, summit view; b, lateral outline [2450], Bc.
- 24 c. Internal cast of the same [2449], Bc.

Orusia lenticularis Wahlenberg, $\times 4$, see p. 354; horizon Ca.

- Fig. 25. Dorsal valve, internal cast, with posterior view, showing casts of convergent crural plates [2509].
26. Ventral valve, internal cast, showing casts of divergent dental plates [2507].
27. Ventral valve, exterior, with lateral outline; and dorsal valve, internal cast, showing casts of parallel crural plates [2516].

Oboius sp. indet. *b*, $\times 6$, see p. 340; horizon Bb₂.

- Fig. 28. Ventral valve, internal cast [1621].

Oboius sp. indet. *a*, $\times 6$, see p. 340; horizon Ba₂.

- Fig. 29. Dorsal (?) valve, external cast [1604].

Acrotreta sp. indet. *b*, see p. 348; horizon Ba₂.

- Fig. 30. Ventral valve, crushed exterior [1606].

PLATE XXIV.

Lapworthella nigra, gen. et sp. nov., $\times 10$, see p. 360; horizon Ad.

- Fig. 1. Exterior of attenuate example, slightly damaged [RR 1221].
2. Internal cast, with shell in section [266].
3. Surface of inner shell, showing (where broken away) a portion of the internal cast [RR 1221].
4. Exterior of wide example, somewhat worn [RR 1221].
5. Interior of outer covering, with portion of inner shell adhering [267].
6. Wide individual, detached from the matrix and somewhat abraded: *a*, summit; *b*, lateral view [270].

Helenia cancellata, sp. nov., $\times 4$, see p. 363; horizons Ac₂ to Ad.

- Fig. 7 *a*. Type-specimen, external impression, with closed end preserved [246], Ac 2.
7 *b*. Same individual, showing marginal groove [245], Ac₂.
8. Another closed end [243], Ac₂.
9 *a*. Another specimen $\times 4$ [246], Ac₂.
9 *b*. The same $\times 10$, with section.
10. Diagram, showing character of sculpture.

Hyolithus (Orthotheca) bayonet, var. *striatus* nov., $\times 4$, see p. 357; horizon Ac₄.

- Fig. 11. One of the ventral faces, with section [231].

Hyolithus (Orthotheca) bayonet Matthew, $\times 4$, see p. 357; horizon Ac₂.

- Fig. 12. Dorsal face [2567].
13. Dorsal apertural margin, external cast [2572].
14. Dorsal face, fragment, with section [2568].
15. One of the ventral faces [2576].
16. Section, more in accordance with Dr. Matthew's description than fig. 14 [1672].

Hyolithus (Orthotheca) compressus, sp. nov., $\times 4$, see p. 357; horizon Ac₂.

- Fig. 17. Type-specimen; *a*, part of an external cast, with outline of curvature [225]; *b*, exterior, with section [226].

Salopiella obliqua, gen. et sp. nov., $\times 4$, see p. 363; horizon Ac_3 .

Fig. 18. Type-specimen [242]; *a*, internal cast or exterior (?), $\times 4$;
b, enlargement of surface, $\times 10$; *c*, section, $\times 10$.

Hyalithellus micans Billings, $\times 4$, see p. 361; horizons Ac_1 to Ba_3 .

Fig. 19. A curved example [252], Ad.

20. A sinuous example [251], Ad.

21. A straight example [1778], Ad.

Hyalithellus micans, var. *robustus* nov., $\times 4$, see p. 362; horizons Ba_1 to Bb_1 .

Fig. 22. A long fragment, with section [1695], Bb 1.

Hyalithellus micans, var. *pallidus* nov., $\times 4$, see p. 362; horizons Ab_2 & Ab_3 .

Fig. 23. A fragment [1261].

Hyalithellus (?) *sinuosus*, sp. nov., $\times 10$, see p. 362; horizon A_1 .

Fig. 24. Type-specimen [1912].

Hyalithus sculptilis, sp. nov., $\times 4$, see p. 356; horizon Ac_3 .

Fig. 25. Type-specimen [2587], with section.

26. Fragment of shell [2585].

27. Character of sculpture [2583], $\times 6$.

Hyalithus crassus, sp. nov., $\times 2$, see p. 354; horizon Ac_3 .

Fig. 28. Type-specimen, dorsal face, lateral view and section [2603].

Hyalithus sp. indet., operculum, $\times 4$, see p. 356; horizon Ac_3 .

Fig. 29. Complete specimen, with section [2655].

Hyalithus strettonensis, sp. nov., $\times 2$, see p. 355; horizon Ab_1 .

Fig. 30. Type-specimen, internal cast, lateral view with section [1903].

31. Dorsal face, external cast [1902].

Hyalithus strettonensis, var. *brevis* nov., $\times 4$, see p. 355; horizon Ab_1 .

Fig. 32. Dorsal face [220].

Salterella (?) *bella*, sp. nov., $\times 10$, see p. 360; horizon Ac_3 .

Fig. 33. Type-specimen, lateral view [523].

Hyalithus (*Orthotheca*) sp. indet., operculum, $\times 4$, p. 358; horizon Ac_3 .

Fig. 34. Exterior [2655].

Hyalithus (*Orthotheca* ?) sp. indet., operculum, $\times 4$, see p. 358; horizon Ac_3 .

Fig. 35. Internal cast, with section [2656].

Scenella elevata, sp. nov., $\times 6$, see p. 364; horizon Ac_2 .

Fig. 36. Type and only specimen, summit and lateral views [RR 1211].

Helcionella cingulata, sp. nov., $\times 4$, p. 364; horizon Ac_3 .

Fig. 37. Type-specimen, incomplete, lateral view and section [827], Ac_3 .

Helcionella oblonga, sp. nov., $\times 2$, see p. 365; horizon Ba_1 .

Fig. 38. Type-specimen, lateral and summit views [1316].

39. Sculpture, $\times 5$ [1314].

Helcionella rugosa Hall, var. *comleyensis* nov., $\times 4$, see p. 365 ;
horizons Ac_1 & Ac_2 .

Fig. 40. Type-specimen, lateral and summit views [1897].

Latouchella costata, gen. et sp. nov., see p. 366 ; horizons Ac_2 & Ac_3 .

Fig. 41. Type-specimen [in the collection of the Rev. W. M. D. La Touche],
internal cast, $\times 5$, Ac_1 .

42. Second specimen, exterior, lateral & summit views, $\times 2$ [2682], Ac_3 .

Latouchella (?) *striata*, sp. nov., $\times 2$, see p. 367 ; horizon Ac_3 .

Fig. 43. Type-specimen, lateral and terminal views [2655].

44. Another specimen, lateral, summit, and terminal views [2659].

Salterella (?) *striata*, sp. nov., $\times 6$, p. 361 ; horizon Ac_3 .

Fig. 45. Type-specimen, lateral view and outline of aperture [139].

[By misadventure this figure has been reversed in position,
giving the appearance of a concave instead of a convex object.]

Leperditia lentiformis, sp. nov., $\times 2$, see p. 367 ; horizon Ac_2 .

Fig. 46. Type-specimen [51 in the Birmingham Museum], two valves in
position : *a*, left side ; *b*, dorsal aspect ; *c*, surface $\times 6$.

Escasona (?) spp. indet., $\times 6$, see p. 368 ; horizon Ac_2 .

Fig. 47. [RR 1332], with profile outlines.

48. [RR 1335], with do. do.

E.S.C. del.

Bernrose, Colla, Derby.

E.S.C. del.

Bernrose, Colla, Derby.

E.S.C. del.

Bemrose, Calla, Derby.

E.S.C. del.

Bemrose, Callia, Derby.