TRANSACTIONS
OF THE
ROYAL ACADEMY OF MEDICINE IN IRELAND.

SECTION OF MEDICINE.

A TRIAD OF CLINICAL RECORDS.

By SIR J. MOORE, M.A., M.D., M.Ch., D.P.H. Dubl.;
Senior Physician to the Meath Hospital.

[Delivered to the Section of Medicine, November 10, 1911].

AN apology on my part is due, and is herewith tendered, to the Fellows of the Royal Academy of Medicine in Ireland for trespassing upon their time and attention with a series of mere clinical notes. But the cases which I desire to report made a profound impression upon me, and their story may prove of some value, at all events to the younger members of the Academy, as a guide to diagnosis and treatment.

CASE I.—*Chronic Perihepatitis and Interstitial Hepatitis—Ascites, Jaundice, and Hæmatemesis.*—Mrs. Teresa O’K., aged thirty-four years, a clerk’s wife, was admitted to the Meath Hospital under my care on April 8, 1909, with the signs and symptoms of purpura hæmorrhagica. She complained of a feeling of fulness in the abdomen, which was distended, and of a constant drowsiness.

Her personal history was not very eventful. At the age of ten years she passed through scarlatina, and three years later through enteric fever. She was married at twenty three, and bore five children, of whom four are alive. The fifth and last was stillborn in the year 1906.

When admitted, Mrs. O'K.'s pulse was 100, her respirations were 20, and her axillary temperature was 99.2° . During her stay of six weeks in hospital her condition was almost continuously subfebrile in the evenings—the highest reading recorded was 101° on the evening of May 2.

To illustrate the hæmorrhagic tendency which was present, I may mention that on April 11 the patient scratched a spot on her right cheek, with the result that there was free hæmorrhage. It was checked by the application of solution of ferric chloride. On April 28, the urinary examination gave the following result:—Quantity, 32 fluid ounces passed in twenty-four hours. Of a yellowish brown colour, acid reaction, spec. grav. 1018. Albumen and blood absent. No sugar. Excess of mucus.

The measurements of the abdomen at the level of the umbilicus were on April 30, 40 inches; May 8, $39\frac{1}{2}$ inches; May 14, $38\frac{1}{2}$ inches; May 20, $36\frac{1}{4}$ inches. She left hospital on the last-named day. While under observation her weight increased from 9 st. $10\frac{1}{2}$ lbs. on April 26, to 10 st. on May 3. It then decreased with the absorption of ascitic fluid to 9 st. 8 lbs. on May 10, only a week later.

The patient left the Meath Hospital, relieved, on May 22, 1909. She was subsequently treated at the Coombe Hospital, where paracentesis abdominis was performed. The fluid, however, accumulated again shortly after she returned home.

Mrs. O'K. came under my care for a second time in January of the present year. On the 25th of that month she entered the Meath Hospital, deeply jaundiced. Her breath was very foetid, owing partly to carious teeth, partly to a spongy state of the gums, which bled freely. There were a few spots of purpura on the body. She stated that her monthly periods had been irregular for some time, that she had been passing less water than usual, and that her skin was very itchy. She was wasted, but the

abdomen was much distended, the umbilical depression being almost obliterated. The superficial veins were dilated over the chest and abdomen. The flanks bulged when she lay on her back, and the other physical signs of ascites were present. She had a craving for alcohol, which had evidently been gratified by her husband and mother through mistaken kindness since her former stay in hospital. Periodical raving occurred at night. Both pulse and respirations were persistently quick. The axillary temperature was unsteady—the readings were generally slightly above normal, though not so high as they had been in 1909. On the evening of February 14, however, 101.4° was recorded. The urine was examined, but was normal save for the presence of bile. There was a fine tremor in the hands. The knee-jerks were present. On February 3, the girth round the abdomen at the level of the umbilicus measured $43\frac{1}{2}$ inches. Paracentesis abdominis was performed on February 14, when 440 ounces of bile-stained serum were drawn off. The specific gravity of the fluid was 1020. It did not contain much fibrin or show other evidence of inflammation. Her weight after the tapping dropped from 10 st. 3 lbs. to 7 st. 10 lbs.

She left hospital on March 7, 1911, and I lost sight of her until Friday, October 6, 1911, when she came into hospital for the third and last time, with all the signs—subjective and objective—already mentioned present in an aggravated form. On admission, the axillary temperature was 98.8° ; the pulse-rate was 108, and the respirations were 24. The night raving now amounted almost to mania. The abdomen was enormously distended, the protruding umbilicus forming a tumour as large as a tennis ball. The gums were bleeding, and she suffered from melæna. To quiet her at night, dilute hydrobromic acid was given, and to check the tendency to bleeding calcium lactate was exhibited in full—15-grain—doses.

The embarrassment to the heart's action and breathing became so great that paracentesis was done on October 10, no less than 580 ozs. of fibrino-serous exudation being drawn off.

Next day it was possible to examine and map out the liver by palpation and percussion. It was found to be much

enlarged, projecting into the abdomen in the nipple line to the extent of $1\frac{1}{2}$ to 2 inches. A hard nodule, or boss, was felt to the left of the xiphoid cartilage, and the surface of the liver was granular or nodular in a marked degree, presenting the typical hob-nail feel of cirrhosis. Hæmatemesis and melæna now became frequent, and the patient gradually sank, and died early on the morning of St. Luke's Day.

The *post-mortem* examination was made a few hours after death by my colleague, Dr. Boxwell, who reports as follows on the findings:—

“ There was very remarkable thickening of the capsules of the liver and spleen. The free surfaces were deeply pitted and indented. These pits could be distinctly felt through the abdominal wall, after removal of the fluid, and the sensation conveyed strongly suggested the irregularities of the typical ‘hob-nail’ liver. A glance at the specimen was enough to show that it was nothing of the kind. The irregularities were merely local depressions in the enormously thickened capsule, and did not extend into the underlying liver substance, the surface of which was comparatively smooth.

“ The liver, however, showed microscopically a cirrhosis of an extreme degree. The new fibrous tissue cut through individual lobules in all directions, and contained multitudes of the so-called new-formed bile-ducts. A dissection of the larger biliary passages and the portal vein failed to disclose any gross obstruction.

“ After a section of the upper free surface of the liver both air and blood could be freely expressed through the portal vein, and a probe passed easily through the common bile-ducts and appeared in the duodenum.”

In the treatment of this case I did not try subcutaneous injections of fibrolysin. The *post-mortem* findings leave no room for regret in regard to this. Arguing from the fact that in interstitial hepatitis the function of the liver is disturbed by constricting bands analogous to cicatrical tissue, Moerlin, in an article entitled “Fibrolysin bei Hepatitis interstitialis,” published in the *München med. Wochenschrift* in 1908 (Vol. LV., July 7, page 1435), and

abstracted in *The Hospital*, Oct. 17, 1908, administered fibrolysin to a patient suffering from cirrhosis of the liver, injecting 2 cc. of the drug subcutaneously in the region of the liver two or three times a week. The affection was in the atrophic stage with commencing ascites. Iodide of potassium and calomel had been tried in succession, but had proved useless. The patient stood the fibrolysin injections well. Shortly after the second injection he felt better, the output of urine was increased and the ascites diminished. The man was subsequently able to leave his bed, with improved appetite, and, after six injections, was able to resume work. Moerlin found that the liver had regained its normal size. In the course of treatment the patient was given Carlsbad salts and hydrochloric acid, but the great improvement was undoubtedly due chiefly to the fibrolysin. I understand that ampullæ are supplied for hypodermic or intramuscular injection containing 2.3 cc.s. apiece (nearly 40 minims of a solution of one molecule of thiosinamin and half a molecule of sodium salicylate).

CASE II.—*Left Pleuro-pneumonia—Empyema—Thoracenteresis—Recovery.*—On Saturday, July 8th, 1911, A. B., a lad of eighteen years, bathed while over-heated and tired. A "chill" followed, and he developed a fever which was regarded as influenza. He progressed favourably until the following Wednesday (July 12th), when he again became feverish, and Dr. Albert Croly detected the physical signs of a left pleuro-pneumonia. I was asked to see the patient in consultation with Dr. Croly on Saturday, July 15, the eighth day from the initial chill and the fourth of a declared pleuro-pneumonia. At this time the patient's tongue was thickly coated, there was a heavy smell from his breath, the abdomen was very tympanitic, though the bowels were regular. He had a troublesome cough, which caused left side pain. With some effort he brought up a viscid sputum, which only slightly partook of the rusty character. The

pulse-rate was 96, respirations were 28 per minute, and the temperature in the mouth was no higher than 101.5° . The percussion note was hyper-resonant over the apex of the left lung, and the respiratory sound was rough and puerile, or compensatory. Posteriorly, there was a moderately large patch of dulness on percussion just below the scapula. Here also some fine crepitations were to be heard, with a distinct friction rub below. In front the percussion note was abnormally clear up to the fifth rib, as if the stomach was distended with gas, as I think it was.

What I did not like about the case was the trifling pyrexia on the fourth day of pneumonia, and also the presence of tympanites. These facts influenced me in replying to the lad's mother, who asked "When would the crisis occur?" I answered: "The case is not a straightforward one of pneumonia, or pneumonic fever, and we cannot look for a 'crisis.' " As a matter of fact, the fever dragged on, after two days of apyrexia, while remaining of moderate intensity. Dr. Croly very properly had the sputum examined bacteriologically for tubercle bacilli, but with a negative result. Nor did the blood give a positive Widal reaction. Our fears as to a possible tuberculosis or enteric fever were consequently allayed, but the patient's state remained unsatisfactory, and, indeed, steadily grew worse. He lost flesh and strength. His heart beat more rapidly, and the rate of the breathing rose to 40 or 44 in the minute. Dr. Croly shrewdly made a diagnosis of an encysted empyema on the left side.

On Sunday, July 30, I visited the patient, who was passing through his illness in the country a few miles south of Dublin—the main object of our consultation being to decide whether he could safely be removed to the epidemic wing of the Meath Hospital. Although at this time he was even gravely ill, the physical state of his heart seemed to justify his removal, and this was effected in an ambulance the following day, Monday, July 31. In the evening of this day the temperature rose to 103.2° —the highest recorded during the whole illness—the pulse was 98, and the respirations were 40.

On the afternoon of Tuesday, August 1, the patient's state

was critical. He had paroxysmal fits of coughing, which caused extreme cyanosis, and so also did any change of position, especially to the right side. His breath was foetid, so was the sputum, and the foetor on coughing was like that of pulmonary gangrene. In the second and third left intercostal spaces an area of dulness on percussion existed, and the other physical signs suggested the formation of a gangrenous cavity in the lung. In the afternoon Dr. Alfred Parsons joined Dr. Croly and me in consultation, my hospital colleague, Dr. Lennon, being out of town. We agreed that there was no indication for thoracentesis at the time having regard to the patient's very weak condition, and it was arranged that the very active treatment which I had adopted in the morning should be continued. Oxygen inhalations, hypodermic injections of one-sixtieth of a grain of sulphate of strychnine, and one-hundredth of a grain of digitalin, were given alternately, each at intervals of six hours. Ten minims of turpentine were taken in a little whisky and hot water every fourth hour. A teaspoonful of Jenner's syrup (*liquor morphinæ*, *aqua laurocerasi*, and *syrupus aurantii*) was taken when cough was troublesome, and 3 grains of salicylate of quinine were given every six hours.

A remarkable change in the physical signs now took place. On the morning of August 3, the greater part of the left side yielded an absolutely dull percussion note, and the heart was seen and heard to beat just outside the sternum on the right side. The dextrocardia had occurred with startling suddenness. It was as if an encysted pleural effusion had escaped into the left pleura at large. I felt that evacuation of the fluid was imperative if the patient's life was to be saved, and accordingly I asked my colleague, Mr. William Taylor, who was on duty in the hospital at the time, to see the patient with a view to operation. He kindly came at once. An exploratory puncture gave exit to foul-smelling pus. Without a moment's delay a drainage tube was introduced, and large quantities of abominably foetid pus were evacuated. The patient's weak state and alarming cyanosis forbade resection of a rib being attempted, although it was open to question whether the drainage tube would continue

to act owing to the close approximation of the ribs. However, our fears were not realised, and drainage went on satisfactorily.

On the day after thoracentesis, 20 cc.s of polyvalent anti-streptococcal serum were injected. On August 5, the pulse fell to 100 from a maximum of 126. The respirations remained fast—40 to 44 per minute. The temperature gradually fell, spiking in the evenings, and remained subnormal for three days from August 14. It then rose again for no apparent reason, spiking for a week, and at one time touching 102.2° . From August 27 to September 14, it was nearly always subnormal, only on September 3 rising to 99.2° in the evening. The sputum became less and less offensive, lessened in amount, and finally almost ceased.

This brief clinical sketch conveys no idea of the perilous and critical state of the patient up to the time that thoracentesis was performed and of the determined struggle for life which he made before and after that operation. He had been a life-long total abstainer, but he drank champagne freely and took teaspoonful doses of whisky in the form of turpentine punch for many days after August 1. All stimulants were stopped long before he left hospital. They were prescribed and taken as any other medicines, and they did the patient good.

A. B. left hospital for the country on September 14. He continued to progress favourably, save for a chill and a passing catarrh in the middle of October, and at the time of writing was apparently in good health.

CASE III.—*Mediastinal Lympho-sarcoma involving the Œsophagus, Stomach, and Liver, and compressing the Trachea.*—At Whitsuntide in the present year (June 5th) Mr. X., a professional man, then in his seventy-first year, failed to recognise the identity of his own son when he saw him in the garden of his house. Two months previously he had experienced great difficulty in filling up the Census

paper, owing to lapse of memory even as to the number of his children. In the light of subsequent events his family regarded this incident as the earliest sign of impaired health.

On Tuesday evening, June 13th, Mrs. X. wrote asking me to see her husband, who seemed to be losing his memory as well as his appetite. Her message ran as follows:—"I should be very glad if you could manage to call on Wednesday morning sometime between ten and eleven o'clock to see my husband. I am most uneasy about him. He is far from well, and I do not think he is fit for his various duties. If you could see him before he went out I know your advice would weigh with him." I visited him next morning. On examination both heart and lungs appeared to be sound. The liver was somewhat enlarged. His tongue was coated. The urine was examined next day—it was clear, acid in reaction, of moderate density, free from albumen. It was high-coloured owing to the presence of bile-pigment. There was no undue arterial tension.

The mental condition was unsatisfactory. Mr. X. seemed dazed, could not remember dates or names, but expressed himself as feeling quite well. As a tonic I ordered him half drachm doses thrice daily of Easton's syrup, and suggested lemon-flavoured sanatogen as a help in his dietary. Acting under my advice, Mr. X. sought and obtained leave of absence from his professional duties, and left town in the middle of July for a change. Early in August he was home again. On August 10th his daughter wrote asking me to call the following morning to see her father, "who seems poorly to-day. He complains of feeling cold, and was sick in his stomach, but could not be induced to remain at home. We took his temperature, but it was up only two points."

Next morning, on calling, I found Mr. X. at breakfast. After a moment or two, he rose from his chair and hurriedly left the room. On returning after a few minutes he apologised, and explained that he was obliged to rush out, and had lost his breakfast in a sudden attack of vomiting without nausea. After examining him I prescribed a simple mixture of solution of bismuth citrate, and tincture of nux vomica. This he took for some days, apparently with benefit.

On August 21st, Dr. Percy Kirkpatrick kindly saw Mr. X.

in consultation with me, and we agreed that he was doing himself harm by going out in the evenings and sitting up late at night, as had been his habit. During the next fortnight the patient had several attacks of œsophageal spasm (œsophagismus) on attempting to swallow food.

On the evening of September 7th I received the following message from Mrs. X.:—"I shall be very glad if you will kindly come and see my husband to-morrow or next day. He *lost his voice some days ago*, and he has had a return of the sickness for which you prescribed some time since." I visited next morning, and found the patient voiceless, speaking only in a whisper. Believing that he had contracted a catarrhal laryngitis, I ordered an inhalation of 20 drops of a mixture of compound tincture of benzoin (4 parts), menthol (1 part), and creosote (1 part), floating on hot water 140° F.), to be used three times a day.

The following Thursday, September 14th, Mr. X. came to to see me in my consulting-room. His laryngeal symptoms persisted, and I advised him to drive straightway to Dr. R. H. Woods, President of the Royal College of Surgeons in Ireland, who had advised him on a former occasion for some throat affection. This he did, and the same afternoon I had the following letter from Dr. Woods:—"34 Merrion Square, E., Dublin, 14th September, 1911.—Dear Sir John, I examined Mr. X., and find he has paralysis of the right recurrent laryngeal nerve and compression of the trachea fairly low down. The right side of his neck is fuller than normal, and the veins on the right side of the chest are dilated. He has, in my opinion, either an aneurysm or a new growth in the right upper region of the chest. I did not examine his chest directly, but from collateral evidence I should be more inclined to expect an aneurysm than a neoplasm.—Yours faithfully, R. H. Woods."

At a subsequent interview Dr. Woods told me that he had detected slight tracheal tugging.

On the morning after the receipt of this most valuable opinion I made a careful physical examination of the patient's chest and circulation. The supra-sternal notch was abolished. There was an area of visible pulsation just above the right clavicle, and in this situation also the heart's sounds

were very audible. No very marked dulness on percussion was present in the right sub-clavicular region or in the upper sternal region. While the radial pulses, right and left, were synchronous, the right was markedly smaller in volume than the left. The arterial pressure tested by Potain's sphygmomanometer was 11 centimetres of mercury in the left radial, but only $10\frac{1}{2}$ centimetres in the right. I may mention that this difference lessened as the disease advanced and as the interference with the systolic wave came to involve the left subclavian artery as well as the right.

With the view of clearing up the differential diagnosis between an aneurysm of the aorta, or innominate artery, and a new growth, I had a radiograph taken in the X-ray room at the Meath Hospital on Saturday, September 16th. The position and extent of the shadow were perhaps in favour of a neoplasm rather than of an aneurysm.

At this time violent fits of coughing and distressing spasms interfered with the act of swallowing, and Mr. X. began to subsist on fluid food alone, and that taken in only small quantities at a time. For the relief of cough the official guaiac resin lozenges and morphine lozenges were prescribed, but the remedy which gave most relief was the sipping of warm water to which a little glycerine and lemon-juice were added. A copious bleeding also occurred about this time—it was not easy to say whether it constituted a hæmatemesis or a hæmoptysis. Sometimes a blood-stained (grumous) sputum came up with the cough. On exertion great breathlessness ensued.

On Friday, September 29th, Dr. Finny very kindly saw Mr. X. in consultation with me. At this time a large swelling existed in the neck on the right side, and the trachea was seen to be bent over to the left by pressure from the mass. Dr. Finny pointed out to me that the tumour was too solid and resistant for an aneurysm; also that the area of pulsation lay below it, and that the pulsation was not excentric. Furthermore, there was now little or no difference in the rhythm or volume of the radial pulses. He expressed a definite opinion in favour of a neoplasm, and approved the treatment upon which I had put Mr. X. on September 23rd—namely, increasing doses of iodide of potassium and

occasional inhalations of nitrite of amyl. I had prescribed these remedies at an early morning visit paid in answer to an urgent note from Mrs. X., dated Saturday, September 23, 7 34 a.m., to this effect: "Will you kindly come to see my husband this morning. He had a terrible attack of breathlessness last night, and we were greatly alarmed."

Dr. Finny's opinion carried conviction to my mind that we had to deal with a new growth rather than with an aneurysm. We agreed that inhalations of the "A. C. E." mixture (alcohol, 1 part; chloroform, 2 parts; and ether, 3 parts) might relieve the dyspnoea and spasms, which were daily increasing in violence, and these inhalations did give much relief during the terrible fortnight of suffering which followed. Dr. Finny again saw the patient during my absence from town on the next succeeding Sunday, October 1st, for it now became necessary to visit Mr. X. once or twice daily.

On Tuesday, October 3rd, Dr. Woods examined the larynx while the patient was in bed. He demonstrated to me most beautifully the complete paralysis of the right vocal cord. He considered that the growth was developing quickly, and gave a very unfavourable prognosis.

For the next fortnight oxygen inhalations did much to relieve the sufferer, and, indeed, to keep him alive. Day by day it became more and more clear that the sands of life were fast running out.

When I visited Mr. X. on Thursday morning, October 12th, I was surprised to find him sitting up in bed and smoking a pipe. As I entered his room he whispered, "I have bucked up a bit." Afterwards I learned that earlier in the morning he had told the nurse in attendance that he had only two days more to live. On Friday and Saturday he was manifestly sinking. A dreadful night of struggle for breath followed. On Sunday morning I was sent for. His hands were now cold. To relieve the urgent dyspnoea I injected hypodermically one-sixth of a grain of morphine and one-thirtieth of a grain of strychnine sulphate. This gave ease, and the end came at half-past three o'clock in the afternoon without further suffering.

On Monday, October 16th, Dr. Boxwell, assisted by Mr.

Robert Weldon, made a partial autopsy in my presence. Appended are Dr. Boxwell's notes:—

“ Only a partial *post-mortem* examination was possible. The growth in the neck, presenting a tense elastic tumour, was first explored. In the process of dissecting the anterior triangle, the scalpel suddenly dipped into a large cavity underlying the distended cervical fascia, and from the opening made fluid resembling pus welled up. The cavity extended straight down into the mediastinum, and was caused by the liquefaction of the cervical and posterior mediastinal glands, which had been matted together, forming an ill-defined mass surrounding the innominate artery and its branches. The glands at the root of the lungs were similarly involved, and on making an opening in the diaphragm it was clear that the glands along the lesser curvature of the stomach were extensively enlarged, while several white masses protruded from the surface of the exposed liver. The left lobe of this organ was removed, and was found to be thickly studded with secondary deposits. While the heart and lungs escaped altogether, the œsophagus had been invaded—apparently from without—almost throughout its length, and where the masses of new growth had broken through the mucosa there were extensive bleeding and fungating ulcers. The spleen was very small, and entirely free from deposits.

“ A remarkable point about this growth was that almost every deposit, however small, was partially liquefied, and it was difficult to find portions of it firm enough to cut.

“ Microscopically the appearance suggested ‘small round-celled carcinoma.’ That is, the cells were those of a small round-celled sarcoma, whereas their method of growth along the lymphatics apparently in solid masses, recalled the distribution and alveolar stroma of a cancer. In the section from the liver the dense fibrous alveolar stroma was particularly evident. I am inclined to look upon it as a variety of lympho-sarcoma. It was certainly not a squamous cancer, and if the cells were epithelial, it was not easy to see whence they originated.”

By an interesting coincidence two cases of sarcoma of

the thymus were reported in the *Lancet* for November 4, 1911, conjointly by Mr. William Sheen, Surgeon to the Cardiff Infirmary, Mr. Cornelius A. Griffiths, Assistant Surgeon to that Infirmary, and Mr. H. A. Schölberg, Senior Pathologist to the same institution. Both cases occurred in young males, aged seven and eighteen years respectively. In both the onset was acute, and dyspnoea, at first in attacks, was very severe. The growths were of an identical nature in both cases. The morbid anatomy and microscopical appearances were those of a small, round-celled sarcoma. The pressure symptoms resembled those observed in my unfortunate patient.

But the case which most resembled that now reported was one which I brought before the Pathological Section of this Academy on Friday, March 25, 1898, under the heading "Cancer of the Œsophagus with Secondary Growths perforating the Trachea and the Right Subclavian Artery." ^a In that case the absence of any constitutional disturbance, and the presence of a whole group of pressure symptoms, led me to the belief that the lesion was most likely an aneurysm of the terminal portion of the arch of the aorta pressing backwards and to the left. The laryngeal stridor with paralysis of the left vocal cord pointed to pressure on the recurrent laryngeal nerve; the dysphagia, to pressure on the Œsophagus; the bronchial catarrh and dyspnoea, to pressure on the left bronchus; the recurring foetid breath and expectoration, to pressure on the nutrient vessels of the left lung. The physical signs of thoracic aneurysm were absent, but then let us recall to mind Sir William Broadbent's division of thoracic aneurysms into two classes according to the predominance of physical

^a Transactions of the Royal Academy of Medicine in Ireland. Vol. XVI. 1898. P. 355, *et seq.* Dublin Journal of Medical Science. Vol. CV. Third Series. No. 317. May, 1898. P. 396.

signs or of pressure symptoms respectively, the former term applying to aneurysms of the ascending aorta and first part of the arch, the latter to aneurysms of the transverse and descending portions of the arch.^a

Writing in 1854, Dr. William Stokes observed that, an intra-thoracic tumour having been discovered, the diagnosis will be between aneurysm and cancer. He showed that a cancerous tumour may not only produce all the symptoms which we observe in aneurysm as proceeding from compression of surrounding organs, but that it may exhibit a diastolic pulsation, accompanied by a distinct bellows murmur. He stated, however, that as aneurysm is so much more frequent than cancer, the chances will be greatly that in any given case of intra-thoracic tumour the disease is aneurysmal.^b

---

MR. WOODS said he had seen the third of these cases in consultation. The patient had a peculiar stridor, which the trained ear could detect as being due to stenosis or to compression of the trachea. The pitch of such stridor varied with the situation of the narrowing. He had seen another case lately with a similar stridor—a boy of thirteen suffering from sarcoma of the thyroid. The first case of the kind he ever saw was one, in Mercer's Hospital, where a gummatous mass compressed the wind-pipe.

DR. W. G. SMITH expressed his surprise at Sir John Moore attaching any importance to "cancerous cachexia." The speaker entirely refused to believe that cancer produces cachexia *per se*, and instanced many uterine and breast cancers of enormous size where no cachexia existed.

^a Heart Disease and Aneurysm of the Aorta. By Sir William H. Broadbent, Bart., K.C.V.O., and John F. H. Broadbent, M.A., M.D. (Oxon.), F.R.C.P. Fourth Edition. London: Baillière, Tindall & Cox. 1906. Pages 434, 453 and 458.

^b Diseases of the Heart and Aorta. Dublin: Hodges & Smith. 1854 Page 603.

Cachexia in cancer is due solely to the growth preventing nutrition.

DR. LITTLE, from the chair, summed up the discussion, and mentioned that diagnostic weight may be attached to the effects of leeching in intra-thoracic tumour. Leeches always afford at least temporary relief in cases of aneurysm.