This article was downloaded by: [New York University]
On: 05 February 2015, At: 21:16
Publisher: Taylor & Francis
Informa Ltd Registered in England and Wales Registered Number:
1072954 Registered office: Mortimer House, 37-41 Mortimer
Street, London W1T 3JH, UK

Annals and Magazine of Natural History: Series 9

Publication details, including
instructions for authors and
subscription information:

[http://www.tandfonline.com/loi/
tnah15](http://www.tandfonline.com/loi/tnah15)

LXIX.—On a new genus of Coccidæ from the Indian region

E.E. Green F.Z.S. F.E.S.

Published online: 02 Sep 2009.

To cite this article: E.E. Green F.Z.S. F.E.S. (1921) LXIX.—On a new genus of Coccidæ from the Indian region , *Annals and Magazine of Natural History: Series 9*, 8:48, 639-644, DOI: [10.1080/00222932108632634](https://doi.org/10.1080/00222932108632634)

To link to this article: <http://dx.doi.org/10.1080/00222932108632634>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not

be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden.

Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

3. SCHUBOTZIA, Bouleng., 1914.

Scales ctenoid. A close-set series of enlarged teeth, with strongly incurved, flattened, rounded crowns, followed by 2 series of minute tricuspid teeth.

Lake Albert Edward.

Schubotzia eduardiana, Bouleng., 1914.

Schubotzia eduardiana, Bouleng. Cat. Afr. Fish. iii. p. 500, fig. 347.

Near *Haplochromis schubotzi*, differing especially in the dentition.

Total length 95 mm.

LXIX.—On a new Genus of Coccidæ from the Indian Region.
By E. E. GREEN, F.Z.S., F.E.S.

CRIBROLECANIUM, gen. nov. (subfamily *Lecaniinæ*).

Adult female with rudimentary limbs and antennæ. Spiracles communicating with the surface by means of a broad enclosed channel, the sides of which are studded with short glandular ducts. Dorsum with numerous, densely chitinous, perforated plates, arranged in more or less symmetrical series. Anal operculum surrounded by a densely chitinous area. Anal ring with ten or more setæ.

Nymph similar to adult, but with the limbs and antennæ still more rudimentary. Anal operculum not surrounded by a densely chitinous area.

Larva with fully developed limbs and antennæ. Dorsum with series of clustered pores in place of the cribriform plates. Male not observed in any stage.

Type, *formicarum*.

Cribrolecanium formicarum, sp. n. (Figs. 1 & 2.)

Fully matured adult female dark castaneous; subcircular, strongly convex, almost hemispherical; densely chitinous. At this stage of development the structural characters are obscured by the heavy chitinization, but the dorsum is seen to be studded with small translucent pores, interspersed with definite denser areas upon which the pores are more closely crowded. Other characters can be observed more clearly by

an examination of the early adult insect. Diameter of fully developed insect 4 to 5 mm.

Fig. 1.

Cribrolecanium formicarum, sp. n.

- a. Early adult female: opt. sect. $\times 30$.
- b. Antenna. $\times 450$.
- c. Posterior spiracle. $\times 130$.
- d. Pores surrounding spiracle. $\times 450$.
- e. Cribriform plate. $\times 220$.
- f. Anterior limb. $\times 450$.
- g. Part of margin. $\times 220$.
- h. Marginal spines. $\times 450$.

Early adult female (fig. 1, a) pale purplish brown; broadly ovate, narrower in front. Derm (after treatment with boiling

potash) soft and colourless, except on well-defined circumscribed areas. Antenna (fig. 1, *b*) rudimentary, without definite segmentation; some longish stout setæ on the apex and sides. Legs (fig. 1, *f*) minute and rudimentary, consisting of a broad basal segment, representing the combined coxa and femur, and a narrower tibio-tarsal segment surmounted by a well-developed claw; both unguis and tarsal

Fig. 2.

Cribrolecanium formicarum, sp. n.

- a.* Anterior leg of nymph. $\times 450$.
- b.* Posterior stigmatic area of nymph. $\times 130$.
- c.* Larva: opt. sect. $\times 50$.
- d.* Larva: group of pores. $\times 450$.
- e.* Larva: antenna. $\times 220$.
- f.* Larva: stigmatic spines and marginal setae. $\times 450$.

digitules are present, the former slightly expanded at their distal extremities. Buccal apparatus large and conspicuous. Spiracular channels closely studded with tubular ducts of the form shown in fig. 1, *d*; the channel usually curved and opening on the venter at a considerable distance within the margin of the body. Dorsum with a subdorsal and sublateral

series of densely chitinous, rounded, perforated plates (fig. 1, *e*), varying in size and form. Derm closely studded with minute pores, and with scattered, larger, thick-rimmed pores; also with scattered spiniform setæ. Valves of anal operculum together forming an oval, the base and outer edge of each valve describing an uninterrupted curve; the whole surrounded by a broad, densely chitinous zone, which is irregularly perforate around its inner margin. Anal ring with at least ten stout setæ. Stigmatic clefts shallow, without specialized stigmatic spines. Margin of body with a close fringe of strong spines (fig. 1, *g, h*), which are interrupted only at the stigmatic clefts.

Nymph very similar to the early adult insect, but distinguishable by the absence of a denser chitinous area surrounding the anal aperture. Antennæ and limbs still more rudimentary, the latter (fig. 2, *a*) being without definite claw or digitules. Cribriform plates as in the adult. Spiracular channels opening directly on to the margin at the stigmatic clefts (fig. 2, *b*). Marginal spines similar to and as large as those of the adult insect. No stigmatic spines. Anal ring with eight setæ.

Larva (fig. 2, *c*) with well-developed limbs and antennæ. The cribriform plates of the adult nymph are replaced by small groups of relatively large pores (fig. 2, *d*), interspersed with which are some isolated pores of a similar structure. Margin of body with simple, short, curved setæ. Stigmatic clefts with two stout spines (fig. 2, *f*), of which the anterior one is lanceolate and acuminate, while the posterior one is clavate and obtuse. Anal ring with six setæ.

Length 1 mm.

Peradeniya, Ceylon.

In hollow branches of *Stereospermum chelonoides*; associated with ants (*Cremastogaster* sp.). The branches had originally been tunnelled by some boring larva (probably of a Longicorn beetle), and had subsequently been occupied by the colony of ants.

Cribolecanium radicicola, sp. n. (Fig. 3.)

Fully mature adult female, pale fulvous (dried examples); rather broadly ovoid (fig. 3, *a*), moderately convex, the medio-dorsal area raised sharply into a broad rounded carina; derm soft and wrinkled, not densely chitinous. Antenna (fig. 3, *d*) small and rudimentary, but distinctly 4- or 5-jointed, the joints separated from each other by broad bands of softer

tissue, the narrow basal joint often incomplete. Legs (fig. 3, *b*) small and rudimentary, but relatively stout; coxa represented by an irregular narrow band, which is often

Fig. 3.

Cribrolecanium radicola, sp. n.

- a.* Adult female. $\times 30$.
- b.* Adult female: anterior leg. $\times 450$.
- c.* Adult female: cribriform plates. $\times 220$.
- d.* Adult female: antenna. $\times 450$.
- e.* Nymph: anterior leg. $\times 450$.

incomplete; femur, tibia, and tarsus all distinct, as broad as or broader than long; claw strongly developed, approximately equal in length to the tarsus; unguis and tarsal digitules slender, gradually dilated towards the extremity.

Spiracular channels broad, opening directly on to or close to the margin, closely studded with tubular ducts. Dorsum with irregularly disposed series of small, densely chitinous, cribriform plates (fig. 3, *e*) varying in size, form, and number of pores, but always much smaller and less conspicuous than those of *formicarum*, each plate with a narrow, sharply defined, paler outer border. Derm of dorsum with smaller and larger pores (the latter thick-rimmed) and with transverse series of spiniform setæ, which are larger and more numerous on the abdominal segments. Anal operculum surrounded by a densely chitinous zone, sprinkled with small pores and larger ovoid lacunæ. Anal ring with sixteen (or more) stout setæ. Margin of body without fringe of spines or setæ. Stigmatic clefts obscure, without stigmatic spines.

Length of average examples 2.5 mm.

Nymph very similar to the adult, but smaller and flatter, and without a denser chitinous area surrounding the anal aperture. Antenna 5-jointed, the basal joint in the form of a narrow band, second joint largest. Legs (fig. 3, *e*) reduced to conical points, with obscure traces of partial segmentation; with a minute apical claw. Cribriform plates as in adult, but often less strongly chitinized. Anal ring with ten setæ. Spiracular channels opening directly on to the margin. No stigmatic spines. No marginal spines or setæ.

Larva not observed.

Coimbatore, India.

On roots of *Cassia* sp. Coll. T. V. Ramakrishna (no. 204), 9. iii. 1921.

LXX.—*Some new or rare British Crustacea.*

By ROBERT GURNEY, M.A.

1. *Canthocamptus echinatus*, Mrazek.

In July 1919 a few specimens of a species of *Canthocamptus* resembling *C. echinatus* were taken at Flordon Common near Norwich, but I was unable at the time, with the scanty material available, to determine its identity with certainty, and I was unable to find the species again on a second visit to the spot. In 1920 the same form was found in considerable