that a wall of antibodies and leukocytes protects the secreting tissue. The essential value to the clinician lies in the statement as it was originally written. Whatever the human economy was planned for, it was not intended to digest pus. Even germs themselves break down and melt tissue by a process closely akin to peptonization, so that that is no bar to their action. The wonder is not that the effects of passing mucus-protected pus-gorms through the alimentary canal are bad and are generally ignored; yet, they are not so bad as one would imagine they would be if the specialist in gastrointestinal troubles will do two things; first, he will assert that mucus in the feces is not necessarily an evidence of lesion of the alimentary tract; second, he will not look into the nasopharynx for the source of the mucus.

Pyorrhea, alveolar abscess, and the like, seem to show their effects more as a toxemia than locally, that swallowed germs are inert is hardly credible. That swallowed germs are inert is hardly credible. They do not seem to be inert when taken out of the feces and cultivated. Of course, in this matter, overgrowth, unfavorable environment, and other factors are to be considered; also that the streptococcus or the staphylococcus are like pet dogs enjoying themselves in rummaging a garbage-can—these germs can become scavengers if they choose. In other words, these cocci are facultative and saprophytic and can live on dead fish quite as well as on the mucous membranes of a human throat. On the nonresistant putrid fish, they produce both an enzyme and a toxin. One can hardly believe that swallowing the toxin that is produced on the fish will cause the rapid death that it brings on, and then, subsequent to declaring such belief, expect any confidence in the idea that the same germs, working upon the contents of the alimentary canal and aided in attacks on its walls by an enzyme, should prove inert.

Providing the patient can swallow, tension, congestion, and all will disappear like magic from a throat if, in addition to local measures of a proper sort, the patient is given 30 grains of compound jalap powder with 1-3 grain of saccharin and 2-3 of a drop of oil of cinnamon; three such powders being given, spaced one hour apart. That the jalap must be up to standard would seem self-evident, were it not for many disappointments which arise from a contrary condition.

X-Rays in the Diagnosis and Treatment of Thyroid and Thymus Enlargement. A. Judson Quimby, and Will A. Quimby, New York, N. Y., Medical Record, January 6, 1917, p. 13.

The writers have found in over 250 cases referred for diagnosis and treatment that the presence of enlarged thymus is so frequent in exophthalmic goitre as to make it necessary to examine the chest for persistent thymus in all cases. It should also be looked for in other types of goitre, especially where the chest symptoms occur.

Enlarged thymus must be distinguished from intrathoracic thyroid enlargement and numerous pathological conditions within the chest. The characteristics of the thymus shadow is dwelt upon and the organ's susceptibility to the therapeutic effects of X-rays is emphasized. They believe that

the marked results obtained in X-ray treatment of exophthalmic goitre are in the cases possessing enlarged thymus, and that the greatest benefit is derived through the treatment of the thymus. The response to X-ray treatment is very prompt in some cases, relief occurring within a few hours, and is most marked in children.

A special warning is given concerning the administration of X-rays and the harmful effects resulting from over exposure. The importance of protecting the surrounding parts is emphasized, especially adjacent glandular structures, such as the salivary glands and breasts. X-ray treatment should only be administered to the younger subject with the greatest care. Some of the finer points in technique may be disregarded in treating the aged. Fifty-three cases were treated by the X-ray in conjunction with a surging high potential current, with evidence of increased benefit and less dangers of burn. They conclude that the relief symptoms justifies X-ray treatment in thyro-thymic enlargement.

The Interpretation of Certain Blood Pressure Ratios. Willard J. Stone, Toledo, Ohio, American Journal of Medical Sciences, Vol. CLIII, No. 2, February, 1917, p. 240.

The pressure-rate is the ratio existing between pulse-pressure and diastolic pressure. Under normal conditions the pulse-pressure is equal to 40-60 per cent. of the diastolic pressure. The diastolic pressure is the index of arterial tension and represents arterial resistance while the pulse pressure represents cardiac work. Two groups are recognized. In the first, vascular changes have produced rigid as well as contracted vessels. The diastolic pressure is consequently increased. The pulse-pressure is increased in normal or increased ratio if the heart is in compensation. This group includes the diffuse types of chronic nephritis. In 42 cases the average diastolic pressure was 132, the pulse-pressure 68, making the systolic 200. The pressure-rate was 51 per cent. Within three years' observation the mortality was 48 per cent. and was due to cerebral hemorrhage, thrombosis, or cerebral edema commonly called uremia. This group was designated as *cerebro-renal*. In the second group of hypertension, designated as *cardiac*, the vessels are rigid but not contracted. The diastolic pressure is therefore not high but the pulse-pressure is increased. This group is made up of valvular and aortic lesions. In 32 patients the average diastolic was 93, the pulse pressure 85, making the systolic 178. The pressure-rate was increased to 92 per cent. Symptoms of nephritis were absent in this group and a cardiac death occurred from dilatation or exhaustion in 25 per cent. within three years of observation. Cardiac dilation was accompanied by a fall in the pulse-pressure and a decreased pressure-rate. The prognostic point of value in the cerebro-renal group is the height of the diastolic pressure. In the cardiac group the prognosis seems to depend upon ability of the heart to sustain the increased pulse-pressure and pressure-rate necessary to adequately supply the periphery with blood.