THE COMPLEX OF IGNEOUS ROCKS AT OATLAND, SANTON,
ISLE OF MAN.

BY FRANK W. WHITE.

(Plates VIII.—XI).

(*Read 4th March, 1910.*)

INTRODUCTION.

The field relations of this complex of igneous rocks have been described by Mr. G. W. Lamplugh, F.R.S., in the Survey Memoir, "The Geology of the Isle of Man" (1903). Petrological notes by Prof. W. W. Watts, M.A., F.R.S., are also included in the volume.

A description of the acid type of rock at Oatland, as revealed by one thin section, had previously appeared under Mr. B. Hobson's name in Vol. XLVII. of the Q.J.G.S., and, later, in "Yn Liaor Manninagh" for 1892 (Vol. I., pt. 10).

Dr. F. H. Hatch briefly refers to the occurrence in his "Text Book of Petrology" (1909), pp. 285-6.

FIELD RELATIONSHIPS.

This igneous complex near Santon, in the S.E. part of the Isle of Man, is intrusive into the thinly-bedded Lonan Flags of the Manx Slate series. The Flags are visible on the East, West, and South-East in the immediate neighbourhood of the igneous mass, and have there been altered by the intrusion to a baked and splintery character. They are minutely spotted and banded, and, under the microscope, appear to consist of fine clastic grains of quartz in an aggregate of fine flakes of mica set in a groundmass of chlorite.

The structural axis of the Island runs N.E.-S.W. from Maughold Head east of Ramsey, through Foxdale to the Calf of

WHITE : OATLAND COMPLEX OF IGNEOUS ROCKS. 111

Man, and though the dip and strike of the bedding are extremely complicated, that of the main cleavage is fairly persistent and uniform over wide areas. The general dip of the cleavage in the "slates" is towards the S.E. on the eastern side of the structural axis of the island. For some considerable distance along the coast-line east of Oatland the cleavage dip is 60° , and at places about midway between Oatland and Foxdale on the W. it has the same direction and amount. At Oatland, however, the dip is only 30° to the S.E. on the eastern side of the intrusion, while on the western side the direction of the dip is reversed.

The actual junction of the igneous rock and the slates is nowhere visible at present (1909).

The intrusive rock itself is exposed in a quarry for road-metal in the centre of the mass, and in very low crags of small area on all sides. The boss is probably of nearly an oval shape where it reaches the surface, the outcrop showing a length of nearly 900 yards from N.E. to S.W., and a breadth of nearly 300 yards from S.E. to N.W. The longer axis is therefore parallel to the structural axis of the island.

Near the centre of the boss a small fault runs in nearly a N.W.-S.E. direction, but this was the only sign noticed of any mechanical disturbance of this character.

At the centre the rock appears to be a pale, mottled greenish-grey, close-textured "granitite," showing chlorite, feldspar and quartz, and an occasional speck or two of pyrites.

But on the outskirts of the boss, in the low crags, the rock is dark-green, with large crystals of augite or hornblende. Pyrites occurs more frequently. The rock resembles many others in the island, and may provisionally be termed a diabase.

In the deepest and most central part of the granitite, opened out recently, were found a few rounded knobs of the diabase as xenoliths, and, near the margin of the quarry, especially on the north-eastern side, were many sub-angular fragments of the same basic rock included in the granitite.

Some of the crags of basic rock are traversed by veins of the granitite, which thin out and become finer towards the circumference of the mass.

112 WHITE : OATLAND COMPLEX OF IGNEOUS ROCKS.

The basic and acid rocks probably have therefore a circumferential junction, though the main junction has not apparently been yet exposed. It is also clear that the basic rock was intruded prior to the more acid rock, though how long an interval elapsed between the intrusions remains to be determined.

Just south of the small fault in the granitite is a very narrow (about 2 ins.) and almost vertical dyke of greenish quartz associated with pads of a soft green, fibrous, asbestos-like mineral (actinolite ?) which closely resembles the "fibrous tourmaline" (Memoir), which occurs in similar veins in the Dhoon granite in the N.E. of the island. This quartz dyke appears in the quarry apparently striking to the N.E., and in a low crag of the basic rock cropping out in the adjoining field in this direction the vein was again found visible traversing both the diabase and the veins of granitite, and associated with the same fibrous mineral.

The granitite at the southern end of the quarry is very soft, decomposed and stained.

A rock which closely resembles a certain facies in the Oatland complex is visible about two miles to the N.N.E., near Ballabunt, and it is noteworthy that rocks with similar affinities occur as bosses in St. Michael's Island, and Langness, also in dykes running N.E. and S.W. along the coast to the east of Oatland.

AGE OF THE INTRUSION.

Unlike the Foxdale granite, and still more unlike the Dhoon granite, the rock at Oatland shows practically no sign of shear movement, and may be ascribed to a somewhat later date. This is confirmed by the effect the igneous mass has had in altering the dip of the cleavage in its vicinity.

But even the most basic modification at Oatland gives no evidence of the presence of either the monosilicate or bisilicate of magnesium, and, in the writer's knowledge, it resembles all the undoubtedly pre-Carboniferous intrusions of the island in this respect, but for the same reason is likewise in strong contrast to the olivine-bearing extrusions and intrusions of Carboniferous or Tertiary age. This would still leave the age of the intrusion of the granitite an open question as evidence seems to point to the lapse of a considerable interval between the intrusion of the more basic and more acid types in the complex.

WHITE : OATLAND COMPLEX OF IGNEOUS ROCKS. 113

For a similar reason, and in view of its close identity with the more basic facies at Oatland the age of the smaller intrusion at Ballabunt is considered to be contemporaneous with the earliest intrusion at Oatland.

DETAILED DESCRIPTION OF THE ROCKS.

A series of thin rock-sections made from specimens taken in various directions revealed the fact that a gradation in character is fairly constant from the most acid to the most basic type. For convenience of description, however, the complex may be resolved into three types of rock, acid, intermediate and basic. Among other considerations, which will be apparent later, we hold the acid facies to be distinguished by the presence of orthoclase, and the basic by the absence of original quartz. The most acid type is found in the centre of the boss, and there is a fairly constant increase of basic characteristics as one proceeds to the margin, although it is not possible to fully test this in every direction, and the megascopic examination of the rocks in the field indicates an increasingly rapid assumption of basic character towards the periphery.

1. ACID TYPE.

(Plate VIII).

The rocks in this section are mostly adamellites, monzonites and granodiorites. They are, however, all characterised by the presence of a considerable amount of quartz, with idiomorphic plagioclase, varying quantity of orthoclase, chlorite pseudomorphs after biotite and a little hornblende. In each case the characters of the minerals will be described in the order in which they appear in the specimens arranged in a series of decreasing acidity.

Quartz.—This has been the last mineral to crystallise. It resembles that of the Foxdale granite in having the granitic habit, while that of Dhoon and the Crosby microgranite (an elvan of the Foxdale mass) is quite granular. It should be remarked, however, that one section shows it as granitic and locally granular, another as mostly granular, and a third as having some undulose extinction. This is probably due to a merely local movement at the time of consolidation. The quartz has the habit of completely surrounding the other minerals and occasionally encloses

114 WHITE : OATLAND COMPLEX OF IGNEOUS ROCKS.

sub-angular fragments of its own species. Fluid pores arranged lineally or in strings increase in number and size from very few to very many. Some contain gas cavities. Staining with iron-ore is shown between some of the crystal faces. Inclusions become more and more numerous, comprising apatite, small flakes of muscovite probably derived from the orthoclase, zircon, epidote (sometimes as disconnected granules in parallel strings, perhaps derived from neighbouring hornblende), zoisite, some large crystals of iron-ore, and an unidentified mineral which appears in two specimens. This latter mineral is of irregular outline and colourless. It has no fluid pores; against it the quartz with its crowd of fluid cavities is everywhere arrested, but it contains one or two negative octahedral crystals, probably of fluid. It is chatoyant by reflected light, is isotropic, and has a refractive index about equal to that of quartz, greater than felspar, and less than biotite.

Measuring off the intercepts of the minerals by Rosiwal's method in the various sections, and calculating their gravimetric proportion in the rock, it was found that quartz decreases from 37 per cent. to 18.4 per cent. in this acid facies.

Orthoclase is always present but in varying quantity. Its outline is usually irregular as in other Manx granites.

Fig. 1.

Fig. 2.

FIG. 1.

Oatland Complex. Acid Facies. × 20.

Much granitic quartz, with a few inclusions of apatite: some plagioclase and corroded orthoclase, also a granular aggregate of epidote, probably after biotite.

FIG. 2.

Oatland Complex. Junction Specimen. × 20.

The portion of the basic xenolith shows an allotriomorphic aggregate of secondary minerals, the fibrous hornblende having a tendency towards the actinolitic habit.

Bordering the xenolith is much clear quartz and a large crystal of decomposed plagioclase.

(Photographed by W. L. Carter.)

WHITE : OATLAND COMPLEX OF IGNEOUS ROCKS. 115

Carlsbad twinning is common, and quartz has sometimes been forced along the twinning plane (Fig. 2). Some of the larger crystals were measured, the dimensions being 2.75×2.35 mm., 2.31×1.6 mm., 3.78×1.45 mm., and 2.5×1.25 mm. In the granodiorite the average size would be about $1.4 \times .65$ mm. There are inclusions of apatite, usually surrounded by iron-ores, grains of magnetite, and a few patches of chlorite. The orthoclase becomes increasingly turbid by decomposition, the products of which, in addition to amorphous matter, are flakes of muscovite which lie nearly normal to each other in some sections (Figs. 1 and 2), and parallel to the sides of the crystal in others. The included iron-ores are sometimes altering to sphene. Clear bipyramidal quartz is an occasional decomposition product. The turbid orthoclase often has a narrow rim of clear felspar substance. When orthoclase and plagioclase are in contact it is evident that the former has always crystallised prior to the latter.

The gravimetric proportion of orthoclase in this facies decreases from 28 per cent. to 6.1 per cent. of the whole rock.

Plagioclase usually shows more regular outlines than orthoclase, and has the habit of projecting into the quartz. The pinacoids show especially good faces, but some faces are apparently corroded by quartz. There are some larger crystals measuring about 2.0×1.15 mm., but the average sizes vary from about $.72 \times .4$ mm. to $1.1 \times .9$ mm. Twinning according to the albite law is common, but there is a gradual assumption of pericline twinning as well. The width of the lamellae in the former type becomes slightly increased. The maximum extinction angles, measured from the twinning-planes show a gradual increase from about 11° to about 16° . The felspar is probably an oligoclase-andesine, but in all the specimens there are some crystals of apparently a medium oligoclase. The individuals are turbid with decomposition products, but usually have borders of fresh felspar substance, sometimes at the ends, sometimes—usually when quartz is adjacent, at the sides as well, and sometimes at the ends and in interior patches. It is considered that the fresh felspar is more acid when added to the ends than that on the sides, and still more than that in interior patches. The new growth of

116 WHITE : OATLAND COMPLEX OF IGNEOUS ROCKS.

felspar round the borders exactly resembles that round the plagioclase in the Foxdale and Dhoon granites.

The decomposition product in addition to amorphous matter is mostly white mica, but much epidote occurs in places, often in veins which completely traverse some of the crystals. The flakes of white mica often project from the turbid patches into the clear felspar, and are sometimes completely surrounded by it.

There are inclusions of apatite usually surrounded by iron ore in small quantity, and iron-ore in octahedral crystals. One large enclosure is an aggregate of dusty-looking greenish-brown sphene with ilmenite, calcite and chlorite. Not only is plagioclase moulded on the orthoclase, but occasionally the former encloses in its centre small but well shaped crystals of the latter with Carlsbad twinning.

The gravimetric proportion of plagioclase in these specimens is variable, showing, in the sections examined, 39.2 per cent., 26 per cent., 36.5 per cent., 25.4 per cent., and 41.2 per cent. respectively of the total mineral content.

Biotite in greenish pseudomorphs of chlorite, with slight pleochroism, low birefringence and straight extinction, occurs with hexagonal basal sections, and longitudinal sections the ends of which are both ragged and granular. Sometimes the colour is greenish-brown, and the pleochroism becomes more marked, showing green, greenish-yellow and very pale yellow. The edges are semi-opaque with what at first sight appears to be magnetite dust, but under a higher power this seems to be due to minute granules of epidote which mutually interfere. The presence of these microscopic grains in the interior of the mineral probably gives rise to the brownish-olive colour that is occasionally presented by sections under crossed nicols. Again, crystals are often bleached, and what seem to be thorns of quartz often appear between the lamellae.

Next to chlorite, epidote is the principal product of altera-

tion. It is in yellowish grains with pleochroism, β brownish-green, α and γ green and pale chrome. Other products are a dark-brown sphene and a blood-red haematite.

Inclusions are abundant, comprising rather large prisms of apatite and zircons up to .07 mm. in length, surrounded by pleochroic halos.

The gravimetric proportion of the chlorite pseudomorphs after biotite is very variable in the different sections in this series.

.

There is another mineral present in variable quantities, "unidentified" in the Memoir, which occurs in such a fashion as to suggest its derivation from a biotite which was originally more ferriferous and titaniferous than the biotite now showing a merely chloritic alteration. It is rarely present in the most acid specimens, but becomes an important constituent in the granodioritic type.

I am indebted to Mr. A. Harker, F.R.S., for the suggestion, based on my written description, that the mineral is merely a biotite which is undergoing a change due to oxidation following chloritization.

In some sections it is frequently difficult to distinguish this mineral from the ordinary biotite. Its outline is sometimes hexagonal. The extinction is nearly straight. In longitudinal sections its ends are frequently ragged, and usually show the same pseudo "resorption border" as the biotite. Its refractive index is high, but apparently less than the chloritized biotite. In plain light it is of a turbid brownish colour and is almost opaque between crossed nicols. Its pleochroism is slight, but oblique sections show light brown and brownish-olive. Under a high power the lines of apparently opaque iron-ore are often found to be really semi-opaque brownish particles, almost ultramicroscopic in size. But in other sections they are undoubtedly iron ores, mostly leucoxene after ilmenite.

118 WHITE : OATLAND COMPLEX OF IGNEOUS ROCKS.

The appearance of some individuals is here illustrated.

Fig. 3.

Fig. 4.

Fig. 5.

Fig. 6.

In Fig. 3 the mineral is partially enclosed by green biotite, and its end is separated from the biotite by granular epidote with pleochroism greenish-yellow and yellowish-green.

In some cases the green biotite encloses much similar epidote, and suggests that one mineral is wholly derived from the other. In other sections layers of green biotite alternate with brown layers of this mineral, the whole showing pleochroism in shades of green. The mineral is also intergrown with a bluish-green

hornblende, but in such a fashion as to suggest that the latter was rather later in crystallising. However in some cases an individual of fibrous amphibole with an extinction of 15° is enclosed. Apatite in prismatic forms is the other usual mineral that is enclosed.

In the process of decomposition, besides varying amounts of epidote, there has usually been a fairly heavy separation of iron-ore, mostly titano-magnetite.

The additional fact that the biotite in the acid facies has commenced to form at about the same relative period as the more oxidised variety, which occurs in the intermediate and basic facies of the complex, affords confirmation of the micaceous character of this mineral if not of its close identity with the less altered biotite.

Hornblende is present in increasing quantity from the centre to the margin of the boss, but in the normally acid facies it does not amount to more than 7 per cent. of the total mineral content and from the most acid specimens in the facies it is nearly absent. Here it is greenish in pseudomorphs of chlorite and is usually associated with the biotite. In one case the alteration product has the structure of helminth. The crystals are sub-idiomorphic with pleochroism light-green—darker green. Its angle of extinction appears to be only about 5° degrees. It occasionally encloses many needles of a mineral which also extinguishes at 5° degrees, and in one individual these lie along three planes, thus—

Fig. 7.

120 WHITE : OATLAND COMPLEX OF IGNEOUS ROCKS.

Alteration products, besides iron-ores in smaller quantity, and the usual chlorite, are secondary dark brown sphene, developing in patches, and grains of epidote. As an inclusion of original mineral, apatite is usually found.

The gravimetric proportion of hornblende in the dioritic end of the acid facies increases from 1.4 per cent. to 7 per cent. of the mass.

THE EFFECT OF THE INCLUDED BASIC XENOLITHS.

The above concludes the description of sections of the acid rocks in their normal conditions, but two sections of rock were examined showing the junction of basic xenoliths, caught up in the acid magma, with the granitic rock.

The first was taken from the deepest and most central part of the boss that was accessible and the second from the north-eastern side of the quarry. Macroscopically the former showed a rounded, the latter, a sub-angular outline.

The former (Plate VIII., Fig. 1) was found to consist of an allotriomorphic aggregate of minute chlorite, muscovite, calcite, epidote, quartz and iron-ores. Of the original minerals, biotite preserved the best forms with epidote enclosed. There were also traces of medium-sized (about .8 mm. by .8 mm.) augites now as chlorite, epidote and calcite, also some twinned hornblende and a patch of prismatic fresh-looking actinolite having the outline of a cross-section of hornblende. The quartz was in very fine threads which extinguished simultaneously over very small areas and were crowded with minute inclusions of the other minerals.

In the other section, confining one's attention to the apparent effect of the xenolith on the character of the minerals in the acid portion of the slide, it was observed that the quartz contained some unusually large crystals of iron-ore, that the amount of plagioclase present was abnormally large while one crystal of augite occurred—probably a xenocryst, with a somewhat irregular shape. Its average breadth was 1.2 mm. It was mostly uralitized, but chlorite was also present in quantity, and practically no trace of the original cleavage remained. A broad border of hornblende surrounded the kernel.

WHITE : OATLAND COMPLEX OF IGNEOUS ROCKS. 121

The hornblende in this part of the section closely resembled that in the intermediate and basic types in presenting two varieties, one with a pleochroism greenish-blue to pale yellowish-pink and extinguishing usually at 22 degrees and 23 degrees; the other with a pleochroism greenish-brown, brownish-green and pale straw. The latter is about three times as abundant as the former. The greenish-blue crystals are often sub-idiomorphic and contiguous to biotite. Actinolitic needles with extinction up to 9 degrees sometimes radiated from the hornblende into the biotite. The brownish-green variety in its larger patches contained some magnetite dust and suggested that it had been derived from augite or the brown hornblende usually included in it.

ORDER OF CRYSTALLISATION IN THE ACID TYPE.

The order in which the minerals commenced to form in the central or acid portion of the boss is the same in all the sections examined and is as follows :—

1. Apatite and zircon.
2. Iron ores.
3. Biotite and hornblende.
4. Orthoclase.
5. Plagioclase.
6. A little hornblende (in the less acid part only).
7. Quartz.

SUGGESTED AFFINITIES.

The chemical composition of most of the specimens, based upon the quantitative mineral composition, suggests that the most acid type resembles the composition of the "potash-granite" from Lamorne, Cornwall, and closely resembles that of the Skiddaw granite and the granite of the Mourne Mountains, while it is not far removed from the potash-soda felsite from Waterford, the Oatland rock being deficient in alumina but having an excess of magnesia.

The granodioritic type mineralogically resembles the quartz diorite from Arran, described by Zirkel, and by Teall, "Brit. Petr.," pp. 263-4.

2. BASIC TYPE.

(Plates IX., X.)

It will assist in the description of the intermediate type of rock, and save repetition, if the mineralogical features of the most basic type are next considered. The rocks in this section grade from hornblende-augite-diorites to a pyroxenite. They are all characterised by the absence of original quartz except in those parts in which there has been an intrusion of this mineral as veins or threads from the more acid magma. In such cases it has had a noteworthy effect in altering the condition of the invaded rock at the planes of junction.

Felspar is practically undistinguishable in the normal rock of this type, but in its place appears an allotriomorphic aggregate of mineral substance "undetermined" in the Memoir. The aggregate is of a dusty greenish-brown or a faint turbid brown colour. Under crossed nicols, owing to its intimate mixture with chloritised hornblende, it sometimes shows an indigo tint. The mineral has a high refractive index which, however, is less than that of either of the hornblendes and the augite. Where the granules were larger they were determined as epidote, zoisite and quartz. In a few cases where the aggregates are completely enclosed by brown hornblende they show good outlines and suggest that they are the products of an extremely decomposed plagioclase felspar. In confirmation of this, the quantity of the minerals present in the aggregates, estimated as plagioclase, was determined, and it was found that the sequence in the three facies then became intelligible. Moreover, the order of crystallisation relatively synchronised with the order as more certainly observed in the intermediate and acid types. Its original feldspathic character was placed beyond doubt by the phenomena observed in a section which showed this rock invaded by a single small vein of quartz. Near this quartz the dusty mineral was found associated with fresh felspar substance. Still nearer the quartz the dusty mineral disappeared and perfectly fresh feldspars were alone found. The extinction angles of these clear feldspars were measured and varied from 18 degrees to 20 degrees. The original felspar was, therefore, probably an andesine.

WHITE : OATLAND COMPLEX OF IGNEOUS ROCKS. 123

The gravimetric proportion of the original plagioclase in the rock was found to decrease, as the basic type became richer in pyroscene towards the margin of the mass, from 36.4 per cent. to 23.6 per cent., and finally only 11.1 per cent.

Biotite in greenish pseudomorphs of chlorite is not found in this type normally. Only in the section showing the fresh quartz vein does a little appear, giving straight extinction.

The more oxidised variety occurs in small but variable quantity. However, large patches of iron-ore have in many cases the appearance of pseudomorphs of this mineral and where the original character of the mineral is more recognisable it has often set free an abnormal quantity of iron-ore—a titaniferous magnetite or ilmenite with lamellae intersecting at 60 degrees. One crystal appears thus—

Fig 8.

The outlines of the less altered mineral are often irregular. It lies in the borders of the augite and in the hornblende and several sections measure about .35 mm. by .15 mm.

Calculating the quantities of pseudomorphs as well as the less altered mineral it was found that the proportion of this richly feriferous biotite showed a gradual increase as the type became more basic from 2.1 per cent. to 8.2 per cent.

Hornblende occurs in two varieties, one pale green and almost colourless in ordinary light, the other giving shades of brown, greenish-brown, golden-brown to a deep yellowish ochre.

124 WHITE : OATLAND COMPLEX OF IGNEOUS ROCKS.

The pale green variety, now mostly chloritic, has rather a fibrous texture and usually encloses a central patch of a tangle of actinolitic blades giving extinction angles of 13 degrees, 16 degrees and 20 degrees. In the pyroxenite, however, it is only just beginning to take on the actinolitic habit. Pleochroism is very faint in shades of very pale green. The chlorite of which the mineral consists, gives, under crossed nicols, a pale indigo to brownish-green colour. The alteration product, besides chlorite, is epidote and zoisite, the former sometimes occurring in parallel strings of disconnected grains. This variety forms about 4.5 per cent. of the total mineral-content.

The brown or ochreous variety often encloses the former and cleavage traces are frequently continuous from one to the other. Its pleochroism varies from light-brown—darker brown to brownish-ochre—pale straw. It is allotriomorphic and occurs not only in isolated patches but is enclosed in streaks and grains by the augite. These, and the zones of brown hornblende surrounding the augite crystals are optically continuous with the latter. This brown variety is compact and its refractive index is higher than the chloritic. Where the former anastomoses into the latter its pleochroism changes from brownish-ochre gradually to bluish-green and green tints. The mineral is frequently in large patches, enclosing idiomorphic aggregates of felspar substance, and extinguishes simultaneously over wide areas. Some large apatites are also enclosed. This variety forms a proportion of the total mass diminishing from 18.2 per cent. to 7.1 per cent.

Near the quartz vein in the above-mentioned section the pale hornblende shows undulose extinction and the chloritic substance of which it consists, instead of showing the usual indigo in polarised light, gives an olive or brownish-olive colour. Close to the quartz the hornblende is clear, with very distinct cleavage and the iron-ore in it is streaky as though partially segregated.

Augite is the mineral most abundantly present in this type. Its form is at first somewhat irregular, approaching the quadrate in section, then outlines showing well-developed clino-pinacoidal faces appear. The dimensions of some crystals almost reach 3 mm. by 3 mm., but the average size is about 2.2 mm. by 2.0 mm.

WHITE : OATLAND COMPLEX OF IGNEOUS ROCKS. 125

Smaller grains are enclosed poecilitically in the hornblende. Twinning is fairly common, but there are very few traces of cleavage left. The mineral is mostly uralitic, and often now chloritic, but frequently it appears as colourless malacolite, with refractive index greater than hornblende, extinction angles of about 44 degrees, and strong birefringence. In one section there is an approach to the diabase type with a few crystals of felspar in a granular state, but with fair outlines enclosed in the marginal zones around the augite. These enclosures range from .1 mm. by .1 mm. to .56 mm. by .26 mm. in section. The augite has therefore continued to grow after the now oxidised biotite and the plagioclase have commenced to form. Magnetite dust is also present in a zonary arrangement near the margins.

The proportion of augite present in this type gradually increases from 38.3 per cent. to 66.8 per cent. of the total content.

In the section showing the intruded thread of quartz, augite is associated with the clear plagioclase crystals. It is here perfectly clear malacolite with high refraction and birefringence. One crystal extinguishes at 47 degrees, and in another the cleavage lines make an angle of 85 degrees.

Iron Ores are present in variable quantity, specially associated with the hornblende and augite. In the former the larger idiomorphic crystals appear to be magnetite and the smaller crystals in the augite are ilmenite now changing to leucoxene.

ORDER OF CRYSTALLISATION IN THE BASIC TYPE.

The allotriomorphic character of so many of the minerals indicates that several of them probably ceased to crystallise almost simultaneously but by a comparison of the sections the following is held to be the approximately correct order in which they commenced to form.

1. Ilmenite, magnetite and apatite.
2. Augite and some brown hornblende.
3. { Felspar.
Biotite.
Brown and greenish hornblende.

126 WHITE : OATLAND COMPLEX OF IGNEOUS ROCKS.

SUGGESTED AFFINITIES.

The augite-hornblende diorite, besides resembling many of the "greenstones" found in the Isle of Man, is, from its estimated chemical composition, comparable with some of the augite diorites in Minnesota and the hornblende-augite diorite from Inchnadamf. It probably approaches the composition of a shonkinite chemically. The most basic type would closely resemble the composition of a pyroxenite from Brandberget, Gran, Norway.

3. INTERMEDIATE TYPE.

This type has variable characteristics, but on the whole combines those features which distinguish both the acid and basic types. Specimens arranged in series of decreasing proportions of quartz do not show a corresponding increase of augite and vice versâ, and no specimens have yet been found with both these minerals absent. This variable character suggests that the intermediate facies is a "mixture rock."

Quartz, when it occurs, is very rarely fractured, and is crowded with fluid cavities. There is also an inclusion of an undetermined mineral of prismatic form with blunt pyramidal ends, cross fracture, refractive index apparently less than that of quartz, weak birefringence and doubtful pleochroism in faint blue and yellow. In the basic xenolith previously mentioned the quartz seems to have separated previously coterminous and interlocking crystals, to have torn off minute ragged pieces of the fibrous hornblende and to have partially re-fused the felspar substance towards the boundaries of the fragment.

Felspar. Where quartz is in small quantity or absent, the feldspathic mineral is practically allotriomorphic. Elsewhere the forms are usually prismatic, some are sub-idiomorphic and some have still more irregular outlines. The average size is about .6 mm. by .3 mm. It is extremely decomposed so that the genus is quite undeterminable, but the alteration products are quartz, calcite, an epidote or zoisite and amorphous matter. No white mica could be found. Inclusions are very rare.

FIG. 1.

Oatland Complex. Basic facies. × 20.

The allotropic compact brown hornblende encloses felspar crystals and large patches of colourless fibrous hornblende. The opaque mineral is titanomagnetite pseudomorphous after biotite.

FIG. 2.

Oatland Complex. Basic facies. × 20.

The large individuals of colourless fibrous hornblende are now chloritised with separation of much calcite and epidote. A few blades of actinolite are just discernible near the centre of the section. Cloudy, allotropic felspathic matter completes the mineral content.

(Photographed by W. L. Carter.)

WHITE : OATLAND COMPLEX OF IGNEOUS ROCKS. 127

Biotite is sparingly present to absent in its pale green variety. One less-altered individual of the other variety appears with rather strong pleochroism in two shades of brown, but it is also turbid in places like the more oxidised crystals. The latter are fairly abundant with pleochroism of brown and olive-brown. In addition to the basal cleavage of a mica there is occasionally a poor cleavage not unlike the prismatic cleavage of an amphibole. The following (Figs. 9 to 12) will give an idea of its outline and that of the associated iron ores :—

Fig. 9.

Fig. 10.

Fig. 11

Fig. 12.

128 WHITE : OATLAND COMPLEX OF IGNEOUS ROCKS.

The appearance is similar to that of the same species in the more basic acid types of rock. Individuals are sometimes pierced by thorns of quartz like the green biotite in the acid rock. The broad cleavage bands would perhaps evidence a scaly character in the altered mineral like the biotite in the plagioclase-augite rock at Tolcarn, near Penzance, where it has been altered by the granite.

The included iron-ores are not quite opaque at their extreme margins, which show a muddy greenish-brown colour. Where crystals of the biotite have been bleached to a greenish-brown colour the iron-ores are not nearly so plentiful.

Hornblende is found in two varieties, as in the basic rock, but the brown variety is usually tinged with greenish shades, and it occurs in varying quantity with the pale green variety, though the proportion of total amphibole is remarkably constant in all the specimens examined, remaining at 21 per cent. to 23 per cent. The darker variety, with pleochroism = dark green—light green—brownish-straw, often lies embedded in the quartz while the hornblende surrounding the augites is lighter in colour, with pleochroism = bluish-green—brownish-yellow and pale straw.

In the basic xenolith above mentioned the hornblende often occurs as large individuals, sometimes twinned. The extinction angle is about 15 degrees.

Augite has characters like those in the basic rock. One chloritic pseudomorph appears to have a distinct "herring-bone" structure. The outlines are usually regular and hexagonal, and the average dimension is 2.5 mm. by 2.5 mm. There are some colourless patches of unaltered malacolite. The products of alteration are usually uralite, chlorite and calcite, the latter along cleavage planes, and a little iron-ore in minute grains.

In the basic xenolith, particular crystals of augite show interesting features. One individual 2.6 mm. by 1.82 mm. has within the outer zone of chlorite a patch of uralite about 1.45 mm. by 1.45 mm., surrounded by a narrow zone of magnetite dust and hornblende with a pleochroism = brownish-green—light-green-straw. In the heart of the crystal is a patch of unaltered

Oatland Complex. Extreme Basic Facies. × 20.

The uralic augite crystal is seen enclosing small patches of pale fibrous hornblende. Compact hornblende almost surrounds the augite. Dusty felspathic matter occasionally with the outlines of plagioclase lies embedded in the pale green chloritised hornblende. An individual of oxidised biotite is by the circumference remote from the augite.

(Photographed by W. L. Carter.)

WHITE : OATLAND COMPLEX OF IGNEOUS ROCKS. 129

malacolite about .5 mm. by .5 mm., with strong D. Refraction. Similar phenomena belong to another crystal measuring 2.4 mm. by 2.3 mm. A still larger crystal, 2.9 mm. by 2.5 mm., shows multiple twinning like the augites in the rock at Poortown, but more hornblende appears to have been originally included and more iron-ores have been liberated. Quartz has penetrated the crystal in several directions.

A few general observations may here be made respecting this sub-angular basic xenolith. Near the margins of the fragment the minerals are arranged—like the whole of the xenolith found in the heart of the boss—as a confused aggregate of minute crystals crowded with amphibole, while the feldspars of the surrounding granodiorite lie tangentially to the outline of the knob. The acid portion of the slide shows an abnormal quantity of the darker oxidised biotite, and, to a less degree, of plagioclase. It appears as though the matrix of amphibole had been dissolved from the surface of the xenolith leaving the biotite and feldspar and a little augite free in the solute. Near the included fragments the hornblende has recrystallised in good idiomorphic forms unlike its usual habit elsewhere in the boss.

ORDER OF CRYSTALLISATION IN THE INTERMEDIATE TYPE.

Similar considerations affect the determination of this as are noted in connection with the basic type.

APPARENT ORDER.

1. Apatite, ilmenite and magnetite.
2. Augite, and, in places, a little hornblende.
3. Feldspar { 3. Br. Biotite } in the more acid
4. Br. Biotite or { 4. Feldspar. } facies of the type.
5. Compact hornblende.
6. Fibrous hornblende.
7. Quartz, when present.

SUGGESTED AFFINITIES.

The intermediate type may be described as an augite-hornblende-biotite-diorite. Mineralogically it resembles the rock from Seathwaite How (Teall, Brit. Petr., p. 276). The calculated

130 WHITE : OATLAND COMPLEX OF IGNEOUS ROCKS.

chemical composition closely agrees with that of the diorite from Netherley, Strathspey, also with the diorite from Benn au Thurian, except that in the latter the ratio between potash and soda is inverse.

THE ROCK AT BALLABUNT.

Two sections from this small intrusion were examined and the results showed that the rock presents close correspondences with that at Oatland, which occurs only two miles to the S.S.W., in a direction parallel to the structural axis of the island.

Quartz has the granitic habit, as at Oatland, and is crowded with fluid cavities.

Felspars have also the habit of those in the more acid rock at Oatland. They are extremely decomposed, apparently to white mica, and cases of twinning can rarely be detected. A few forms gave extinction angles of about 13 degrees.

Biotite of green colour in pseudomorphs of chlorite enclose octahedra and irregular grains of titanomagnetite together with a few veins of epidote.

Augite occurs in some quantity, now uralitised and of a very pale green colour. It is almost always twinned. Some subidiomorphic green hornblende and tangles of actinolite, the blades of which often cross at angles of 84 degrees, occur in the interior of the crystals. The refraction of the actinolite is less than that of the surrounding hornblende. Some apatite is included.

Hornblende of a dark green colour occurs on the borders of the augite and usually, but not always, optically continuous with it. A little compact green hornblende also occurs apparently not connected with the augite but frequently moulded on the biotite. The crystals are usually more compact than those at Oatland.

Iron Ores. Ilmenite is the most abundant but scales of haematite occur in the augite and the green hornblende.

GRAPHIC REPRESENTATION OF THE MINERAL CONTENT OF THE ROCKS.

In this diagram the length of the intercept of any mineral on the vertical line will give the percentage of that mineral in a given rock section. Twelve sections were measured, numbers 1 to 6 and 9 to 12 being from the Oatland complex, and numbers 7 and 8 from the Ballabunt Rock. In each case also the order in which the minerals are placed, reading from top to bottom, will give the order in which they successively crystallised in bulk.

From Vachas, Chas. U.S. Nat. Mus. Vol. VII, Plate VI

WHITE : OATLAND COMPLEX OF IGNEOUS ROCKS. 131

The order of crystallisation of the minerals was the same in both slides, viz. :—

1. Apatite and iron ores.
2. Augite.
3. Felspar.
4. Biotite.
5. Hornblende.
6. Quartz.

The gravimetric mineralogical composition of the two sections proved to be

			1.		2.
Quartz	..	..	10.03	..	10.09
Felspar	..	..	37.95	..	39.49
Augite	..	..	29.99	..	32.57
Hornblende		..	14.70	..	12.34
Biotite	..	..	4.53	..	2.88
Iron Ores	..	..	2.80	..	2.63

SUGGESTED HISTORY OF THE INTRUSION.

Following the great E. and W. earth pressure, which caused the complex folding in the island, came the intrusion of an igneous rock magma, probably already highly differentiated, as a sill of basic rock.

Afterwards, and the petrological evidence on the whole points to the lapse of a considerable period, came a second intrusion from the same magma reservoir, probably laccolitic in habit, which lifted the overlying rocks and by intimate mixture with the already consolidated basic rock formed at the plane of junction a hybrid or "mixture" rock while the acid magma penetrated the surrounding crust of basic rock in innumerable tongues, veins and *Fasern*.

The last effort of the igneous intrusion to reach the surface resulted in the injection of the remaining super-heated silica-magma as a quartz dyke (similar to those at Foxdale, described by Lomas) which dissolved some of the last crystallised and fibrous amphibole and altered the basic rock at its conterminous parts into a fresh-looking augite-diorite.