A SECOND BRACHYDACTYLOUS FAMILY.

By H. DRINKWATER, M.D., F.R.S. (Edin.), F.L.S.

I HAVE already published accounts of three families showing an inherited symmetrical shortness of the fingers and toes in individuals who are also below the average in stature. In one—the “Brachydactylous Family”¹—the fingers are reduced to about half the normal length: in the other two families they are intermediate in length between these very short ones and the fingers of average normal individuals, a condition which I have designated by the term “Minor-Brachydactyly”².

In both types, the shortening was shown to be due chiefly to an abortive condition of the middle phalanx. The main features are represented in the outline illustrations in Fig. 1, where *A* shows the bones of the middle finger of a normal adult, *C* the Brachydactylous condition, and *B* the Minor-Brachydactylous.

In *C* the middle phalanx (2) is seen to have become ankylosed to the terminal phalanx (3).

In the Summer of 1913, my friend Dr J. D. Lloyd informed me that he knew of some people resident in his neighbourhood whose hands closely resembled those of one or other of the above-mentioned families; and he not only afforded me an opportunity of examining some of them at his own surgery, but very generously consented to allow me to carry out whatever investigations I might be disposed to undertake with regard to them.

Examination at once made it evident that the abnormalities in these new cases are identical with those described in my paper on “Brachydactyly”; there is, namely, the maximum amount of shortening of the digits, and the same characteristic shortness of stature. The essential peculiarities are so exactly similar that one cannot resist the

¹ “An account of a Brachydactylous Family,” *Proc. Roy. Soc. Edin.* Vol. xxviii. Part 1.

² “Account of a Family showing Minor-Brachydactyly,” *Journal of Genetics*, February, 1912. “Minor-Brachydactyly” No. 2, *Journal of Genetics*, February, 1914.

conviction that the two families must have descended from the same original stock.

The connection between them unfortunately cannot now be established, for there is not a single surname common to the two families. This difference in names may be accounted for by the

Fig. 1. Normal and Brachydactylous Phalanges. (Natural size.)

A. Normal. B. Minor-Brachydactylous. C. Brachydactylous.

fact that the inheritance in the second family has been almost exclusively through the female line, so that the surname has changed (through marriage) at each generation.

Both families reside in the same part of the country: a fact which would seem to give some support to the theory of their common origin.

Dr W. C. Farabee was the first to describe the condition of Brachydactyly in a family that he studied in 1903¹ in North America. Is

¹ Published in March 1905 by the Peabody Institute, Harvard, U.S.A.

there any blood relationship between this American family and either of the English families? The cases observed by Farabee were apparently identical as regards the anatomical and hereditary features with those described in my first paper in August 1907. I was not aware of Farabee's work until my own investigations were nearly completed, and there was not time to decide the question of relationship before presenting my paper to the Royal Society of Edinburgh. His reply to my enquiries did not come for several months afterwards, for he happened to be in some out-of-the-way part of Peru engaged on a scientific expedition. However, it was found that no surname was common to the two families; nor was it possible to make my chart fit in with his: so that, assuming that there had been a common ancestor of the Brachydactylous type (and this hardly admits of doubt), he or she must have lived prior to the earliest member now traceable in either of the two families. A study of Farabee's account made me think it likely that his family had descended from "an abnormal member of the English family, four generations back, who had migrated to America, but of whom no tidings have since been received by his relatives in this country," but as this man's name does not occur in Farabee's family, he was obviously not the connecting link. So far, therefore, I have not been able to prove any blood relationship as existing either between my two families or between the first of these and Farabee's.

There still remains the question whether the second family (about to be described in the present communication) is related to Farabee's. Farabee informed me that the name amongst his people which most nearly approached any name in the English family was *Hyde*. Now the man marked 2 in the chart on page 327 was named *Benjamin Hyde*. In November of 1913 when the chart was almost complete, I forwarded Dr Farabee a copy of the earlier generations, numbering each individual 1, 2, 3 etc. My letter was a long time in reaching him, for again he was away from home on another scientific expedition—this time in Brazil. I asked if there was a *Benjamin Hyde* in the American family? At the end of May I received a letter from him from Barbados, dated May 16th, 1914, in which he says:

"Your letters of November 26th, 1913, and February 9th, 1914, have just reached me. For the past ten months I have been in the interior of Northern Brazil and Southern British Guiana, out of touch with the rest of the world...*You have settled the whole question. 2 is Benjamin Hyde.* His mother had short fingers, and she was the only one of her family who had. She had eleven children, but I was

unable to learn even the sex of three of them marked ??? as they died young. I had hoped to look up the English branch of this family, but am glad to know you have found it. You will have pleasure in the comparative study...."

Now, as the name *Benjamin Hyde* is a very rare combination, there can be little doubt that Farabee's family and this second English family are blood relations and belong to the same stock; the only difficulty with regard to it is the fact that in the English there is one generation more than in the American family. This may be owing to the fact that it is eleven years since the American chart was constructed, and also because marriage may not have occurred at such an early age over there as in this country, for in the English family many of the women have married whilst still "in their teens."

Moreover the relative proportion of abnormal males and females is exactly the same in these two families, viz. 61 per cent. of women in the American family and 61 per cent. in the English family. This may be a mere coincidence, but nevertheless it seems rather remarkable, and differs from the proportions found in my first family, where there were 19 females to 25 males.

Fig. 2 shows Farabee's chart of the American family, and Fig. 3 the English family.

Fig. 2. Farabee's Chart.

It will be observed that the second generation in the English family is incomplete compared with the second generation in Farabee's chart, but there is no real discrepancy. It was Benjamin Hyde's sister, who removed from America and settled in this country, who became the ancestor of the English branch. My informants know that her maiden name was Hyde, that she had a brother named Benjamin Hyde, and that their mother had *several* other children, but do not know their type or sex. As the rest of the family remained in America and lived in

Farabee's neighbourhood, he would be likely to get a more complete record than I could obtain, and hence the difference in the second generation in the two charts is accounted for.

It will be observed that Farabee has here indicated the descendants of *abnormals* only; my chart includes the offspring of *normals as well as abnormals*.

The rest of this paper refers exclusively to the English family which I have recently studied.

The abnormality, wherever it exists at all, affects *both* hands and *both* feet, as well as the stature. Both hands and both feet are always affected symmetrically.

The chart includes 50 abnormal individuals, of whom 34 are living. Through the great kindness of these people I have been able to obtain photographs of the right hand of 30 of them, and radiographs of the hand and foot of 29, as well as a few full length portraits;—a most satisfactory record, when one considers their very natural reluctance to do anything calculated to draw attention to them individually. In fact there are only two adults whom I have failed to persuade to pay a visit to the photographer and radiographer.

AN EXAMPLE OF MENDELIAN INHERITANCE.

Students of Mendelism will at once recognise that Brachydactyly as illustrated by this family conforms in a remarkable degree with certain laws enunciated by Gregor Mendel. One of these laws states that a "dominant" character is transmitted only by a member showing that character, and not by a member showing the "recessive" character, and that the recessives always breed true to the recessive character.

In this family the normal individuals are "recessives," and according to the theory *cannot* bear abnormal (short-fingered) children; and the chart shows that the normals have had normal children only. In other words, every abnormal child has an abnormal parent. As there has been no inter-marrying of abnormals, it follows that each abnormal child has had one abnormal and one normal parent—one showing Brachydactyly, and one with the normal type of hand and foot.

When abnormal individuals—the offspring of parents of two distinct types—marry *normals*, the offspring (in conformity with Mendel's Law) should be of both types, in approximately equal numbers—50 per cent. of normals and 50 per cent. of abnormals. That is what Mendel found

in the case of two types of sweet-pea, and his results have been repeatedly confirmed by other observers. The normals (recessives) should produce normals only: the abnormals (dominants) should produce normals and abnormals in about equal numbers. The chart at once shows that the *types* produced accord with this theory. What about the relative proportion of the two types among the children of the abnormals? The short-fingered children of abnormal parents ought to be equal or very nearly equal in number to the normal-fingered children of the same parents.

As a matter of fact this equality is not usually *exact*, and theory does not require it to be so, for there is a *chance* element which comes into play which makes this precise number uncertain and variable. According to Mendel the germ cells of any individual, such as one of these abnormals, are of two kinds. One kind of cell carries the factor which can produce the abnormality, and is inherited from the abnormal parent; the other kind of cell, being inherited from the normal parent, lacks this factor: and these two kinds of germ cells are present in the individual *in equal numbers*. If this be true, it must of necessity follow that the particular kind of germ cell which will take part in any given fertilization, will depend, so far as we can tell, upon a chance meeting; just as there is a chance of drawing a black or a white marble out of a hat which contains an equal number of each. It is a lottery with equal chances for both kinds. Sometimes one kind will predominate, sometimes the other, but in the long run the numbers will be approximately, if not exactly, equal.

In this family there are 50 abnormals, so that according to Mendel's theory there ought to be *about* 50 normals. The actual number of normals is 48.

Thus, instead of the exact 50 per cent. of abnormals which *might* occur in strict accordance with theory, we have 51.02 per cent. This is sufficiently near to constitute this family a striking instance of Mendelian inheritance in the human subject. This remark applies also to the other instances of Brachydactyly published by Farabee and myself, as well as to my illustrations of Minor-Brachydactyly.

If a normal member of this family were to ask whether if married to another normal (relative or non-relative) there would be any risk of having Brachydactylous children, one would be justified in replying that there was no risk whatever of such children resulting from the union, for, not in a single case has a short-fingered child been born of two normal parents. This abnormality thus differs in a remarkable

manner from certain hereditary diseases (such as colour-blindness and hæmophilia) which are known to appear in the male children of a woman, who, though showing no sign of these diseases herself, is yet able to transmit them from her affected male parent to her children.

There are six generations shown in the chart. All the individuals of the first, second and third generations are dead; of the abnormals of the fourth generation, only two are still living (Nos. 8 and 9); of the fifth and sixth generations, nearly every member (of both types) is still alive.

Of the 34 normals living at the present time, I am able to give fairly complete details of 30, and very incomplete details of the remaining four.

I am greatly indebted to Dr Geoffrey Williams of Wrexham, Dr John of Stoke-on-Trent, and Drs Bythell and Boydell of Manchester for the excellent radiographs, without which it would have been impossible to arrive at a correct interpretation of the exact nature of this interesting abnormality.

"The hands¹ and feet, as already stated, are abnormal in each affected individual, and the feet are, if anything, more abnormal than the hands, at least as regards their digits. The middle phalanx is practically or virtually—though not actually—absent from each finger and toe. The metacarpal bones are short and otherwise abnormal, but the metatarsus is scarcely, if at all, affected. Nor is the variation limited to the hands and feet, for *all* the individuals, with the exception of young children.....are below the average stature, as shown by a reference to the table of measurements" (page 338). It will be well to study each feature in turn in the following order: hands, feet, stature, etc., as measured and as revealed by photography and radiography.

FACTS REVEALED BY PHOTOGRAPHY.

A. *Hands: External aspect.*

Length. "The most conspicuous feature is the shortness, especially of the fingers; these are only slightly more than *half* the normal length, sometimes even less than half, whilst the hand looks abnormally broad. The middle finger, measured on the palmar surface from the

¹ All quotations such as this, where the source is not mentioned, are from my paper on Brachydactyly read before the Royal Society (Edin.) and serve to show features common to the two families.

base or metacarpo-phalangeal crease to the tip, is normally as long as the width of the palm at the knuckles; in these people it is approximately half," the average length in the adults being $\frac{1}{2}$. "Sometimes it is shorter than the first finger." The extreme shortness in one or two cases is due chiefly to the metacarpal bone. The hand, measured from the carpal end of the radius to the tip of the first finger, has an average length of about 5 inches. The width of the hand exceeds the normal in numerous instances.

Markings. "The skin creases are peculiar. Each finger shows only one crease, corresponding to the space between the first and third phalanx¹. It is *single*, like the one present in ordinary fingers opposite the second joint." The crease "opposite the first joint is double in most normal hands."

"Whilst it would be difficult to tell that there are only two phalanges in each finger, from an inspection of its dorsal aspect, the palmar view shows clearly that such is the case; and this is confirmed by radiography in the majority of instances."

"The palm shows two peculiar lines." "One runs straight across the hand transversally...it appears to be a union of the palmist's lines of "heart" and "head."

"The second line, often the deeper of the two, starts from the middle of this transverse line and runs to the space between the first and second fingers." This is shown very distinctly on Nos. 20, 38, and 39 (Plates XII and XIII). It is occasionally faintly visible in normal hands, but as a rule is quite absent.

"The skin is loose, and the whole hand is soft and flabby."

"The palm is very compressible, owing to the wide intermetacarpal spaces." The fingers are all "double jointed," i.e. they can easily be flexed dorsally by slight pressure applied directly to their palmar surfaces.

"A slight lateral pressure makes the palm half an inch narrower, without altering the plane of the metacarpals. The narrowing is not due to a transverse folding of the hand."

"On flexing and extending a finger at the basal (metacarpo-phalangeal joint) *slowly*, one can feel that the opposing surfaces are not uniformly curved as in the normal hand. At certain points the phalanx seems to slide over a ridge. This is what one would anticipate from an examination of the radiographs." "The ring-finger in several instances is bent at the middle, so that the tip points towards

¹ A bone of the finger.

the middle finger." This inclination is sometimes so marked that, in the position of rest, the ring-finger lies partly in front of the middle one. Strangely enough the little finger tends to project away from its neighbour, so as to leave a large gap between the two (Pl. XII, Nos. 27, 23, 41).

The skin of the back (dorsum) of the hand is very coarsely reticulated.

The mouth of the sweat glands is conspicuous.

The nails are well formed in every individual.

Strength of Grip. It is considerably below the normal average.

B. *The Feet.*

"Here the one main peculiarity is the shortness of the toes—each one apparently having only two phalanges. They are also broad and straight, with very little tendency to the extreme flexion of many ordinary toes."

In the photograph of the plantar aspect of the foot of the child (Pl. XII, No. 43), the toes are shown fully extended. Frequently the length scarcely, if at all, exceeds the width, as in the three smaller toes in this child.

FACTS REVEALED BY RADIOGRAPHY.

A. *Hands.*

The X-rays show that the middle phalanx is not really absent; a fact which could not possibly have been ascertained from the most careful ocular and digital examination. Without the aid of radiography one would conclude that there is an absence of the middle phalanx, and even the X-rays might lead to the same conclusion if the hands of *some* adults were the only ones examined. The young man, Pl. XIV, No. 37, for instance, has only two bones in each finger, and at first sight it looks as though the middle bone—the second phalanx—is altogether missing, whilst the other bones appear about normal; but if the terminal phalanx is carefully compared with the corresponding bone in a normal hand it will be seen that the base is much thicker than it should be. In most of the abnormal fingers, this thickened base is seen to form almost a perfect cube, being as broad as long. Whenever this condition is present, it will be found that there is no vestige of the *separate* middle phalanx.

What is the nature of this cubical basal portion of the terminal phalanx? and what has become of the second phalanx?

In the *middle* finger of every abnormal adult member of this family, the second phalanx is always present as a *separate* bone, but invariably it is so far reduced in length as to become a cube; and the terminal (third) phalanx is of the normal pyramidal shape. The ring-finger also, in certain individuals, resembles the middle finger in these particulars, having a normal-looking third phalanx, and a separate, short, cubical middle phalanx. Whenever the terminal phalanx has a cubical base, this base corresponds in shape with the second phalanx present as a separate bone in the middle finger. In fact "this cubical basal portion is the second (middle) phalanx that has become ankylosed¹ to the terminal phalanx. The pyramidal distal portion of these bones corresponds to the ungual phalanx and the basal cubical portion to the middle phalanx."

"It does not exist as a separate bone in either the index or little finger in a single adult."

"What, then, has happened to the middle phalanx? It varies in two respects from the normal:

1. It is always very short.
2. It generally becomes ankylosed to the base of the terminal phalanx.

The fact that the middle phalanx is abortive, but not completely absent, is proved conclusively by an examination of the radiographs of the hands of young children, before ankylosis has occurred, and in whom the middle phalanx is seen to be invariably present, but at the same time abnormal. Each phalanx, in the ordinary (normal) hand, consists at first of cartilage (gristle) and is gradually transformed into bone, or, as it is termed, becomes ossified, such transformation being limited for some time to two parts, one forming the great bulk or *shaft* of the phalanx, and the other a thin plate or disc at the base. This disc is termed the *Epiphysis* of the phalanx.

That narrow strip of the cartilage which is situated between the bony shaft and the bony epiphysis usually remains unossified until about the twentieth year in each phalanx. The epiphysis of each phalanx is seen to be situated at its base; but the epiphysis of each metacarpal (palmar) bone is at the distal extremity. So long as the strip of cartilage between the shaft and the epiphysis remains

¹ Joined by bony union.

unossified, the bone can increase in length, but after its ossification further growth does not occur.

The disc seen in some of the radiographs at the base of the terminal phalanx *looks* very much like an epiphysis, but is really the second phalanx in a very abortive condition.

In some instances in children there is only one piece of developing bone visible between the shafts of the third and first phalanges; normally, three can be seen: namely, (1) the epiphysis of the third, (2) the shaft of the second, and (3) the epiphysis of the second phalanx, and it is only possible by a comparison of several hands to say what this single bone represents. I believe it always represents the shaft of the second phalanx.

In no single instance does the radiograph show the presence of an epiphysis at the base of the second phalanx.

It is thus clear that there is no real absence of the second phalanx in any individual, but merely a rudimentary condition, and that at a certain stage of development there is a union of this with the terminal phalanx.

"The essential feature of the abnormality apparently consists in an absence of the epiphysis at the base of the second phalanx. It is possible that the epiphysis is also missing, in some instances, from the third phalanx, and that the two phalanges (second and third) consist at first of a single piece of cartilage."

Premature union of shaft and epiphysis is of general occurrence in this family. Such union should not occur until about 20 or 21 years of age. In No. 41, a girl of 15, it is already complete, and the radiograph would pass for that of an adult hand. In this instance, moreover, the union is not recent, for the epiphyseal lines are already almost entirely obliterated. The hand of the boy (Pl. XV, No. 31), aged 8 years and 4 months, shows a very interesting peculiarity. The epiphysis of the third phalanx, though still separate from its own shaft, has already united with the shaft of the second phalanx in the first and little fingers. This seems to point strongly to the opinion expressed above, that these two phalanges are sometimes represented by only one piece of cartilage. *This premature union of the shaft and epiphysis is no doubt one great factor in producing the characteristic shortening.*

"The index and fourth fingers seem more aberrant than the second and third, as they (the former) never show the middle phalanx as a separate bone in the adult." In Pl. XV, No. 27 (*aet.* 19), the union has occurred in the little finger: in the index finger the union is not yet complete, but it is evidently taking place.

"Functionally, the fingers are all reduced to the bi-phalangeal condition, and thus come to resemble thumbs."

The Thumb. "The change in the *thumb* consists of a shortening of the first phalanx which is reduced to a cube. There is no attempt at ankylosis however." In some children the epiphysis is present, but in others it is apparently absent, for if present union must have occurred before the seventh year. The terminal phalanx has the epiphysis in every individual.

"The so-called metacarpal bone of the thumb is in reality the first phalanx, for here the epiphysis is clearly seen at the *base*, whereas in the other metacarpals the epiphysis is seen at the *distal* end of the bones."

"The Metacarpals vary more or less in different individuals, but, as a rule, where ossification is complete, they are abnormal." "The head of each metacarpal is distinctly nodulated in many cases....The middle one is the shortest of the four," in No. 25, and in this woman the whole palm shows the maximum degree of shortening. In Pl. XV, No. 27, the metacarpals of the middle and little fingers are much shorter than the other two.

"*Sesamoid* bones are well shown in several cases."

Spaces. The intermetacarpal spaces are generally increased in width, especially between the *heads* of the bones. Sometimes it is about three times the normal width. This accounts for the easy compressibility of the palm laterally.

MEASUREMENTS: AVERAGE IN INCHES.

The following table gives the average length of the middle finger, the hand, and the stature in eleven abnormal adult women (8 married and 3 spinsters).

Middle Finger	Hand	Stature
$1\frac{3}{8}$	$4\frac{7}{8}$	$57\frac{1}{2}$

In normal women these measurements are about 3, $6\frac{3}{4}$ and 64 respectively.

There is only one abnormal adult male member of the family now living, and I have not yet seen him. The only measurement he has sent me is that of the middle finger, which he says is $1\frac{5}{8}$ " in length.

There are three abnormal males between 17 and 21 years of age, and their averages are as follows¹:

Middle Finger	Hand	Stature
$1\frac{7}{8}$	$5\frac{9}{16}$	$61\frac{1}{2}$

¹ In the first family: these measurements were

1—15/16	$5\frac{1}{2}$	61.
---------	----------------	-----

The longest middle fingers measure 2 inches (Nos. 8 and 28). The shortest measures $1\frac{1}{4}$ inches (No. 21), and three measure $1\frac{1}{2}$ inches (adults).

The tallest male is $64\frac{1}{2}$ inches in height (No. 28); the tallest woman is 59 inches (No. 26); the shortest woman 52 inches (No. 9).

B. *Feet.*

The shortness of the feet is "due solely to the abortive middle phalanx, and the somewhat stunted growth of the first."

"The so-called first phalanx of the big-toe is considerably shortened"; in most cases it is reduced to a cube, and in some the shortening goes even further.

"The middle phalanx in the other toes has become ankylosed to the terminal one in every adult." This ankylosis "occurs at an earlier age in the foot than in the hand, though the hand as a whole is more abnormal."

"There is an absence of the basal epiphysis of the first phalanx of the big-toe, though it is conspicuous in the first phalanx of each of the other toes (in children), so that this bone appears to be really homologous with the second phalanx of the other toes."

Metatarsus. There is very little wrong with the metatarsal bones, in which respect they differ from the metacarpals. In two cases (Pl. XIV, Nos. 27 and 28) the third and fourth are the shortest of the series.

Thigh and Leg. These are shorter than normal, but the numbers especially of the normals are too few from which to calculate a reliable average. There is no doubt, however, that the short

Stature is entirely accounted for by the shortness of the lower limbs. When the adults are seen seated they do not strike one as being particularly short people, but on standing up it is at once apparent that they are considerably below the average height. The spinal column is therefore approximately of the normal length.

This fact seems fairly clear from the photographs of the two brothers (Pl. XI): the one on the right (No. 37) is $17\frac{1}{4}$ years old, $59\frac{1}{2}$ inches in height, and is Brachydactylous; the other one is 14 years old, 3 inches taller, and is not Brachydactylous. The difference in the length of the legs is very obvious, which no doubt accounts for their difference in stature.

The infant (No. 48) is three-quarters of an inch taller than his normal twin sister, in this respect presenting a striking and unexpected

contrast to the rule that holds good in adults and youths, so that the arrest of growth occurs in some later period of life, probably beginning at or soon after the age of 2 years.

The shortness of hands is well shown in Pl. XII, Nos. 23, 37 and 41, where they are represented with normal relatives for the sake of comparison.

Symmetry.

"In every instance the hands are exactly symmetrical, as shown both by photography and radiography (Pl. XV); and I believe the same rule holds true with regard to the feet, though *both* feet were not examined in all cases." This remarkable symmetry obtains even in those cases that show special peculiarities of any of the bones, either in shape or ossification, or in the presence and time of union of the epiphysis.

Persistence.

There is no apparent tendency to revert to the normal type, for the hands and feet of No. 9 are no more abnormal than those of her grandchildren; neither is there any other indication of the abnormality disappearing: on the contrary the numbers seem to be on the increase, the abnormals in the last four of the six generations in the English branch being as follows:

3 in the third generation.			
8	„	fourth	„
12	„	fifth	„
23	„	sixth	„

One would expect *a priori* that individuals whose fingers are short, stumpy and below the average in strength would be so much handicapped in the "struggle for existence" that they would be swamped by the general population, and in an uncivilized state of society this surely must have occurred; but the conditions of modern society have afforded them such chances of survival that their numbers are steadily increasing, yet it seems clear that they *are* handicapped to some extent, for all the employed men and women are engaged in occupations where there is no great need for manual dexterity: their social position is below that of their normal relatives.

Marriage.

Marriage is the fashion amongst these short-fingered people and they "go off," as one of the women expressed it, before their normal relatives. The great majority of those shown in the chart as not having had children died in infancy. Several adult normals remained unmarried. Why should this be? There must be some special fascina-

Measurements of Abnormals in inches.

Initials	Number in Chart	Sex	Age	Middle finger	Hand		Ulna	Tibia	Height
					Length	Width			
Mrs E. T. ...	9*	female	64	$1\frac{1}{2}$	5	3	$7\frac{1}{4}$	10	52
Mrs M. ...	16	"	37	$1\frac{3}{8}$	$5\frac{1}{4}$	$3\frac{1}{2}$	—	—	$58\frac{5}{8}$
M. H. ...	17*	"	33	$1\frac{3}{4}$	5	$3\frac{1}{2}$	—	—	$58\frac{1}{2}$
M. H. ...	18*	"	31	$1\frac{3}{4}$	$5\frac{1}{4}$	$3\frac{1}{4}$	9	$11\frac{1}{2}$	$60\frac{1}{2}$
Mrs M. A. L. ...	19*	"	43	$1\frac{3}{4}$	$5\frac{1}{4}$	$3\frac{1}{2}$	$7\frac{3}{8}$	—	$55\frac{1}{4}$
Mrs R. F. ...	20*	"	41	$1\frac{1}{2}$	$4\frac{3}{8}$	$3\frac{1}{8}$	9	13	$59\frac{1}{2}$
Mrs S. E. J. ...	21*	"	39	$1\frac{1}{4}$	4	$2\frac{1}{4}$	$6\frac{1}{4}$	9	56
Mrs O. R. ...	23*	"	26	$1\frac{5}{8}$	$4\frac{5}{8}$	$3\frac{1}{2}$	$7\frac{1}{2}$	$10\frac{1}{2}$	$57\frac{1}{2}$
W. H. ...	24	male	39	$1\frac{5}{8}$	—	—	—	—	—
Mrs O. G. F. ...	25*	female	37	$1\frac{1}{2}$	$4\frac{1}{2}$	$3\frac{1}{8}$	8	$11\frac{3}{4}$	57
Mrs R. T. ...	26*	"	28	$1\frac{5}{8}$	5	$3\frac{1}{2}$	$7\frac{3}{8}$	—	59
H. H. ...	27*	"	19	$1\frac{3}{4}$	5	$3\frac{1}{2}$	$8\frac{1}{2}$	$11\frac{3}{4}$	59
S. H. ...	28	male	18	2	6	4	—	—	$64\frac{1}{2}$
E. M. ...	29*	female	$12\frac{3}{4}$	2	$5\frac{1}{8}$	3	—	—	58
E. M. ...	30	"	$10\frac{1}{2}$	$1\frac{1}{10}$	$4\frac{1}{10}$	$2\frac{7}{8}$	—	—	$52\frac{3}{4}$
J. M. ...	31*	male	$8\frac{1}{12}$	$1\frac{1}{2}$	$4\frac{1}{8}$	$2\frac{3}{8}$	—	—	$47\frac{1}{2}$
P. M. ...	32*	"	$5\frac{3}{12}$	$1\frac{1}{4}$	$3\frac{3}{8}$	$2\frac{3}{8}$	—	—	42
W. L. ...	33	"	17	$1\frac{3}{4}$	$5\frac{1}{2}$	$3\frac{5}{8}$	$9\frac{1}{2}$	—	$60\frac{1}{2}$
E. L. ...	35	"	$10\frac{1}{12}$	$1\frac{1}{8}$	$4\frac{1}{2}$	$2\frac{1}{2}$	—	—	43
L. L. ...	36*	female	7	$1\frac{1}{4}$	4	$2\frac{1}{2}$	$5\frac{1}{4}$	$8\frac{1}{2}$	43
O. F. ...	37*	male	$17\frac{1}{2}$	$1\frac{7}{8}$	$5\frac{1}{2}$	$3\frac{1}{4}$	9	13	$59\frac{1}{2}$
W. F. ...	38*	"	8	$1\frac{1}{10}$	4	$2\frac{1}{2}$	$6\frac{3}{8}$	9	$45\frac{1}{2}$
E. F. ...	39*	female	6	$1\frac{1}{10}$	$3\frac{5}{8}$	$2\frac{3}{8}$	6	$7\frac{5}{8}$	$42\frac{1}{2}$
A. F. ...	40*	"	$2\frac{1}{2}$	$\frac{7}{8}$	$2\frac{3}{4}$	2	$4\frac{1}{2}$	6	$32\frac{1}{4}$
S. J. ...	41*	"	15	$1\frac{5}{8}$	5	3	$7\frac{1}{2}$	11	$54\frac{1}{2}$
A. J. ...	42*	"	$3\frac{3}{4}$	$1\frac{1}{8}$	3	$2\frac{1}{10}$	5	$6\frac{3}{8}$	32
E. R. ...	43*	"	$4\frac{1}{2}$	$1\frac{1}{4}$	$3\frac{1}{2}$	$2\frac{1}{4}$	$5\frac{1}{8}$	8	$38\frac{1}{4}$
W. H. ...	44	male	8	$1\frac{1}{2}$	—	—	—	—	—
D. H. ...	45	"	$1\frac{1}{2}$	1	—	—	—	—	—
J. A. F. ...	46	"	12	$1\frac{3}{8}$	$4\frac{1}{4}$	$2\frac{3}{8}$	$7\frac{1}{2}$	11	$51\frac{1}{2}$
A. F. ...	47*	"	$6\frac{3}{4}$	$1\frac{1}{10}$	$3\frac{5}{8}$	$2\frac{1}{4}$	$6\frac{1}{4}$	8	41
A. F. ...	48	"	$1\frac{3}{12}$	$\frac{7}{8}$	$2\frac{5}{8}$	$1\frac{3}{4}$	$3\frac{1}{2}$	$4\frac{3}{4}$	$28\frac{1}{4}$
C. C. T. ...	49*	"	7	$1\frac{5}{8}$	4	$2\frac{1}{2}$	$5\frac{1}{2}$	$9\frac{1}{2}$	$44\frac{1}{4}$

* Individuals marked by an asterisk are illustrated on Plates XII—XV.

tion that compensates for the manual defects, but what it is I am unable to say. It might be suggested that their amiability and obliging disposition (of which this paper is sufficient evidence) will afford an explanation, but I should not like to say that the normal members fall short of them in this respect.

Of the first family there are 27 or 28 abnormals now living: so that altogether there are at least 60 people of the Brachydactylous type in England at the present time.

The normals do not appreciably differ in the above dimensions from the general population, and I give the details of only a few children and adolescents for comparison with their short-fingered relatives.

Measurements of Normal Members of the Family.

Initials	Sex	Age	Middle finger	Hand		Ulna	Tibia	Height	
				Length	Width				
A. J.	...	male	17	3 $\frac{1}{8}$	7 $\frac{1}{2}$	3 $\frac{1}{2}$	9 $\frac{1}{2}$	12 $\frac{1}{4}$	67 $\frac{1}{2}$
W. J.	...	„	12	2 $\frac{1}{2}$	6	2 $\frac{3}{4}$	7 $\frac{1}{4}$	10 $\frac{3}{8}$	51
A. F.	...	female	16	2 $\frac{1}{2}$	6	2 $\frac{7}{8}$	9	12	58 $\frac{3}{4}$
F. W. F.	...	male	14	2 $\frac{3}{4}$	6	3	8 $\frac{1}{4}$	11 $\frac{1}{2}$	54 $\frac{3}{4}$
L. O. F.	...	female	1 $\frac{8}{12}$	1 $\frac{1}{2}$	3	1 $\frac{3}{4}$	3 $\frac{3}{4}$	5 $\frac{1}{2}$	27 $\frac{1}{2}$
E. F.	...	male	19	2 $\frac{3}{4}$	6 $\frac{3}{4}$	3 $\frac{3}{8}$	9	12	60 $\frac{1}{2}$
J. F.	...	„	14	3	6 $\frac{1}{2}$	3 $\frac{1}{2}$	9 $\frac{1}{2}$	14 $\frac{1}{4}$	62 $\frac{1}{2}$
R. F.	...	female	10	2 $\frac{1}{2}$	5 $\frac{3}{4}$	2 $\frac{3}{4}$	7 $\frac{1}{2}$	11	50
E. H.	...	„	20	3 $\frac{1}{8}$	7	3 $\frac{1}{4}$	—	—	61 $\frac{1}{2}$
H. H.	...	„	17	3	7	3 $\frac{1}{2}$	—	—	59 $\frac{1}{2}$
S. M.	...	male	2 $\frac{1}{2}$	1 $\frac{3}{4}$	—	—	—	—	33 $\frac{1}{4}$

Part of the expense of this investigation has been defrayed by the "Earl of Moray Fund for Original Research" of the University of Edinburgh. I very gratefully record my thanks to the Trustees for their generous help.

EXPLANATION OF PLATES.

The numbers are those given in the chart on p. 327 and also in the Table on p. 338.

PLATE XI.

Fig. 1. Shows three brachydactylous females; a child, its mother and grandmother (No. 43, 23 and 9). The short stature of the women is evident when compared with that of the author which is 5 feet 8 $\frac{1}{2}$ inches.

Fig. 2. Shows two brothers. The one on the right is brachydactylous and two years older than the one on the left, who is normal.

PLATE XII.

Photographs of hands and a foot. The numbers refer to those in the chart. Their ages and other particulars are given in the table of measurements on page 338. The same applies to the following plates.

PLATE XIII.

All these hands are brachydactylous.

PLATE XIV.

Radiographs of hands and feet, all abnormal.

PLATE XV.

Radiographs of hands. Both hands are shown in order to illustrate the bilateral symmetry.

Fig. 2.

Fig. 1.

No. 23 and normal.

No. 37 and normal.

No. 20.

No. 43.

No. 39.

No. 27.

No. 41 and normal.

No. 9.

No. 21.

No. 17.

No. 23.

No. 18.

No. 25.

No. 19.

No. 26.

PLATE XIII

No. 32.

No. 42.

No. 36.

No. 43.

No. 38.

No. 47.

No. 40.

No. 49.

No. 20.

No. 39.

No. 37.

No. 20.

No. 27.

No. 31.

No. 41.

No. 18.

No. 27.

No. 29.

No. 31.