Original Articles.

THE CHLOROPHYLL TEST OF GASTRIC MOTILITY.*

By FRANKLIN W. WHITE, M.D., BOSTON,

Instructor in Medicine, Harvard University; Assistant Visiting Physician, Boston City Hospital.

THERE is no need to emphasize the importance of the motor function of the stomach. It is well recognized that it is the stomach's chief function, or at least the one which makes most trouble, if it is deranged. A simple, exact test of motility has, therefore, great value.

Our tests of secretion are simple and accurate. Our tests of the motor function of the stomach are not as simple nor as accurate, especially for the finer changes in motility.

The chlorophyll test was devised by Boas¹ two years ago, with the hope of avoiding some of the defects in previous tests of gastric motility, and the present work was done to make a trial of this test in a considerable series of cases, to see how far this hope has been justified.

Some of the defects in previous tests, which the chlorophyll test aims to avoid, are as follows: the Leube meal, which is much used, is a rather heavy meal (plate of soup, steak or roast beef, potato, a roll, a glass of water), and is often taken with little appetite by sick patients, which must affect the motility of the stomach. A patient who has been on a light diet often dislikes the sudden change to the heavy Leube meal and may have difficulty in tolerating it, and vomiting may result. In a considerable number of tests of motility, rather coarse food, such as raisins, green vegetables and so forth, are given. Those are not without a disagreeable effect in some cases of ulcer.

In those tests of motility which are made by aspirating the residue in the stomach after the course of a number of hours or over night, there may be a considerable addition to the bulk of the residue due to gastric secretion or hypersecretion. Thus the element of secretion enters to confuse what was aimed to be a test of motility. The ten- and twelve-hour tests for gastric residue do not show the finer changes in the motility of the stomach.

The x-ray is often used at present for testing gastric motility. The expense and the necessity for repeated visits are sometimes an objection. Also the bismuth often leaves the stomach at quite a different rate from the food.

To illustrate this briefly: A recent patient with complete achylia gastrica had no residue after a test breakfast at the end of one-half hour, on three separate occasions (the stomach

was washed out each time to make sure it was empty). The stomach showed a six-hour residue after a bismuth meal.

Another patient with catarrh of the stomach, who showed a three-ounce residue of food eight hours after a Leube-meal, showed an empty stomach four and a half hours after a bismuth meal. Such illustrations might be multiplied.

With some of these defects of previous tests in mind, we can easily appreciate the value of a simpler and more accurate method.

The technic of the chlorophyll test is as follows:

The test is made in two parts.

First part: The patient, on a fasting stomach, drinks 400 c.c. of water, which has been colored pale green by the addition of twenty drops of a watery chlorophyll solution. At the end of half an hour, the residue is aspirated with the stomach-tube, and the amount noted.

Second part: In order to make the test more accurate and that no residue might escape, the stomach is then washed out with a 1% watery solution of sodium-bicarbonate. The amount of this solution used for lavage depends upon the amount of residue aspirated at the end of one-half hour. Let the thirty-minute residue equal x . The stomach is washed out with 400 minus x c.c. The residue and the wash waters are put together and made up to 400 c.c. (if necessary) and the green color of this mixture compared with dilutions of the original chlorophyll water taken (400 c.c. water, 20 drops chlorophyll solution), and in this way the per cent. of the colored water, which has passed out of the stomach in one half-hour, is estimated.

This test was based upon Moritz' observation, that 500 c.c. of water leaves the normal stomach in one-half to three-quarters of an hour, and to Boas' own observation in normal cases, that 400 c.c. of water taken into the fasting stomach, leaves a residue in a half hour of 50 to 60 c.c.

The chlorophyll solution is tasteless, easily obtained and is not absorbed from the stomach. It is, however, easily precipitated by saliva and mucus and the patient is warned not to swallow saliva after drinking the chlorophyll water.

The advantages claimed for this test of motility are, that the chlorophyll water gives little stimulus to secretion, causes no pyloric reflex and is a pure test of motility. The watery residue is easily recovered. The water is easy for the patient to take and it constitutes a rapid, simple method for the doctor.

In making a systematic trial of this test, we were especially interested to see if these claims were justified, and whether or not the test was equal or superior in accuracy to the older tests, whether it would detect finer grades of motor disturbances. In our series of 50 cases, we have systematically compared the test with the Leube seven-hour test. The x-ray has been used in many of the cases, but primarily for other purposes, the detection of defects or deformities in the outline of the stomach. While the x-ray

* Read at the seventeenth annual meeting of the American Gastro-Enterological Association at Atlantic City, June 22 and 23, 1914.

findings are given in our table, they are not emphasized as a guide to the accuracy of the chlorophyll test.

We have found the technic of the test simple and easy for both doctor and patient. Careful aspiration of the residue is very necessary, but this is easy with the watery solution.

The second part of the test, which was designed to increase its accuracy, *does not* do this in our opinion and we have omitted it for the following reasons: We have found that the chlorophyll pigment clings to the mucus in the stomach and makes the second part of the test inaccurate. This is readily seen in the table of the cases by comparing the amount of residue in c.c. in the first column with the amount of color-residue in the second.

In Case 1 we see a 5 c.c. residue, well within normal limits, and a 50% color-residue far above the normal (which is about 10%). The same thing is seen in a varying degree in Cases 3, 4, 5, 6, 8, 19 and 22. The amount of color retained in the stomach in these cases runs more or less parallel with the amount of mucus in the stomach, to which it could cling. The mucus precipitated and held back pigment, while the water escaped. In short, the amount of residue in c.c. in the stomach was a far better guide to the speed of emptying than the color test. Several other stains were tried, in hope of finding one which would prove tasteless, and not be absorbed or held back by the mucus, but with poor success.

The cloudiness of the mixture of residue and wash water, the presence of minute food particles, the occasional reflux of bile, sometimes made accurate color comparison difficult. Filtration was not practicable, as thereby most of the pigment was filtered out.

It was found that complete aspiration of the residue was the rule, making the second part of the test unnecessary and often misleading, and this colorimetric part of the test was omitted after the first 22 cases, and our estimate of the motor power of the stomach based upon the number of c.c. of residue at the end of one-half hour. This omission makes the test shorter and easier for both patient and doctor. It is not absolutely necessary to add the chlorophyll to the water, in making the test, but it is a convenient way of identifying the residue with the liquid taken.

We have not found the chlorophyll a stimulus to gastric secretion, neither did it check it and, if marked hyper-secretion was present, we found it wise to empty the stomach in the morning, before making the chlorophyll test.

RESULTS.

The following table gives our results. First in the 45 cases where the chlorophyll and Leube tests agreed, in patients with normal and with abnormal motility, then in the 5 cases where somewhat divergent results were obtained

with the two tests. In judging of the accuracy of the test, we have used chiefly the amount of residue seven hours after a Leube meal, for comparison. In 90% of cases, the results are the same as with the Leube meal and in the remaining 10% the differences are all minor ones. No important change in motility was missed and in this way the test was satisfactory for diagnosis. A few individual cases, 5 out of 50, have shown some difference in results with the different tests used. In Case 46 there was a slightly increased residue with the half-hour chlorophyll test and a normal residue with the 7-hour Leube test. There was evidently a slight, early retention of food in the stomach. The same is true to a less degree of Case 50. In Case 47, one of ptosis, the chlorophyll residue is close to normal. The Leube 7-hour residue was something more than normal. The stomach emptied quite well at first and later showed fatigue. In Case 49, a duodenal ulcer, the chlorophyll test was normal and the Leube 7-hour test showed considerable residue. There was marked hyper-secretion in this case, which undoubtedly explains the considerable residue at the end of seven hours, and the differences between the two tests. As a pure test of motility, I believe the chlorophyll was more accurate than the Leube test, in this case.

The same thing is true to a less degree in Case 48, one of gastric ulcer.

Comparing the results of different tests of gastric motility is not a simple matter. It is difficult to say what is the "standard test." The Leube 7-hour test has been chosen for this series, because it is one of the best known and most widely used and most satisfactory tests which we have had.

Gastric motility is rather complex. The x-ray has especially helped us to understand this of late years. The activity of peristalsis is not uniform at different periods of digestion. In one case, it is very active at an early stage, later fatigued and slow. In another, it is sluggish at the beginning and later "gets its second wind."

We do not believe that the occasional slight differences which we have found between the chlorophyll and the Leube tests are important in diagnosis and we do not believe that the chlorophyll test will detect finer differences in motor function than the Leube test.

The theoretical objection may be raised, that it is better to combine the test of motility with a test of secretion, since the latter affects the time of the emptying of the stomach and that a "real meal" is more natural than a long drink of green water. These objections do not seem to be valid. The practical results of this test in diagnosis have been good, sometimes apparently better than the controls. It is a satisfactory quantitative test of a very simple type, the amount of residue varying with the grade of disturbance.

Kemmerling² and Wartensleben³ have recently reported results which resemble ours. Both found difficulties in the colorimetric part of the test and Wartensleben, working in Boas' own

clinic, advises omitting the second part of the test. The per cent. of cases in which the chlorophyll test ran parallel with other tests of motility was smaller, in both series, 62% Kemmerling, 78% Wartensleben (90% in our series). This difference is largely due to their frequent comparisons of the chlorophyll test with the bismuth meal and x-ray test of motility. I do not believe that the bismuth test of motility should be used as a criterion to judge the value of other tests of motility for the reason stated above, namely, that bismuth often leaves the stomach at an entirely different rate from ordinary food or drink.

A. BOTH TESTS AGREE.

(1) Normal Motility.

	CHLOROPHYLL	TEST.	Leube	X-ray
	Residue	Residue	7 Hr.	Empty
	in c.c.	Color	Residue	in
		% Mucus		Hours
1. Neurosis	5	50	+++	0
2. Achylia	9	10	0	0
3. Colitis	0	45	+++	0
4. Adhesion cecum	10	20	++	0
5. Neurosis				
6. " "	12	20	++	0
7. " "	6	0	0	0
8. Duod. ulcer	45	20	+	0
9. Neurosis	5	15	+	0
10. " "	24	10	+	0
11. Gallstones	15	—	—	0
12. Duod. ulcer	0	—	—	0
13. Neurosis	10	—	—	0
14. Fatty liver	12	—	—	0
15. Catarrh	0	—	—	0
16. Constipation	10	5	—	0
17. Cholecystitis	5	5	—	0
18. Gastr. ulcer	0	0	—	0
19. Chronic appen-				
dix	40	20	+	0
20. Constipation	20	15	—	0
21. Chronic appen-				
dix	5	10	—	0
22. Gastr. ulcer	20	30	++	0
23. Neurosis	0	5	—	0
24. " "	0	5	—	0
25. Phthisis	2	5	—	0
26. Cancer	10	—	—	0
27. Neurosis	40	—	—	0
28. " "	60	—	—	0
29. Ptois	10	—	—	0
30. Gall-stones	50	—	—	0

(2) Abnormal Motility.

31. Tabes	90	15	+	8
32. Catarrh	180	30	—	90
33. Cancer	540	—	—	390
34. " "	300	—	—	412
35. " "	435	—	—	340
36. Duod. ulcer	240	—	—	490
37. Gastr. ulcer	170	75	++	35
38. Ptois	180	—	—	40
39. Benign stenosis	205	—	—	130
40. " "	290	—	—	175
41. " "	200	—	—	200
42. Duod. ulcer	210	—	—	160
43. Benign stenosis	340	—	—	180
44. Ptois	120	—	—	145
45. Benign stenosis	300	—	—	420

B. DIVERGENT RESULTS.

46. Gastr. ulcer	90	—	—	0
47. Ptois	65	—	—	57
48. Gastr. ulcer	3	—	—	25
49. Duod. ulcer	30	—	—	120
50. Ptois	75	—	—	0

SUMMARY.

The second colorimetric part of the test is better omitted and the estimate of the motor power of the stomach based on the number of c.c. of residue found at the end of one-half hour.

We may answer the question whether it is better than older methods of testing gastric motility by saying that it is simpler than most; makes little demand on the appetite, and is very easy on the stomach. It is readily used in patients who are vomiting, or who are on a very simple light diet, where a big meal is uncomfortable or impossible. The sources of error are few and probably unimportant and no considerable disturbance of motility escapes detection. It is an easy method of following results of treatment of gastric stasis.

It is probably not superior in accuracy and not likely to replace the best of the older tests, but it is simpler and easier for both doctor and patient in a large group of cases and gives us an extra "string to our bow."

REFERENCES.

- ¹ Boas, J.: Beitrag zur Motilitätsbestimmung, Deutsch. med. Woch., No. 10, 1912.
- ² Ueber die Boas'sche Chlorophyllprobe zur Bestimmung der Magenmotilität, Arch. f. Verdauungskr., 1914, Band xx, S. 49.
- ³ Ueber Motilitätsbestimmung des Magens mit besondere Berücksichtigung der Boas'sche Chlorophyllmethode. *Idem*, S. 66.

A NOTE ON THE INCREASE OF TOTAL NITROGEN AND UREA NITROGEN IN THE CEREBROSPINAL FLUID IN CERTAIN CASES OF INSANITY, WITH REMARKS ON THE URIC ACID CONTENT OF THE BLOOD.*

By H. M. ADLER, M.D., BOSTON,

Chief-of-Staff, Psychopathic Hospital, Boston; Assistant Professor of Psychiatry, Harvard Medical School.

AND

B. H. RAGLE, BOSTON,

Interne, Psychopathic Hospital.

I. CEREBROSPINAL FLUID DETERMINATIONS.

THE cerebrospinal fluids examined by us were obtained from patients at the Psychopathic Hospital by lumbar puncture. The earlier speci-

* Being Contributions from the Psychopathic Hospital, Number 50 (1914.16). (*Bibliographical Note*.—The previous contribution, Number 49 (1914.15), by Mary L. Gerrin, entitled "Impressions of a General Hospital Nurse on Beginning Work at the Psychopathic Hospital, Boston," published in the BOSTON MED. AND SURG. JOUR., Vol. clxxi, No. 13, Sept. 24, 1914.)