could concentrate on this subject, they would do industry a real service, for nearly all industrial operations such as punching, shearing, forging, milling, spinning, and, of course, the turning of metals, are plastic flow phenomena.

During experiments I carried out with heavy lathes in 1908 for the purpose of finding the most economical high-speed steel to use, I encountered some chips which were not only straight but actually presented concavity to the tool face, and I have one of these chips now. They were produced at very high speed on steel, and are mentioned in the discussion of a paper read before the Siemens' Stafford Engineering Society in 1908 (Proc., vol. 1, p. 93), on "The Plastic Deformation of Solids."

Brewster's beautiful photo-elastic method and Professor Coker's important applications of it enable the stresses during elastic strain of the tool and material in the region A to be computed, but Taylor, in the work cited, has shown how a tool should be forged and supported on the saddle to give it maximum life and maximum strength.

Unfortunately for engineering industry in this country, nearly all lathes are built with the vertical space between the upper surface of the tool rest and the line of centres far too small to enable Taylor's important conclusions to be put into practice.

ALAN POLLARD.

Imperial College of Science and Technology,
November 29.

I GATHER from Mr. H. S. Rowell's letter published in NATURE for December 9 that, while interested in the subject of the flow of metals in shavings, he is not altogether familiar with the work that has already been done on the subject. In a comprehensive "Mémoire sur le rabotage des métaux" (which cannot be so well known as I have hitherto believed) published more than forty years ago, M. H. Tresca investigated the question of the curling of shavings, both experimentally and mathematically, the actual flow of the metal (expressed by a *coefficient de réduction*) being especially selected for study under very varied conditions. The following quotation indicates only part of the scope of the work: "Ces phénomènes sont aussi ceux dans lesquels, pour la première fois, les métaux les plus durs, tels que l'acier, le fer, se comportent en réalité comme le plomb, comme le savon, comme le cire, nous dirions presque comme les liquides, tant est complet le rapprochement que l'on doit faire entre les rides de nos différents copeaux et de véritables vagues de métal."

The memoir is published as one of the "Mémoires présentés par divers savants à l'Académie des Sciences de l'Institut de France," tome xxvii., 1883. Those familiar already with the beauty of the results obtained will pardon this effort to direct the attention of others to the work.

E. N. DA C. ANDRADE.

Artillery College, Woolwich,
December 11.

The Secondary Spectrum of Hydrogen.

SINCE the negatively charged hydrogen atom is known to exist, from work on positive rays, it seemed likely that Silberstein's particular solution of the three-body problem, applied by him to the case of neutral helium (*Astrophys. Jour.*, September 1922) should also be applicable in this case. Consequently the formula used by him was modified so as to apply to hydrogen (charge E instead of $2E$, and hence N instead of $4N$), and also a small but important correc-

tion was made to the value of N so as to take account of the fact that with two electrons instead of one, the correction to the mass of the electron for the finite mass of the nucleus is no longer the same.

It was assumed as a first approximation that the electrons would be arranged antipodally, and consequently the forces would be again central. So Curtis's value of N for hydrogen was corrected so as to apply to a nucleus of infinite mass:

$$N_{\infty} = N_H \left(1 + \frac{m}{M} \right).$$

Frequencies were then calculated from the formula

$$\nu = N_{\infty} \left(\frac{1}{n_1^2} + \frac{1}{n_2^2} - \frac{1}{m_1^2} - \frac{1}{m_2^2} \right).$$

These frequencies were then sought for in the secondary spectrum of hydrogen; it is known that negatively charged atoms are to be found in hydrogen at fairly high pressures with intensity quite comparable with that of the positively charged atom ("Rays of Positive Electricity," p. 39). As a result it was found that 47 lines in the secondary spectrum agreed with the calculated values within an absolute error of one unit of frequency, taking integral values of n_2 and m_1 from 1 up to 10, and values of m_2 from 1 to 15, while n_1 was taken as 2 and 3.

This means that the frequencies can be looked on as a kind of "summation tone," being the sums of a Balmer or a Paschen frequency and a frequency in the infra-red.

It was also found that in several cases a physical similarity of behaviour was common to "series" of the lines grouped according to the m 's and n 's concerned, though this was not exclusively true. As a standard of reference for the observed frequencies the values obtained by Merton and Barratt (Phil. Trans. A, 1922, pp. 388-400) were employed.

As typical may be given the following:—

Formula.	Calculated.	Observed.	Error $\delta\nu$.	Character.
$\frac{3 \cdot 9}{2 \cdot 6}$	16934.9	16934.51	+0.39	2 + + CD
$\frac{3 \cdot 10}{2 \cdot 6}$	17192.3	17192.14	+0.16	6 + + CD + + HP + He
$\frac{3 \cdot 12}{2 \cdot 6}$	17527.6	17527.47	+0.13	3 + + CD + HP + + He
$\frac{3 \cdot 13}{2 \cdot 6}$	17638.8	17639.89	-1.09	0 + + CD + + He
$\frac{6 \cdot 6}{3 \cdot 3}$	18289.8	18288.26	+0.54	0
$\frac{6 \cdot 8}{3 \cdot 3}$	19623.4	19622.74	+0.66	0 + He
$\frac{6 \cdot 10}{3 \cdot 3}$	20240.7	20240.71	-0.01	3 + LP

In the foregoing table, the figures in the last column refer to intensity and the symbols to the physical properties of the lines as given by Merton and Barratt (*loc. cit.*).

It is hoped to complete these and similar calculations shortly and also to investigate the conditions under which these lines should be enhanced.

A. C. MENZIES.

Physics Laboratory, The University, Leeds,
December 8.

Science and the Empire.

THE admirable sentiments expressed in the leading article in NATURE of December 16 will undoubtedly be re-echoed by every scientific worker in the country. In stating, however, that the British Science Guild is the only organisation which exists to undertake the propaganda work "for the extension of an understanding of the influence of scientific research and its results," the very effective propaganda which is being carried out by scientific workers themselves under the

ægis of the National Union of Scientific Workers is overlooked.

Of this body you say "it is a Trade Union affiliated, we believe, to the Labour Party, and it exists to secure suitable conditions of work and payment for its members rather than for the extension of natural knowledge." In that statement truth and error are intermingled. The National Union of Scientific Workers is a registered Trade Union; it registered as such when industrial and Civil Service joint councils on the Whitley plan were being set up and when it was announced that none but members of Trade Unions would be given representation on those bodies. The Employers' Federations registered as Trade Unions also and for the same reasons. But the National Union of Scientific Workers is not affiliated to the Labour Party or to any political party; it has no political funds, and it imposes no restraints upon the political activities of its members, three of whom stood for Parliament at the recent election, one in the Conservative interest and the other two as Labour candidates.

Again, while it is true that the National Union of Scientific Workers exists to secure suitable conditions of work and payment for its members—and all other scientific workers incidentally—it considers that the best way to do this is by raising the professional standard of scientific workers by improved training and education, and making them aware of their importance as citizens on one hand, and on the other, by pointing out to employers and captains of industry that it is an economy to employ the best scientific workers, to encourage research, and to assist the universities. In order to persuade private employers, corporations and governing bodies to deal justly with scientific staffs, it is true that the Union would be prepared to follow the methods employed by such bodies as the British Medical Association; but it believes with the British Science Guild that the attitude of the general public towards science is due to ignorance or apathy. Accordingly, it puts propaganda efforts, designed to cure these diseases, in the forefront of its programme, hoping thereby to increase the demand from industry and the State for the best scientific knowledge. It is ready to co-operate with any other body for this purpose, and to assist any political party with its advice on matters appertaining to science and scientific workers. It believes, however, that scientific workers themselves must be their own propagandists, and that the first step towards really effective action is unity in the profession of science.

A. G. CHURCH,
General Secretary.

National Union of Scientific Workers,
25 Victoria Street,
Westminster, London, S.W.1,
December 18.

[THE National Union of Scientific Workers is an occupational organisation; therefore its propaganda efforts, useful as they are, are naturally regarded by the public as arising from self-interest. The British Science Guild, on the other hand, requires no technical or other qualification for membership; and, as was pointed out in our article, it bears the same relation to scientific workers that the Navy League does to the Royal Navy. It seems to us that a body of this type, in which citizens engaged in many and diverse departments of national life are concerned, can afford much more effective and disinterested support of science than is possible by any group consisting of members of the profession alone. That was the main point of the article to which Major Church refers, and we see no reason to depart from it.—EDITOR, NATURE.]

The Hermit-crab (*E. bernhardus*) and the Anemone (*C. (Sagartia) parasitica*).

IN NATURE of December 2, p. 735, I described observations and experiments on the common hermit-crab (*E. bernhardus*) with its messmates, the anemone (*C. (Sagartia) parasitica*) and the polychæte worm *Nereis fucata*. By the kindness of Mr. Hugh Main, it has been pointed out that the observations mentioned above with regard to the natural position of the anemone confirm those of J. Sinel (p. 39, "An Outline of the Natural History of our Shores," 1906). Sinel states that "the woodcuts that appear in many text-books—even our high-class ones—which represent this anemone and its congener, are in one respect incorrect. The anemone is always represented as upright—palm-tree like—on the top of its equipage, as if its chief object were display—or a ride. . . . I have invariably found the anemone affixed to the rear of the shell and in such a position that when the hermit is at a meal or even moving about, the margin of the tentacles just touch the ground, like some patent sweeping-machine. It, no doubt, finds this position a paying one."

Sinel's unique and fascinating book contains a fund of information hidden away in a popular description of natural history on the shore. It is plain that, owing probably to the popular character of the book, many naturalists have passed over important original observations described therein by Sinel, whose knowledge of the biology of the shore has probably never been equalled.

Sinel's observations were previously unknown to me, but the agreement in the two sets of independent observations is valuable in opposing a traditional error, and will be sufficient to establish the correctness of the interpretations; the natural position of the anemone on the hermit-crab was clearly first shown by Sinel.

J. H. ORTON.

Marine Biological Laboratory,
Plymouth, December 13.

Winter Thunderstorms.

MAY I through your columns again ask for reports of thunderstorms occurring in the British Islands between January 1 and March 31? With the help of your readers and of observers of the British Rainfall Organization I was able to collect a mass of information on winter thunderstorms for 1916, 1917, 1918, and 1920, from which it appears that on more than 40 per cent. of the days in question, thunderstorms occurred somewhere in the British Islands. In collaboration with the Meteorological Office I propose to collect information again. The chief points to be noticed are the times at which the storms occur, and especially the times of passage of such storms as pass overhead; whether a severe storm or whether there are only one or two flashes of lightning or only one or two claps of thunder; whether there is a change of wind or a drop of temperature with the storm; whether there is rain, hail, or snow; in the case of lightning seen at night the direction in which it occurs; and any other information the observer thinks of interest. Reports are wanted especially from the west and north of Scotland, and from the south-west, west, and north-west of Ireland, but any information however slight from any district in the British Islands will be of great use to the investigation. Reports should be sent by postcard or letter to my address (not to the Meteorological Office).

C. J. P. CAVE.

Stoner Hill, Petersfield, December 20.