This process of disappearance and recurrence has continued at about the same intervals up to the present time. As long as she was inoculated at regular intervals, the discharge remained absent, but from four to six weeks after inoculations were discontinued a recurrence was sure to come. This represents that class of chronic otitis media which is only benefited by vaccine treatment. It is not an uncommon experience to encounter cases under this class in which the discharge diminishes about 50 per cent. under a course of inoculations, but at no time disappears entirely. Of the seventeen patients with chronic otitis media treated by vaccine therapy, three recovered completely, five were improved, and nine showed no improvement at all.

In this class of cases vaccine therapy should be tried when the ordinary measures fail. Its use should be considered as a means capable of producing some measure of good in about 50 per cent. of cases, with an occasional remarkable result.

CONCLUSIONS

In conclusion, I feel safe in saying that in subacute otitis media no other means of treatment is capable of producing such uniformly good results as vaccines. Their early use would greatly diminish the large number of chronic cases, so common to every physician, and so annoying to the individual.

The results in chronic cases, on the other hand, are so variable, that its use as a routine should not be advised. It should be tried, however, when other measures have failed.

VANADIUMISM

WALTON FOREST DUTTON, M.D.
CARNEGIE, PA.

Vanadium is a metal named by Sefström, in 1830, in honor of the northern goddess Vanadis (the German Freya). The metal is of no importance technically, although various forms of it are valuable. The pentoxid is used in photography as a developer. Vanadium chlorid and trioxid are used as mordant in printing fabrics; the trioxid is used also in the manufacture of steel, in making easily malleable and ductile alloys. Vanadium pentoxid has been used in diseases due to defective metabolism; there is not, however, enough clinical evidence to justify its use as a medicine.

Vanadiumism is a chronic intoxication caused by ingestion or absorption of some forms of vanadium, either industrially, medicinally, or accidentally. The principal lesions are found in the lungs, kidneys, and gastro-intestinal tract. The lungs are highly congested and show a marked destruction of alveolar epithelium. The kidneys reveal congestion due to continuous irritation and, at times, acute hemorrhagic nephritis. The gastro-intestinal tract shows evidence of irritation and inflammation.

Persons employed in establishments where the various kinds of vanadium are produced, and more especially vanadium trioxid, suffer from chronic poisoning. For example, laborers in reduction plants, who are exposed to the dust and fumes, are most susceptible on account of their occupation. The medicinal use of small doses of the pentoxid will frequently cause poisoning. Accidental poisoning may come from using or wearing clothing dyed by vanadium chlorid, trioxid, or pentoxid.

Anemia is an early symptom (a peculiar cachexia), not altogether unlike chlorosis. There is, at first, marked

increase in the number of corpuscles and hemoglobin, which is shortly followed by reduction of cells and hemoglobin. At times, there is disintegration of the red cells. The cough is a prominent and characteristic symptom. It is dry, irritating, and paroxysmal, becoming so intense that hemorrhages are frequent and severe, even causing death. The workers in the manufacture of the various products of vanadium are very susceptible to tuberculosis and, unless removed from the labor, rapidly succumb to the disease. Emaciation, irritation of the nose, eyes, and throat are always in evidence. Anorexia, nausea and diarrhea (often absent), followed by obstinate constipation, indicate gastro-intestinal involvement. Albumin, casts and blood, are often present in the urine.

Continuous exposure to the poison leads to fine tremors of the extremities, headache, neuroretinitis, amaurosis, vertigo, hysteria, and melancholia. The drug is eliminated by the kidneys; it may be found also in the feces, sputum, and saliva. Dr. Omar T. Cruikshank, in his experiments on animals, has been able to observe the toxic effects which coincide with those found in man.

The diagnosis is not difficult when the history of exposure to the various forms of vanadium is at hand. Accidental origin is often obscure and diagnosis is not, at once, determined.

Watts gives the test for vanadium in the following:

All vanadium compounds heated with borax or phosphorus salt in the outer blowpipe flame produce a clear bead, which is colorless if quantity of vanadium is small, yellow when it is large; in the inner flame the bead acquires a beautiful green color.

Vanadic and chromic acids are the only acids whose solutions are red; they are distinguished from one another by the vanadic acid becoming blue and the chromic acid green by deoxidation.

When a solution of vanadic acid, or acidulated solution of an alkaline vanadate, is shaken up with ether containing hydrogen dioxide, the aqueous solution acquires a red color like that of ferric acetate, while the ether remains colorless. This reaction will serve to detect the presence of one part of vanadic acid in 40,000 parts of liquid.

In the absence of grave renal, blood, nervous, and lung involvement, the prognosis is good. Where there is active inflammation of the kidneys and lungs the prognosis is unfavorable, but depends greatly on the extent of the lesions. Fatal terminations frequently ensue, as a result of continuous exposure to the poison.

The prevention of vanadiumism is difficult in vanadium works, owing to the carelessness of employees and employers. Means to allay and carry off the dust and fumes should be employed constantly. Perfect ventilation and the use of respirators are imperative. The nasal and oral cavities should be thoroughly cleansed with some efficient alkaline spray, such as the ordinary Dobell solution, followed by a mentholated oil spray. The stomach should be washed out, and later the intestines freely evacuated with a saline laxative. The cough may be allayed by giving terpin hydrate $\frac{1}{4}$ gr., heroin $\frac{1}{8}$ gr., and creosote $\frac{1}{2}$ minim, every two hours with counter-irritations of iodine or mustard applications over the chest. The inhalation of stimulating vapors is salutary. Iron, calisaya, and strychnin will meet the needs of anemic, nervous and debilitated conditions. Cod-liver oil may be given with advantage. Active outdoor exercises are essential. Turkish, Russian or cabinet baths may be given to aid elimination.

401 Fourth Avenue.