In der Ephemeride ist gegeben RA., Dekl. und geozentrische Distanz für jeden fünften Tag von Mai 1 bis zum Schluß des Jahres 1914.

Zur Zeit der Sonnenfinsternis 1914 Aug. 21 ist die Lage der Dreieckspunkte die folgende:

$$\begin{array}{lcl} \alpha & L_4 = 10^h 17^m 9 & L_5 = 8^h 49^m 3 \\ \delta & +12^\circ 8' & +17^\circ 43' \\ \Delta & 1.308 & 0.976 \end{array}$$

Berechnet man die Distanzen der Dreieckspunkte von der Sonne zur Zeit der zentralen Finsternis im nördlichen Europa, so erhält man:

$$\begin{array}{l} L_4 = 4^\circ 36' \text{ östlich von der Sonne} \\ L_5 = 17 \text{ } 41 \text{ westlich } \text{ » } \text{ » } \end{array}$$

Der zweite Dreieckspunkt hat, wie man sieht, eine ziemlich günstige Lage.

Ephemeride für 0^h m. Z. Berlin.

L_4				L_5			
1914	RA.	Dekl.	Δ	RA.	Dekl.	Δ	
Mai	1	2 ^h 1 ^m 6	+13° 54'	1.394	1 ^h 7 ^m 8	+5° 5'	0.838
	6	3 0.8	17 23	1.364	1 34.0	7 11	0.950
	11	3 39.2	20 30	1.306	2 2.0	10 7	1.068
	16	4 18.7	23 1	1.218	2 35.8	13 37	1.178
	21	4 56.4	24 27	1.099	3 14.7	17 17	1.268
Juni	26	5 27.0	25 8	0.965	3 57.0	20 36	1.325
	31	5 47.2	24 41	0.832	4 40.5	23 9	1.349
	5	5 54.3	23 39	0.720	5 23.4	24 47	1.345
	10	5 49.4	22 8	0.639	6 4.6	25 29	1.322
	15	5 36.0	20 29	0.596	6 43.1	25 22	1.285
Juli	20	5 21.1	18 58	0.588	7 18.7	24 35	1.237
	25	5 10.3	18 10	0.612	7 50.9	23 12	1.183
	30	5 7.8	18 1	0.657	8 20.1	21 31	1.120
	5	5 12.9	18 27	0.717	8 45.2	19 28	1.053
	10	5 25.2	19 19	0.788	9 6.3	17 20	0.980
	15	5 44.2	+20 21	0.866	9 22.4	+15 15	0.902

1914	RA.	Dekl.	Δ	RA.	Dekl.	Δ	
Juli	20	6 ^h 8 ^m 8	+21° 19'	0.948	9 ^h 32 ^m 0	+13° 28'	0.824
	25	6 38.6	22 0	1.033	9 33.0	12 15	0.750
	30	7 13.1	22 11	1.118	9 23.1	12 5	0.693
Aug.	4	7 51.8	21 36	1.198	9 3.1	13 10	0.674
	9	8 33.9	20 3	1.265	8 41.5	15 1	0.712
	14	9 17.8	17 25	1.308	8 32.1	19 28	0.822
	19	10 1.3	13 48	1.318	8 40.3	17 50	0.943
	24	10 42.0	9 33	1.294	9 2.2	17 32	1.080
Sept.	29	11 19.0	5 3	1.243	9 31.0	16 11	1.199
	3	11 52.2	+ 0 32	1.184	10 2.0	14 3	1.294
	8	12 22.1	- 3 34	1.105	10 33.0	11 22	1.366
	13	12 47.6	7 23	1.030	11 3.1	8 20	1.416
	18	13 13.0	10 44	0.953	11 32.2	5 6	1.448
	23	13 34.0	13 34	0.878	12 0.5	+ 1 45	1.463
	28	13 51.2	15 46	0.803	12 28.4	- 1 42	1.464
Okt.	3	14 3.5	17 12	0.733	12 56.1	5 8	1.459
	8	14 9.0	17 42	0.671	13 23.8	8 33	1.421
	13	14 7.5	16 58	0.621	13 47.6	11 58	1.364
	18	13 58.3	15 4	0.595	14 20.5	15 0	1.316
	23	13 44.0	12 12	0.602	14 49.3	17 52	1.235
Nov.	28	13 32.5	9 24	0.653	15 17.4	20 18	1.133
	2	13 26.6	7 33	0.775	15 42.4	22 5	1.012
	7	13 40.1	8 8	0.876	15 59.4	22 49	0.878
	12	14 1.3	10 7	1.015	15 59.7	23 5	0.760
	17	14 31.0	13 7	1.145	15 42.5	19 45	0.678
Dez.	22	15 5.6	16 22	1.250	15 18.6	16 34	0.671
	27	15 42.5	19 27	1.325	15 4.0	14 26	0.723
	2	16 20.2	21 57	1.374	15 2.8	13 51	0.804
	7	16 58.1	23 48	1.401	15 12.3	14 30	0.893
	12	17 35.3	24 57	1.411	15 29.0	15 52	0.978
	17	18 11.8	25 27	1.406	15 50.6	17 33	1.055
	22	18 47.3	25 17	1.389	16 16.0	19 19	1.124
	27	19 22.3	24 33	1.358	16 44.5	20 58	1.185
	32	19 54.1	-23 17	1.315	17 15.7	-22 23	1.236

Observatoriet Lund, 1914 Mai 13. H. Henic.

Observatoriet Lund, 1914 Mai 13.

H. Henic.

Positions of Variables and Asteroids Discovered on Photographs of Star-fields.

Taken with the 40-inch Reflector of the Lowell Observatory. By C. O. Lampland.

In the course of the examination of an extensive series of negatives of star-fields — these observations forming part of an investigation by Director Lowell, not yet completed — a number of variable stars and asteroids have been found. During the past year about 800 negatives have been made and examined in connection with this work. The negatives were all taken with the 40-inch reflector, and the examinations and the measurements were made with a Zeiß comparator equipped with a Blink-Mikroskop.

All the areas observed have been photographed in duplicate and the exposures were from two to three days apart, though at times longer intervals elapsed between the duplication of the negatives. The plates used were Seed's »30« and Lumière's »Sigma«, but mostly the former on account of their greater freedom from defects. At the beginning of the work the plates were exposed thirty minutes. Later it was found that the exposure time could be reduced to

eight minutes and still give sufficiently strong images of the fainter stars required in the examination of the fields. The size of the plates used was five by seven inches. While this is a much larger area than the field of good definition of the instrument, the large plates have been of considerable advantage in the examinations, in checking — on the overlapping ends of negatives of adjacent fields — images having indications of variability, asteroids, and defects. The settings at the telescope were made on the intersections of the hour-circles and parallels of declination (at intervals of four minutes in right ascension and one degree in declination) in Schönfeld's and Gould's Durchmusterung charts, so that the plates in their longest direction have considerable overlap, the linear scale of the negatives corresponding to one degree of arc being 3.8 inches.

The estimates of the magnitudes must be considered as being rather rough. Many of the images are outside of

the field of good definition, with the added difficulty that no faint stars with well determined magnitudes were to be had for comparison. Due to the overlap of negatives of adjacent fields many of the stars here given appear on four plates.

As many of the stars are comparatively faint, the measured positions should have been accompanied with small charts of the surrounding fields. When additional negatives

can be obtained of some stars, such fields will be prepared and published.

The observations — the work at the telescope and the examinations of the negatives with the comparator — are the joint labors of Mr. *E. C. Slipher* and myself. The writer must be held responsible for such deficiencies as may appear in other parts of the work.

Variables.

	α 1913.0	δ 1913.0	1913	1913	Obs.
29.1914 Librae	15 ^h 15 ^m 27 ^s	-20° 8'3"	May 5 < 15 ^m	May 7 15 ^m	L
30.1914 »	15 21 55	-4 0.6	July 28 < 15	Aug. 30 14-14 ^m 5	S
31.1914 »	15 22 11	-21 55.7	May 8 15.5	May 10 not vis.	S
32.1914 »	15 22 38	-22 7.1	May 8 13	May 10 < 15.5	S
33.1914 »	15 37 49	-16 11.2	May 26 14.5	May 29 13	L
34.1914 »	15 38 2	-15 52.6	May 26 13.5	May 29 < 15	L
35.1914 »	15 42 20	-12 8.6	July 3 12.5	July 24 13	L
36.1914 »	15 42 56	-5 32.1	July 29 14.5	Sept. 2 15.5	S
37.1914 »	15 44 16	-16 27.6	May 26 < 15	May 29 14.5	L
38.1914 »	15 48 4	-18 18.8	June 27 13.5	July 3 15	L
39.1914 »	15 51 38	-18 35.6	June 27 15.5	July 3 14	L
40.1914 Scorpii ¹⁾	15 58 54	-12 52.5	June 30 13	July 4 13	L
41.1914 »	15 59 1	-12 56.5	June 30 15	July 4 14.5	L
42.1914 »	16 2 5	-19 23.8	June 27 15.5	July 2 15	L
43.1914 »	16 3 47	-11 52.7	July 3 15.5	July 24 13.5	S
44.1914 »	16 3 49	-19 15.0	June 27 15	July 2 not vis.	L
45.1914 »	16 2 24	-16 32.7	May 26 16?	May 29 15	L
46.1914 »	16 4 34	-15 23.6	June 3 14	June 6 > 15	L
47.1914 »	16 4 54	-14 23.5	May 30 < 14	June 2 < 15	L
48.1914 »	16 6 34	-15 2.0	June 3 15?	June 6 16?	L
49.1914 »	16 7 23	-21 43.3	May 31 14.5	June 2 12.5	S
50.1914 »	16 10 1	-16 27.7	May 27 14.5	May 29 < 15	L
	α 1875.0	δ 1875.0			
51.1914 Librae	15 17 19	-28 16.5	June 25 12.5	June 28 14.5	S
52.1914 »	15 18 23	-25 38.4	July 1 14	July 22 < 15	S
53.1914 »	15 33 25	-29 20.9	June 26 12.5	June 30 14	L
54.1914 »	15 34 38	-26 9.5	June 4 15	June 7 13.5	L
55.1914 »	15 35 12	-28 25.7	June 23 14	June 26 < 15	L
56.1914 Lupi	15 35 9	-29 48.3	June 27 14.5	July 2 < 15	L
57.1914 Librae	15 37 10	-28 19.6	June 23 < 14	June 26 < 15	L
58.1914 »	15 38 10	-29 11.9	June 26 13.5	June 30 15	L
59.1914 Scorpii	15 43 27	-24 29.1	May 31 14	June 3 12	S
60.1914 »	15 44 11	-23 35.6	May 31 < 15?	June 3 13.5	S
61.1914 »	15 47 51	-29 5.1	June 25 14	June 28 < 15	L
62.1914 »	15 50 21	-24 7.9	June 2 < 15	June 5 14	S
63.1914 »	15 50 21	-28 58.8	June 25 13.5	June 28 not vis.	L
64.1914 »	15 51 17	-28 3.0	June 25 < 15?	June 28 14	L
65.1914 »	15 52 54	-28 43.5	June 25 < 15	June 28 13	L
66.1914 »	15 55 32	-27 44.8	June 25 14.5	June 28 15.5-16	L
67.1914 »	15 56 25	-24 49.8	June 3 < 15	June 6 14.5	L
68.1914 »	15 57 29	-25 8.8	June 3 < 15	June 6 14	L
69.1914 »	15 58 32	-25 32.4	June 3 < 14	June 6 14	L
70.1914 »	15 58 35	-23 2.0	May 29 15.5	May 31 14.5	L
71.1914 »	15 58 38	-28 35.3	June 25 14.5	June 28 13.5	L
72.1914 »	15 58 44	-28 1.3	June 25 15	June 28 16	L
73.1914 »	15 58 47	-23 3.3	May 29 15	May 31 16?	L
74.1914 »	15 59 20	-28 56.3	June 25 13	June 28 15	L
193.1904 » ²⁾	15 59 41	-23 19.6	May 29 15	May 31 13	L
75.1914 »	16 0 47	-28 4.8	June 25 15	June 28 16	L
76.1914 »	16 2 51	-23 8.3	May 29 not vis.	May 31 14	L

< = fainter than, > = brighter than. L = *Lampland*, S = *Slipher*.

¹⁾ Discovered on comparison of negatives with *Wolf-Palisa* chart.

²⁾ Vergl. A. N. 167.161. *Red*.

Asteroids.

Planet	1913 M. M. T.	α 1913.0	δ 1913.0	Mag.	Obs.	Planet	1913 M. M. T.	α 1913.0	δ 1913.0	Mag.	Obs.
May						June					
147 Protogeneia	5 11 ^h 44 ^m	15 ^h 13 ^m 36 ^s	19° 2'5	13-14	S	new 1913 c	3 13 ^h 21 ^m	16 ^h 2 ^m 39 ^s	23° 49'2	13.5-14	S
184 Dejopeja	5 12 31.5	15 19 23	20 6.1	12	L	663 Gerlinde	3 15 20	15 58 7	14 50.1	13-14	L
1913 SS	6 11 29	15 25 25	13 30.9	14	L	663 Gerlinde	4 14 44	15 57 24	14 42.5	13-14	L
184 Dejopeja	7 12 19.5	15 17 50	20 0.3	12	L	160 Una	5 11 18	15 32 5	23 55.6	12.5	S
new 1913 a	8 12 41	15 46 51	22 13.8	15?	S	new 1913 d	7 12 15.5	15 31 12	27 5.2	13-13.5	S
	10 14 32	15 44 45	22 3.9	14.5?	S	new 1913 e	10 12 54.5	15 27 13	28 25.6	8-9	L
663 Gerlinde	26 13 29.5	16 4 25	15 55.4	12	L	new 1913 f	25 10 57.0	15 20 23	28 35.0	13.5	S, L
322 Phaeo	28 11 25	15 52 34	22 33.7	13.5	S	new 1913 g	26 12 32.5	15 57 2	20 5.8	14	L
162 Laurentia	29 12 9.5	15 32 29	22 42.1	12-12.5	S	new 1913 h	26 12 32.5	15 59 28	19 39.6	13.5	L
438 Zeuxo ¹⁾	29 12 46.5	15 41 27	22 41.0	12	L		27 11 43	15 58 53	19 41.7	14	L
77 Frigga	29 12 46.5	15 40 27	22 51.1	12	L	new 1913 i	27 13 22	16 6 3	18 22.7	14.5	L
322 Phaeo	30 11 44	15 50 40	22 23.7	13.5	S	379 Huenna	27 13 22	16 6 3	18 34.4	13-14	L
162 Laurentia	31 9 39	15 30 56	22 38.7	12-12.5	S	1913 f	28 10 30	15 19 5	28 26.6	13	S, L
77 Frigga	31 10 8	15 38 45	22 45.4	11.5	L	379 Huenna	28 12 56	16 5 30	18 33.0	14	L
438 Zeuxo	31 10 8	15 39 35	22 41.3	11.5	L	July					
June						379 Huenna	2 13 22.5	16 3 32	18 29.6	14	L
new 1913 b	3 9 43	15 43 43	24 1.1	14	S	663 Gerlinde	3 11 16	15 43 36	11 56.9	12.5	L

¹⁾ 438 Zeuxo: GröÙe im B. J. falsch, muß 11^m4 statt 12^m9 heißen. *Berberich*.

Lowell Observatory, Flagstaff, Arizona, 1914 January.

C. O. Lampland.

Orbite de l'étoile variable X Vulpeculae. Par M. Luizet.

Après quelques années d'interruption, j'ai observé de nouveau X Vulpeculae en 1912 et 1913; j'ai utilisé les mêmes étoiles de comparaison qu'en 1906, et l'échelle de lumière résultant de l'ensemble des observations diffère peu de celle obtenue avec les premières (A. N. 4195):

$$\begin{aligned} A &= BD + 26^{\circ}37'25'' & L &= 17^{\text{deg}}4 \\ a &= +26\ 37'39'' & & 11.8 \\ b &= +26\ 37'46'' & & 3.7 \\ c &= \text{Anonyme} & & 0.0 \end{aligned}$$

Les maxima ci-dessous ont été tirés des 74 observations faites pendant ces deux années:

Maxima.					
E	T. m. Gr.	O-C	E	T. m. Gr.	O-C
404 1912 Juill.	9.85	+0.26	432 1913 Janv.	2.85	+0.33
410 Août	16.60	+0.09	433 »	9.10	+0.26
411 »	22.80	-0.02	455 Mai	28.10	+0.24
412 »	29.20	+0.06	457 Juin	9.60	+0.10
419 Oct.	12.25	-0.13	458 »	15.90	+0.09
420 »	18.55	-0.15	461 Juill.	4.70	-0.07
422 »	31.50	+0.17	464 »	23.80	+0.07
423 Nov.	6.70	+0.05	477 Oct.	13.80	-0.08
428 Déc.	9.15	-0.10	478 »	20.20	+0.01
430 »	20.95	+0.07	479 »	26.70	+0.19
431 »	27.50	+0.30			

Les comparaisons O-C montrent qu'il n'y a pas lieu de modifier actuellement les éléments que j'avais obtenus en 1906:

$$\begin{aligned} \text{Max.} &= 2417040.732 \text{ (t. m. Gr.)} + 6131896 \cdot E \\ M - m &= 1.95. \end{aligned}$$

Pour obtenir la courbe moyenne de lumière, j'ai d'abord divisé les 126 observations, ordonnées au préalable suivant

les valeurs croissantes des intervalles Δt comptés à partir du maximum, en 25 groupes de 5, sauf le dernier qui en contient 6; puis, afin de diminuer la grandeur des écarts de ces 25 points à la courbe régulière tracée au milieu d'eux, et faciliter, par conséquent, le tracé de cette courbe, je les ai combinés de la façon suivante: soient $a, b, c, d, e \dots$ les moyennes de ces 25 groupes; la valeur b a été remplacée par $B = \frac{1}{4}(a+2b+c)$

ce qui revient à faire les moyennes $\frac{1}{2}(a+b)$ et $\frac{1}{2}(b+c)$, et à en prendre de nouveau la moyenne; de même c a été remplacée par $C = \frac{1}{4}(b+2c+d)$ etc.

Les résultats de ces combinaisons sont les suivants:

Δt	L	Δt	L	Δt	L	Δt	L
0.069	13 ^{deg} 69	2.066	9 ^{deg} 47	3.955	6 ^{deg} 75	5.652	11 ^{deg} 36
0.326	13.58	2.324	9.10	4.145	6.79	5.878	13.01
0.614	13.41	2.580	8.70	4.398	6.72	6.052	13.71
0.892	12.75	2.814	8.06	4.666	6.34	6.172	13.79
1.197	11.52	3.067	7.52	4.904	6.33		
1.463	10.55	3.399	7.22	5.155	7.34		
1.755	9.99	3.730	6.93	5.413	9.27		

La courbe de lumière tracée au milieu de ces 25 points montre que X Vulpeculae est une Céphéide régulière, c'est-à-dire que sa variation lumineuse ne présente pas d'arrêt, ou de maximum secondaire, pendant la diminution d'éclat.

Éléments orbitaux. X Vulpeculae est trop faible pour pouvoir être étudiée spectrographiquement; mais comme elle offre tous les caractères visuels des Céphéides brillantes que le spectroscope a montré être des systèmes binaires, on peut admettre qu'elle a une constitution analogue à la leur, et chercher à déterminer ses éléments orbitaux par la méthode que j'ai indiquée ¹⁾. Je rappellerai que cette détermination est basée:

¹⁾ M. Luizet. Les Céphéides considérées comme étoiles doubles, avec une monographie de l'étoile variable δ Céphée. Annales de l'Université de Lyon. Gauthier-Villars, Paris.