SURGERY

UNDER THE CHARGE OF

J. WILLIAM WHITE, M.D.,

FORMERLY JOHN RHEA HARTON PROFESSOR OF SURGERY IN THE UNIVERSITY OF PENNSYLVANIA AND SURGEON TO THE UNIVERSITY HOSPITAL,

AND

T. TURNER THOMAS, M.D.,

ASSOCIATE PROFESSOR OF APPLIED ANATOMY AND ASSOCIATE IN SURGERY IN THE UNIVERSITY OF PENNSYLVANIA; SURGEON TO THE PHILADELPHIA GENERAL HOSPITAL AND ASSISTANT SURGEON TO THE UNIVERSITY HOSPITAL

Post-traumatic Ossification of the Region of the Elbow-joint.—LEHMANN (*Dent. Zeitschr. f. Chir.*, 1914, cxxvi, 213) made a study of this condition on the basis of thirty-seven cases. He says that the elbow more than other joints shows a predilection for post-traumatic ossification. This is in part explained by its anatomical formation. These ossifications were found after traumas of all kinds, but especially after luxations. The trauma is the primary cause, but an individual predisposition must be assumed. The view of Machol that only reduced dislocations are followed by ossification is not borne out, although an injudicious method of reduction increases the tendency considerably. The bone formation can develop from the intramuscular connective tissue as well as from the periosteum. Practically there is no difference between the ossification of myogenous and that of periosteal origin, since both occur alongside of each other. The neighboring soft tissues participate in the structure of the callus. A proper treatment of the recent injury is the best prophylaxis against the ossification. It should aim at the most complete rest and preservation of the injured soft tissues, and should favor absorption. Vigorous massage and passive movements should be avoided. A fibrolysin cure is to be recommended for the persistent thick induration. The complete ossification can undergo spontaneous absorption, so that conservative treatment should receive much consideration. An operation is indicated only when there are long continued and severe disturbances of function and when there are complications from pressure on nerves and bloodvessels. In general, operations should not be done before the termination of the process of ossification.

The Function of the Periosteum in Bone Transplants, Based on Four Human Transplantations without Periosteum and Some Animal Experiments.—McWILLIAMS (*Surg., Gyn., and Obst.*, 1914, xviii, 159), on the basis of his clinical and experimental work, reached the following conclusions: If the periosteum be excised from an area of bone and a section of bone within this area, going into the medullary cavity, be removed (not, however, involving the whole diameter of the bone) and the endosteum scraped away as well, then it is found that the cavity fills in perfectly with new bone, which of necessity must come from the