17. *The CARBONIFEROUS SUCCESSION in GOWER (GLAMORGANSHIRE), with NOTES on its FAUNA and CONDITIONS of DEPOSITION.* By ERNEST EDWARD LESLIE DIXON, B.Sc., A.R.C.Sc., F.G.S., and ARTHUR VAUGHAN, M.A., D.Sc., F.G.S. (Read March 9th, 1910.)

[PLATES XXXVIII-XLI.]

CONTENTS.

	Page
I. Introduction (with Maps on pp. 478-79)	477
II. Lithological Characters of the Zones (E. E. L. D.)	480
(1) Eastern District.	
(2) North-Western District.	
(3) South-Western District.	
(4) Tabulated Synopses of the Sequence:—(i) in each District; (ii) generalized for the whole Area.	
III. Note on the Pseudobreccias of the D-Zone (E. E. L. D.)	507
IV. Note on Lagoon-Phases and the Origin of Radiolarian Cherts (E. E. L. D.)	511
(1) Introduction and Definition.	
(2) The <i>Modiola</i> Phases—Calcareous Lagoon-Phases—of Gower.	
(3) The Radiolarian Phase—Cherty Lagoon-Phase—of Gower.	
(4) Meaning of the Difference between the <i>Modiola</i> Phases and the Radiolarian Phase of Gower.	
(5) Topographical Position of Lagoon-Areas.	
(6) Earth-Movements accompanying Lagoon-Conditions.	
(7) Lagoon-Phases outside the South-Western Province.	
V. Interpretation of the Lithological Sequence (E. E. L. D.)	532
(1) Preliminary Remarks.	
(2) Variations in Depth and other Conditions of Deposition over Gower as a whole at Successive Times; Conclusions; Earth- Movements.	
(3) Relations between the Different Districts of Gower at Suc- cessive Times; Conclusions; Earth-Movements. Summary of §§ III, IV, & V; Acknowledgments.	
VI. Notes on the Delimitation of the Upper from the Lower Avonian in the South-Western Province	542
(1) From the Stratigraphical Standpoint (E. E. L. D.).	
(2) From the Faunal Standpoint (A. V.).	
VII. Faunal Lists (A. V.)	543
VIII. Palæontological Notes (A. V.)	553
(A) Corals.	
(B) Brachiopods.	
IX. Summary of Palæontological Sections	565

I. INTRODUCTION [E. E. L. D. & A. V.].

THE object with which the Avonian rocks of Gower (figs. 1 & 2) described in the present paper were examined was primarily the comparison of their faunal sequence with that of Bristol and other parts of the South-Western Province. In this connexion valuable pioneer work on parts of the sequence had already been done by

[478]

Dr. W. B. Gubbin¹; and it was desired that our knowledge should be extended to the whole development, as also to the relationship of the zones that are based on brachiopods and corals² with the representative of the Pendleside Series which had been recognized in Gower by Dr. Wheelton Hind.³ For in Gower alone of the whole province was such a representative known.⁴ Between 1899 and 1902 the area had been mapped by the officers of H.M. Geological Survey, Dr. A. Strahan, Mr. R. H. Tiddeman, and Mr. B. S. N. Wilkinson, and our work was greatly facilitated by the knowledge then obtained, which was generously placed at our disposal and has since (in 1907) been published in the official maps and memoirs.⁵

Our joint field-work, the examination of the Mumbles-Bishopston ground and the southern coast, was carried out in 1905, and in the following year one of us (E. E. L. D.) extended the work into the north-western part of the area in connexion with the Geological Survey; a summary of the information then obtained has since been published.⁶

II. LITHOLOGICAL CHARACTERS OF THE ZONES [E. E. L. D.].

The Avonian rocks of Gower (see map, fig. 2, p. 479, based on the Geological Survey maps previously mentioned) form part of the southern margin, the 'South Crop,' of the South Wales coal-basin. They present several outcrops, ranging west-north-west on the whole, as they have shared in the powerful Armorican folding about axes having that direction, which is the dominant structural feature of the whole area. The folds are chiefly elongated periclinal, and the several Avonian outcrops along the limbs are continuous one with the other.

These outcrops represent the deposits of an area of the Avonian sea which extended for some distance both eastwards and westwards (that is, parallel to the coast-line lying at no great distance to the north), and northwards and southwards; and the bearing of this fact on the development in different districts will be pointed out.

In the east, between Mumbles and Cefn-y-Bryn, only one major fold—the Cefn-y-Bryn pericline—appears above the sea, and the Lower Carboniferous rocks which form its eastward-pitching 'nose,' though presenting several outcrops on account of minor folding, may be grouped together as the Eastern District. But in the west, rocks of the same age occur over a wider extent of country. At

¹ Proc. Bristol Nat. Soc. ser. 4, vol. i (1905) p. 42.

² A. Vaughan, Q. J. G. S. vol. lxi (1905) p. 181.

³ Geol. Mag. 1902, p. 485; *ibid.* 1904, pp. 402, 585-87.

⁴ Recently, Dr. Hind has correlated some beds at Tenby with the Pendleside Series, Proc. Geol. Assoc. vol. xxi (1909) p. 179.

⁵ The Swansea sheet (N. S. 1-inch map 247) and 'The Country around Swansea' 1907; the Worms-Head sheet (N. S. 1-inch map 246) and 'West Gower & the Country around Pembrey' 1907. These memoirs will be referred to as the 'Swansea Memoir' and the 'West Gower Memoir' respectively.

⁶ West Gower Memoir, pp. 14-17.

present, those in the most widely separated outcrops are 4 or 5 miles apart across the strike, but they must have been deposited at a much greater distance from one another, seeing that the intervening ground is traversed by several deep folds. The northernmost outcrop, west of Llanrhidian, which is continuous along the strike with that at Mumbles, will therefore be distinguished, under the name of the North-Western District, from the southernmost, which lies between Port-Eynon Bay and Worms Head, and constitutes the South-Western District. No attempt, however, will be made to describe the whole outcrop of any part of the series.

Note.—(i) In the following descriptions the dolomites, unless otherwise described, are ‘contemporaneous,’ that is, they have been deposited as calcite or aragonite, but altered to dolomite by the waters of the Carboniferous sea,—shortly, therefore, after deposition. They are concluded to have had this origin on account of a variety of evidence, part of which has been published in the Swansea Memoir, pp. 13–20, pls. i & ii. ‘Vein-dolomites,’ due to a later mineralization, and described in the same memoir, are frequent, but have in some cases no obvious relationship with any visible vein, appearing, then, to have resulted from diffuse percolation. Such cases may occur in rocks which have been partly altered ‘contemporaneously.’ (ii) Thicknesses, unless stated to have been measured directly, have been estimated from outcrop, dip, and height; the locality is added in each case.

(1) Eastern District.

Lower Avonian.

K = *CLEISTOPORA* Zone.

Lithological characters.—So far as known, Lower Limestone Shales, that is, grey shales with subordinate fossiliferous limestones,¹ the latter weathering soft and ochreous. For Lower Limestone Shales underlie, probably at no great distance, the beds in Threecliff Bay referred to horizon β , and the position of horizon β is immediately above the *Cleistopora* Zone.

Limits.—Relations to beds above and below not seen.

Exposures.—Poor; near Penard Castle, and at Southgate.

¹ The ‘limestone flags,’ 200 feet thick, exposed along Penard Pill below Parkmill, and referred to this zone by Dr. Gubbin, Proc. Bristol Nat. Soc. ser. 4, vol. i (1905) pp. 52–53, belong (on the evidence of the fauna which we have obtained from them) to Z., together, probably, with horizon β . Their proximity to the Old Red Sandstone, mentioned by Dr. Gubbin, is due to a fault.

$Z = ZAPHRENTIS$ Zone.

Z_2 = subzone of *Zaphrentis konincki* Edw. & H., Carr.

Z_1 = subzone of *Spirifer clathratus* M'Coy.

Horizon β .

Lithological characters.—The lowest part of the Main Limestone¹; in descending order:—

Z .—Dolomites similar to the *Laminosa* Dolomites, that is, dark grey, finely crystalline, with small scattered nests of calcite and dolomite; interbedded with unaltered, thinly-bedded, crinoidal limestones, which are abundant in the lower part but die away to a great extent, upwards, at about 100 feet above the base of the Zone. Through the lowest 50 feet or so occur impersistently-tabular cherts and beekitized fossils.

Horizon β .—More thickly-bedded, crinoidal limestones; several yards exposed, but base not seen.

Fauna.—See pp. 544–45.

Thickness.—Of horizon $\beta + Z_1$:—about 150 feet (Threecliff Bay), on the assumption that the unexposed basal part of horizon β is not much more than 10 feet thick. Of Z_2 alone not determinable without difficulty; of the dolomite-group, $Z_2 + Laminosa$ Dolomites, 320 feet (Threecliff Bay).

Limits.—Horizon β , at the junction of the *Cleistopora* and *Zaphrentis* Zones, is probably also the junction of the Lower Limestone Shales and the Main Limestone. The junction of Z_1 and Z_2 is not marked by any appreciable lithological change; the top of Z_2 could not be localized, owing to the prevalent dolomitization, but evidently is also not defined lithologically.

Typical exposure.—Threecliff Bay, east side:—horizon β and the rest of the zone in sequence.

C_1 = Lower Subzone of the *SYRINGOTHYRIS* Zone.

2. *Caninia* Oolite.

1. *Laminosa* Dolomites.

The presence or absence of horizon γ in Eastern Gower could not be determined, owing to the dolomitization prevalent through Upper Z and Lower C_1 .

Lithological characters and thickness.—In descending order:—

2. *Caninia* Oolite. As a rule a remarkably pure limestone, distinguished from neighbouring horizons by its light-grey, often white colour and thick bedding, but especially by the purity

¹ This is merely a convenient name for the main limestone-mass of the Lower Carboniferous; no relationship to the Main Limestone of the North of England is implied.

of its oolitic structure¹—other material, such as crinoid-ossicles and smaller organisms, being largely confined to subordinate bands, while larger fossils amount to little more than a few beds of brachiopods or *Bellerophon* and an occasional *Syringopora*. Some bands, generally current-bedded, show evidence of contemporaneous erosion in included, sometimes rounded, fragments of oolite. Minute to microscopic ooliths are frequent, in places to the exclusion of others, especially towards the top. There, also, more often than elsewhere, the rock is dark in colour. Chert is unknown, unless represented by big, light-grey, concentrically-zoned siliceous nodules (Threecliff Bay); these, however, have not been critically compared with the siliceous aggregates found later in North-Western Gower. The rock passes up from the dolomite below, through intermediate phases in which the ooliths are less dolomitized than the matrix; and in places dolomite recurs in bands in the lower part.

Thickness.—160 feet (Threecliff Bay); 125 and 70 feet (Caswell Bay anticline); 150 feet (Longland Bay).

1. *Laminosa* Dolomites. Dark-grey or black, finely-crystalline dolomites,² chiefly representing fairly thin-bedded, crinoidal limestones, with here and there a few such beds still unaltered; towards the top, where the dolomite is rather paler and more thickly-bedded than below, it may be seen in places that it represents oolite. As a rule, dolomitization has obscured or obliterated most of the original constituents, including, among the fossils, some of the crinoid-ossicles, highly-resistant though they be. A feature of the crinoidal dolomites and others of the same type (as, for instance, those in Z) is the sporadic occurrence, throughout their mass, of small nests, up to several inches in diameter, of a coarsely-crystalline mosaic of glassy calcite and milky-white dolomite; on wave-worn surfaces these nests contrast strongly with the surrounding black rock. Many of them appear, from traces of organic structure still preserved, to have originated, partly through the alteration and recrystallization of calcareous mud, inside fossils with internal cavities, such as gasteropods and certain corals.

Thickness indeterminable (see Z₂).

Fauna.—Little known (p. 546), the *Laminosa* Dolomites having lost most of their fossils, and the *Caninia* Oolite never having had many.

¹ Under the microscope the grains, like those of most of the oolites of the Gower Avonian, show a radial-fibrous as well as a concentric structure, and differ thus from the S₂-pisolites. They are approximately spherical, with a diameter which in many is about 0.4 mm. and is seldom greater than 0.6 mm., but in the smallest measures 0.13 mm. or less.

² The completeness of the dolomitization may be gauged from the fact that an average sample, analysed for commercial purposes, was found to contain 20.35 per cent. MgO, the percentage in CaCO₃.MgCO₃ being 21.9.

Limits.—The boundary between the two groups is merely the somewhat uneven level at which the dolomitization characteristic of the lower shows greatest diminution when followed upwards. Unevenness of the level on a large scale would explain the inconstancy in thickness of the *Caninia* Oolite, should it be found that the total thickness of C_1 is fairly constant.

The top of the *Caninia* Oolite is sharply defined, and appears, from the character of fragments in the overlying bed, to have been slightly eroded, in places at least, prior to the deposition of C_2 , though it is not channelled or corroded as at West Williamston (Pembrokeshire).¹

Typical exposures.—Threecliff Bay, east side. Caswell Bay, east side:—several outcrops due to folding and faulting. This disturbance, which amounts to an anticline the crest of which has sagged down into a deep syncline with an overthrust from the north for its northern limb, has been described and figured by Dr. A. Strahan.² The ‘thick bed of light-coloured oolite’ referred to by him is the *Caninia* Oolite, and the ‘dark and thin-bedded limestones’ beneath are the *Laminosa* Dolomites together, possibly, with the top of Z_2 . Thus the *Caninia* Oolite has three complete outcrops and, in addition, a partial outcrop due to a strike-fault in the southern limb of the anticline. Longland Bay, north side:—the *Laminosa* Dolomites with, probably, the top of Z_2 ; in and near Rothers Tor, the *Caninia* Oolite.

Upper Avonian.

C_2 = Upper Subzone of the *SYRINGOTHYRIS* Zone.³

2. Standard limestones.

1. *Modiola* phase.⁴

Lithological characters.—Of the beds in descending order:—

2. Standard limestones, that is, limestones with a standard marine fauna, largely crinoids, corals, and brachiopods. The limestones are chiefly light-grey, rather thickly-bedded, and highly fossiliferous, but are in some cases dark and more thinly-bedded, especially those containing many gasteropods. Beds of dark, finely-crystalline dolomite and partings of shaly limestone occur, but are quite subordinate. In the gasteropod-beds the gasteropods are replaced in many cases by dolomite, even where the surrounding matrix has remained quite unaltered. Selective dolomitization of this kind is found only in organisms which have been originally aragonitic, such

¹ ‘The Country around Haverfordwest,’ Mem. Geol. Surv. (in the press).

² Swansea Memoir, p. 8, fig. 1.

³ As explained in § VI, the dividing-line between Lower and Upper Avonian is now taken at the base of C_2 .

⁴ The meaning of this term is explained on pp. 512 *et seqq.*

as gasteropods and orthocerates, but occurs at many Avonian horizons throughout the South-Western Province. In view of the instability of aragonite, this phenomenon is readily explicable, but implies that the organisms were still aragonitic when dolomitized. Further, the wide occurrence of the phenomenon, taken in conjunction with the short life of organic aragonite under the conditions accompanying vein-dolomitization, supports the conclusion, based on other considerations, that the dolomitization was 'contemporaneous.' Where there is no independent evidence of 'contemporaneous' dolomitization, it is found that organic aragonite has been replaced by coarse calcite-mosaic.

In places, the lowest bed contains fragments of the highest of the deposits of the underlying *Modiola* phase.

1. *Modiola* phase. A variable group of limestones, dolomites, breccias, and more or less calcareous shales or clay-mudstones, thinly bedded to finely laminated or papery; grey, dull buff, or greenish in colour (where free from Triassic reddening). The limestones, some of the argillaceous varieties of which are nodular and concretionary,¹ are either fine-grained to compact, or very finely oolitic (these resembling parts of the *Caninia* Oolite) or, occasionally, 'pisolitic.' Crinoidal limestones are rare, and are merely thin laminæ. Many of the fine or compact limestones and dolomites fracture conchoidally, and resemble 'chinastone-limestones'; but some are finely-laminated: they are calcite- or dolomite-mudstones (see p. 516). Either kind, but, it appears, more frequently the dolomitic, has recrystallized in places to calcite- or dolomite-mosaic of fine to coarse grain. Some finely-crystalline dolomites, however, may be 'contemporaneous' replacements of originally-calcitic rocks.

Ordinary detrital matter is frequent. The mudstones contain fine clayey material, which may exceed the calcareous, or, in some cases, fine quartz-sand. Breccias, of limestone and clay-mudstone fragments, measuring up to several inches in length, set in an argillaceous, calcareous, or fine-sandy matrix, may occur at any horizon. The fragments are similar in structure and fossils, either to some of the intercalated beds, or, in the case of some in the basement-bed, to the *Caninia* Oolite below, and evidently have been derived from both sources by contemporaneous erosion. Undoubtedly far-travelled material, other than clay and sand, appears to be absent, and the fragments show little, if any, rounding.

Some beds, however, have a brecciated structure, due in part to the penetration of an alien matrix into calcite-mudstone, apparently along irregular cracks. The origin of the structure

¹ The concretionary structure here mentioned is the type usual in argillaceous formations, not the peculiar structure of the 'concretionary beds' of the Avon.

is uncertain, but the mud appears to have been brecciated *in situ* by desiccation and cracking, without complete separation into fragments (see p. 514, footnote 6).

It may be noted that where a calcite-mudstone has succeeded a breccia with uneven upper surface, it has filled up the inequalities and formed a level surface; in the same way, but on a larger scale, a compact limestone, the lowest bed of a thin *Modiola* phase, fills up the inequalities in the upper surface of the conglomerate at Pendine (Carmarthenshire), described by Dr. Strahan.¹

True chert is unknown, but quartz in the peculiar form of minute, colourless and milky-white, zoned nodules of the size of sand-grains, has been noticed in a microscope-slide of one of the laminated limestones.

The following section illustrates the character of the phase and its relations to the beds above and below:—

Junction of C_1 and C_2 .—Southern outcrop on the east side of Caswell Bay.

	<i>Thickness in feet.</i>
9. Crinoidal limestones, resting sharply on	
8. Calcite-mudstone, white-skinned, filling inequalities in	
7. Breccia: consisting partly of black calcite-mudstone, some of which is fragmentary, and some, though penetrated by the matrix, apparently not completely broken up. The matrix is mudstone and very fine sandstone. With Bed 8.	5
6. Thinly-bedded, fine-grained limestones, including oolites. With Beds 4 & 5	11
5. 'Pisolite,' consisting of black, flattened, concentrically-laminated, calcitic 'pisoliths,' up to 0.75 inch in diameter, set in grey finely-crystalline dolomite; a few inches thick.	
4. Thinly-bedded to laminated, compact, argillaceous limestones.	
3. Dark limestone and black shale, passing into	1½
2. Clay-mudstone, with angular or subangular fragments, up to several inches in diameter, of fine-grained, dark oolite such as is found at the top of the <i>Caninia</i> Oolite; lenticular beds of black limestone in the upper part. Rests sharply on	1½
1. Fine-grained oolite, light or dark grey.	
Total thickness of <i>Modiola</i> phase	<u>18¾</u>

Bed 1 is the top of the *Caninia* Oolite (C_1); Beds 2–8 constitute the C_2 *Modiola* phase; and Bed 9 is the base of the standard limestones of C_2 .

The base of C_2 presents the features described above, which, in conjunction with the faunal peculiarities, show that it constitutes a *Modiola* phase, wherever it has been examined in the eastern district, except in Longland Bay. There it consists of a group, a few yards thick, of dark, finely-crystalline dolomites, similar to the *Laminosa* Dolomites but containing

¹ 'The Country around Carmarthen' Mem. Geol. Surv. 1909, p. 81.

abundant specimens of *Bellerophon*; this, though sharply separable, on account of its far more intense dolomitization, from the immediately succeeding beds (limestones with dolomitized *Bellerophon*), forms part of the standard limestones.

Fauna:—

2. Standard limestones.—Rich, especially in corals and brachiopods; crinoid-remains ubiquitous. In their fossils the lowest beds differ somewhat from the rest (pp. 542–43, 546–47).
1. *Modiola* phase.—Poor, consisting in most beds merely of fish-fragments, abundant ostracods, a *Spirorbis*-like annelid, or abundant *Calcisphæra* (?).¹ The calcareous alga (*Girvanella*) may be present, poorly preserved, in some coats of the ‘pisoliths.’ Standard forms—crinoid-ossicles and foraminifera—are known only from rare laminae, in which, however, they are abundant.

Thickness.—Total, between 200 and 250 feet (Threecliff Bay); 350 feet (Longland Bay). The discrepancy does not appear to be due to faulting. Of the *Modiola* phase alone, 16 to 20 feet (direct measurement).

Limits.—The change at the top of the *Modiola* phase to standard limestones is abrupt, and is accompanied in places by contemporaneous erosion, which has been sufficient locally, as may be seen in Caswell Bay, to cut out some of the highest beds of the phase. The top of the subzone, that is, C-S, the junction of C₂ and S₁, is not marked by any appreciable lithological change. For this reason, it is frequently useful to group the subzones C₂ and S₁ as one division, C₂ + S₁.

Typical exposures:—

2. Standard limestones.—Threecliff Bay, east side. Caswell Bay,—several exposures, the southernmost on the east side being the best. Longland Bay, east side.
1. *Modiola* phase.—Threecliff Bay, east side, and Caswell Bay,—several complete exposures at each place, the display in the syncline being particularly good. Note:—The outcrops of the *Modiola* phase are easily found by reference to the underlying *Caninia* Oolite.

S = *SEMINULA* Zone.

S₂ = subzone of *Productus* ‘*cora*’ D’Orb., mut. S₂.

S₁ = subzone of *Caninia bristolensis* Vaughan.

The subzones were not delimited, nor examined so closely as in the South-Western District; but the sequence was seen to be much the same as in that district.

¹ The organisms provisionally referred, here and on subsequent pages, to *Calcisphæra* are isolated spheres of calcite, with an average diameter of 0·05 to 0·1 mm., the walls of which are thin and have a radial-fibrous structure.

Lithological characters.—In descending order:—

S_2 :—2. *Modiola* phase. Chiefly 'pisolites,'¹ oolites, foraminiferal limestones, and calcite-mudstones, the latter including the unfossiliferous white-weathering variety called, from their porcellanous appearance, 'chinastone-limestones'; fossils abundant (individuals, not species), notably *Seminula* in beds with a dark, fine-grained matrix; some beds dolomitized. Contemporaneous erosion is evidenced in some of the oolites by irregular fragments, also of oolite, measuring up to 2 feet in length; in such beds dolomite, if present, is more abundant in the matrix than in the fragments.

1. Standard limestones. Oolites, with some pisolites and fine-grained limestones; in many, few macroscopic fossils, but others are coral-² brachiopod-limestones, etc. Contemporaneous erosion is evidenced in the same way as in the overlying group. Dolomites recur at intervals.

S_1 :—Limestones, largely made up of crinoids, corals, brachiopods, and, in some dark impure beds, gasteropods; dolomitic beds at intervals; chert-nodules and beekitized fossils, extending, with the gasteropod-beds, possibly into the base of S_2 as in the South-Western District.

Fauna.—See pp. 547-48. The *Modiola* phase has not yielded any of the peculiar forms generally present in such phases.

Thickness.—Of the whole zone, about 1250 feet (Longland Bay).

The subdivisions were not separately measured, but S_2 was by far thicker than S_1 .

Limits.—The upper limit of S_1 was not examined. The two parts of S_2 , though separable in a broad way, pass insensibly one into the other, the incoming of the pisolites abundant in the upper part being quite gradual. The upper limit of S_2 is marked by a rapid and considerable lithological and faunal change, thus:—

Junction of S_2 and D_1 .—Generalized from several exposures.

D_1 : Pseudobreccias of limestone or dolomite, with a typical D_1 -fauna from the base upwards; in places a thin basal oolite.

Passage-beds: Current-bedded, fine-grained, and in part oolitic limestones with *Cyathophyllum*; 15 to 20 feet.

S_2 : *Modiola* phase. Limestones of the types previously mentioned; *Seminula ficoides* in crowds, at intervals up to the top.

¹ Much of the 'pisolite' might more properly be termed 'contemporaneous breccia,' for it consists largely of more or less irregular masses, of various sizes, of dense, amorphous-looking limestone, some only of which show a pisolitic structure. These latter consist of a layer of dense calcite, of fairly uniform thickness, around an organic nucleus, generally a *Seminula* (double-valved) or a coral-fragment, but they rarely, if ever, have a concentric structure of several well-defined layers, though such a structure is frequent in the S_2 -pisolites of the Avon. Under the microscope it is seen that the dense material may contain *Girvanella*-like tubules, but also that much of it is structureless and of exceedingly fine grain, being devoid, apparently, of any necessary connexion with these tubules.

² None of the coral-limestones are unbedded through great thicknesses, and thus differ from coral-reef limestones.

Typical exposures.—Threecliff Bay, east side. Pwll-du Bay. Whiteshell Point:—lower part of the zone. Longland Bay, east side:—the whole zone, in sequence with those above and below. Limeslade and Broadslade (Bracelet) Bays:—much of S_2 , the junction with D_1 at Tutt and at Mumbles Head (in both the Middle Head islet and the adjacent cliff). Bishopston Valley:—several exposures, including one of the same junction at the Sker.

$D = DIBUNOPHYLLUM$ Zone.

The development of the upper part of this zone in Gower differs in important respects from that in the Bristol and Dublin districts, and has therefore been subdivided on somewhat different lines. In Eastern Gower the following groups are present. Above the lower part of the zone, which is a normal development of D_1 , comes a group of limestones, connected faunally with both D_1 and D_2 , and, therefore, referred to as D_{1-2} . This forms the top of the Main Limestone, and is followed by a calcareo-argillaceous group, the 'Upper Limestone Shales' of the Geological Survey,¹ which probably represents part of D_2 , but certainly extends to a higher horizon and is therefore known as D_{2-3} . A resemblance of some of its fossils to forms which are common in the *Zaphrentis* Zone led at one time, before the D_3 -fauna of County Dublin was known, to some controversy as to the systematic position of D_{2-3} . It is now clear, however, that, as Dr. Wheelton Hind maintained,² it occurs at the top of the Viséan.

It is succeeded by an almost purely argillaceous group, which is included in the Millstone Grit by the Geological Survey and has been recognized by Dr. Hind as a representative of the Pendleside Series. And, though it is possible that the Pendleside Series is a phasal equivalent of part of the sequence of coral-and-brachiopod zones of some regions, the name *Posidonomya* Zone (P), already in use for it, will be retained.

$D_1 = \theta\phi$ Subzone.

Lithological characters.—Chiefly light-grey, thickly-bedded limestones, many partly dolomitized. The limestones are predominantly of the type, peculiar in Gower to the D-zone, characterized by the remarkable 'pseudobrecciated' structure which has been described by Mr. Tiddeman³ and forms the subject of a separate note (p. 507). Besides much obviously subsequent

¹ The disadvantages attaching to other names, such as 'Bishopston Beds,' which have been applied to this group, have been stated by Dr. Strahan in the Swansea Memoir, p. 21. It may be added that, as delimited by Mr. E. B. Wethered & Prof. C. Lloyd-Morgan, the 'Upper Limestone Shales' of the Avon,—for which Buckland & Conybeare originally proposed the name,—include lower horizons than those of Gower.

² Geol. Mag. dec. 5, vol. i (1904) pp. 585–87.

Swansea Memoir, p. 10. He observes the persistence and high horizon of these rocks, and groups them together as the 'Mumbles Head Beds.'

dolomite, there are occasional large irregular bodies of dolomite, similar in its coarse grain and its association with clay to the dolomite characteristic of pseudobreccias, but evidently of somewhat different origin. Another peculiar feature of the pseudobreccias, noted by Mr. Tiddeman and difficult of explanation, is the occurrence of numerous pits, as a rule circular, 18 to 30 inches in diameter and a foot deep, but also irregular and larger, distributed evenly over the upper surface of some of the beds (see Pl. XXXVIII, fig. 1, taken from an example in the North-Western District); these pits are filled with grey, buff-weathering clay or, less frequently, with coarse, light dolomite, and, at some levels at least, underlie thin beds of clay continuous with their infillings. A few cases have been observed of a pseudobreccia passing laterally, in a short distance, into clay with limestone-rubble.

Subordinate to the pseudobreccias are limestones of ordinary types. In places a coarse-grained oolite marks the base of the zone, and other limestones, dark or light, are intercalated at higher horizons. The zone is, however, further distinguished by the presence of thin coaly layers, also noticed in the first place by Mr. Tiddeman (*loc. cit.*). Some, though thin and valueless like the others, resemble many true coal-seams in their character and association with underclay, and have probably originated in the same way. Thus, an unusually good exposure at Colts Hill, of one lying at a considerable distance from both the top and the bottom of D_1 , showed in descending order:—

[Section near thin coal in D_1 .—Southern quarry at Colts Hill.]

Thickness in feet.

4. Limestone pseudobreccia, with a standard marine fauna.	
3. Coal smut and finely-laminated, carbonaceous shale; in all about	$1\frac{1}{2}$
2. Underclay with frequent rootlets and much pyrite; passing upwards into shale (3).....	$3\frac{1}{2}$
1. Limestones, pseudobrecciated or ordinary. The upper surface of the highest is soft, but uneven (with projecting corals), and it is uncertain whether it shows a passage or a non-sequence into (2).	

There is no need to postulate a different origin for the smut-bed at Oxwich Point, though found by Mr. Tiddeman to contain many small gasteropods.

Fauna.—See pp. 548–49.

Thickness.—Between 400 and 450 feet (Pwll-du Bay).

Limits.—The faunal change to D_{1-2} is not accompanied by any noticeable lithological change.

Typical exposures.—Pwll-du Bay and Head. Mumbles Head (including the Middle Head and the lighthouse islets):—much of the subzone, the dolomitic pseudobreccias conspicuous. North-north-eastern face of Mumbles Hill:—dolomitic pseudobreccias, in several cases ‘pitted.’ Oystermouth (Colts Hill southern quarry):—

pseudobreccias, unweathered, contrasting strongly in appearance with the weathered rocks on the coast. Bishopston Valley :— the outcrops of the D Zone in the sides of the valley are too narrow to accommodate its probable thickness, the cause of the discrepancy being, doubtless, unexposed strike-faulting.

D₁₋₂.

Lithological characters.—Much as in D₁, but the pseudo-breccias disappear in the upper part. The following section shows the sequence at Colts Hill, and also the relations to the beds (D₂₋₃) above :—

D₁₋₂ and D₂₋₃.—Colts Hill.

	<i>Thickness in feet.</i>
D ₂₋₃ :—‘Black lias,’—thinly-bedded, black, argillaceous limestones and shales, one of the latter a foot thick; cherts.	
D ₁₋₂ :—3. Thickly-bedded, fine-grained limestones including oolite; the highest few feet dark, with black cherts, the rest light grey and chertless; pure, except for some dolomite and occasional shaly partings; about	50
2. Dark, shaly nodular limestone, slightly dolomitized, the nodules elongated	1-2
1. Thickly-bedded, light-grey limestones, including pseudobreccias, continuous with those of D ₁ ; base not determined owing to poorness of exposure, but thickness probably about	60

Fauna.—See pp. 549 & 551. The *Productus* Band at Colts Hill occurs at the top of Group 1. *Caninia cornucopie*, mut. D₂₋₃ was observed in the dark limestone immediately below D₂₋₃.

Thickness.—Uncertain, but not more than 120 feet (Colts Hill).

Limit.—The change to D₂₋₃, which is considerable as regards both fauna and rock-types, is almost abrupt, and is emphasized by weathering, as may be seen at Colts Hill: D₁₋₂ remaining hard limestone, but D₂₋₃ becoming soft, clayey ‘rottenstones.’ The change is foreshadowed in the highest few feet only of D₁₋₂; there the fauna includes one of the corals abundant in D₂₋₃, while the rock is dark and contains dark cherts.

Exposures.—Pwll-du Head: top of cliff 300 to 400 yards south of High Penard. Colts Hill northern quarry and an old face between this and the southern quarry.

D₂₋₃.

[The ‘Upper Limestone Shales’ of the Geological Survey.]

Lithological characters.—Limestones and shales with cherts, the whole known locally as ‘black lias.’ The limestones are thin—generally less than 3 feet thick—and evenly bedded (fig. 3, p. 492); almost invariably dark-grey or black, with a white skin or crust of impurities where weathered; crinoid-débris

Fig. 3.— D_{2-3} limestones and shales in the 'Black Lias' Quarry at Oystermouth.

T. C. Hall photo. (Geol. Surv. Colln.).

and fine-grained matrix in varying proportions. They are argillaceous and some, at least, are dolomitized—the rhombohedra being, as a rule, minute. The shale-bands and partings are more or less calcareous and, though abundant, generally less than 2 feet thick. The chert, largely made up of sponge-spicules, is abundant in some beds, either as black flinty nodules, more or less sharply defined from the matrix, or diffused and difficult to detect except under the microscope; it also occurs as beekite, replacing fossils, in some beds apparently in this form alone.

‘Rottenstones’: relations to the ‘Black Lias.’—The ‘black lias’ characters are retained only where the rocks are unweathered. In places the beds, both limestones and shales, have been completely altered by decalcification and oxidation to light-grey or yellow-brown, so-called ‘rottenstones’¹ and clays. The ‘rottenstones’ are either friable and largely argillaceous, or brittle, like porous earthenware, and partly siliceous—the latter evidently representing diffusely-cherty limestones. Many are cavernous with moulds of unsilicified fossils, and some display in sharp relief many details, hidden in unweathered material, of the structure of silicified fossils. Wavellite² occurs in cavities and joint-planes in the higher ‘rottenstones,’ as in the overlying radiolarian cherts.

The ‘rottenstones’ and clays differ so materially at first sight from ‘black lias,’ that their equivalence is masked; but the view put forward by Mr. Tiddeman,³ that they have been derived by process of ordinary weathering from such beds, is confirmed by the relations of the two types exposed in the northern quarry at Colts Hill. There, D₁₋₂, which is carried by the dip from top to bottom of the quarry, is followed by ‘rottenstones’ and clays with cherts near the top, but by ‘black lias’ with the same fauna at the bottom. Other palæontological evidence, also, points to the equivalence of the two types: the fauna of the Bishopston ‘rottenstones’ being approximately the same as that of the Oystermouth ‘black lias’ (p. 551).

An explanation of the extensive character of the weathering which the ‘rottenstones’ have undergone is probably to be sought in the fact that the latter, so far as known, are situated as at Bishopston, that is, at or near the top of the partly dissected plateau which constitutes the whole of this part of Gower; whereas ‘black lias’ is found at lower levels, as at Oystermouth. It is believed that the conditions of minimal run-off and denudation which characterize the top of the plateau have lasted long enough to allow the rocks there to be completely weathered to a considerable depth. In the valleys, where denudation is rapid, the outcrops at lower levels are unweathered.

¹ None of the beds have been worked for rottenstone; they are not so fine-grained as the material of that name dug for polishing-powder on the ‘North Crop.’

² First noticed by Logan: see *Mem. Geol. Surv.* vol. i (1846) p. 134.

³ Swansea Memoir, pp. 25–26.

Fauna.—Abundant (p. 550).

Thickness.—At Oystermouth at least 175 feet, and possibly as much as 350 feet, if, as is probable, the effects, in duplicating outcrop, of the strike-fault shown on the Geological Survey map between Oystermouth Castle and Colts Hill are negligible.

Limit.—The junction with the Pendleside Series, P, is nowhere clearly exposed. D_{2-3} and P are seen together in the Bishopston road-cutting, but their relations cannot be determined with certainty, as both are weathered and crumbling and their steeply-dipping junction strikes along, and largely beneath, the road. The section suffices to show, however, that D_{2-3} with its abundant fauna is succeeded by a comparatively barren radiolarian-chert group (the base of P), the change, which is great and far-reaching both faunally and lithologically, being abrupt, if not marked by an unconformity (see p. 529). Of unconformity, however, there is no evidence, stratigraphical or palaeontological, although the junction contrasts strongly, in its abruptness, with the gradual passage by alternation at Loughshinny.¹

Typical exposures.—Oystermouth:—The Black Lias quarry (fig. 3, p. 492) affords an excellent section of a thickness of 86 feet of beds in the condition of 'black lias' from near the top of D_{2-3} downwards; the sequence is without notable lithological or faunal change, and only one limestone, the thickest (3 feet), is sufficiently pure to make lime. In Colts Hill northern quarry the lowest beds only are exposed. 'Black lias' also forms the intervening ridge on which Oystermouth Castle stands.

Bishopston:—In the cutting—chiefly along its south side—on the Clyne Common-Bishopston Valley road, the top of D_{2-3} , to a thickness of at least 45 feet,² is exposed in the condition of 'rottenstones.' The sequence is obscured by weathering and foundering, but has yielded a rich fauna, largely as moulds.

P = *Posidonomya* Zone.

This zone is represented by the Pendleside Series, first recognized in Gower by Dr. Wheelton Hind.³ He correlated, however, with this series not only the beds now under discussion, but also the D_{2-3} 'rottenstones.' The composite group, thus constituted, had previously been called 'Bishopston Beds' by De la Beche⁴ and 'Gower Shales or Series' by Phillips.⁵ As regards the beds above D_{2-3} , Dr. Hind's correlation has been confirmed⁶ by the subsequent

¹ C. A. Matley & A. Vaughan, Q. J. G. S. vol. lxiv (1908) p. 413.

² Details are recorded by Mr. Tiddeman, Swansea Memoir, pp. 22, 24.

³ Geol. Mag. 1902, p. 485; *ibid.* 1904, pp. 402, 585-87.

⁴ Mem. Geol. Surv. vol. i (1846) pp. 133-34.

⁵ 'Manual of Geology' 1855, pp. 169-70.

⁶ Of the fauna quoted by him, *Glyphioceras diadema* (Beyrich) alone may be regarded as of zonal value; and confirmation is, therefore, important.

discovery in them of *Glyphioceras spirale* (Phill.)¹ and *Posidonomya becheri* Bronn (p. 552). The importance of distinguishing between the 'rottenstones' and the overlying beds has been recognized by the officers of the Geological Survey, who have included the latter in the Millstone Grit, the 'rottenstones' (as previously mentioned) being 'Upper Limestone Shales.'

Lithological sequence.—In descending order:—

2. Dark, well-laminated, non-calcareous shales, with some iron-stone-nodules and thin sandstones; in the lower part with black cherts²; according to De la Beche,³ there is a thin limestone (4 inches) in the middle. Very thick.
1. Radiolarian cherts⁴ with interbedded shales, the cherts showing the fine dark and light stripes, due to thin lamination, frequent in radiolarian deposits, and the shales, also, well-laminated and non-calcareous, but soft and in part weathering to a chocolate colour. Under the microscope it is seen that the cherts are exceedingly fine-grained, and that the dark and light laminæ differ one from the other in grain and in the relative proportions of their organic and inorganic constituents, the latter chiefly the finest quartz-silt. Many of the laminæ are lenticular, some of them sharply so. Wavellite occurs along joints. The total thickness is slight, the group forming, as it were, basal beds to the shales above.

Apparently, the series includes no recurrence of the fossiliferous limestones characteristic of the underlying Avonian.

Fauna.—2. Largely unfossiliferous, but some beds are crowded with thin-shelled goniatites⁵ preserved as casts.

1. Of the abundant radiolaria and subordinate sponge-spicules in the cherts, the former correspond closely with Culm forms.⁶ The shales are largely unfossiliferous, but some have yielded a few lamellibranchs (p. 551) and indeterminate plant-remains. Thickness.—Unknown, though probably at least 1500 feet, as *Glyphioceras spirale* (Phill.) has been found in abundance at apparently that distance above the base. Of this amount probably much less than 50 feet belongs to the radiolarian-chert group.

Limits.—The radiolarian cherts, doubtless, pass up into Group 2; but the junction does not appear to be satisfactorily exposed, and no repetition of similar cherts above is known. The upper

¹ By Mr. J. Pringle of the Geological Survey, Swansea Memoir, pp. 25 & 30; the specimens, 'J. P. 1260, 1265-1276' in the Survey collection, were found in a stream-bank on Barland Common, 150 yards west of Old Castle.

² Swansea Memoir, p. 25.

³ Mem. Geol. Surv. vol. i (1846) p. 133.

⁴ Dr. Strahan has described in the Swansea Memoir, pp. 22-25, how the discovery by Dr. G. J. Hinde of the radiolarian origin of these cherts resulted from De la Beche's comparison of the beds with the cherts of Coddan Hill. He appends Dr. Hinde's description of the microscopic characters of the rocks and their organisms.

⁵ Swansea Memoir, pp. 25 & 30.

Dr. G. J. Hinde in the Swansea Memoir, p. 25.

limit of Group 2 has not been determined, but is not conspicuous lithologically, as the succeeding beds, up to the Coal Measures, are, also, largely shales.

Where examined.—Bishopston, north side of road-cutting:—the radiolarian cherts. Barland Common:—the shales (Group 2); the most complete section is now overgrown, but was described by De la Beche¹ and has been discussed by Dr. Strahan.²

(2) North-Western District.

The sequence of the Avonian rocks in North-Western Gower is similar on the whole to that in Eastern Gower, and attention will, therefore, be directed chiefly to the differences between the two districts. The *Cleistopora* Zone, however, is poorly exposed in the Eastern district, and is described in some detail.

Most of the zones were examined along Burry Pill, the banks of which, north and north-west of Cheriton, afford a fairly complete sequence from a horizon low in the *Zaphrentis* Zone up to the middle of D₁, and much of D₁ was, also, found well displayed in Tor-gro, a range of cliffs between Burry Pill and Landimore. The thicknesses, estimated from outcrop and dip, in these localities are stated in Table I (facing p. 505).

Cleistopora Zone,³ K.—This zone, although not completely exposed, especially towards the top, probably corresponds exactly with the Lower Limestone Shales, the resemblance to the development in those parts of the South-Western Province where this relationship holds good being close. It consists in descending order of:—

			Thickness in feet.
Probably K ₂ .	4. Grey shales, with thin limestones containing crinoids and brachiopods; upper part not seen; total thickness probably between 250 feet and		300
K ₁ or K ₂ .	3. Limestones, including fine-grained dark oolite; at least		30
	2. Like 4 on the whole; with Km		150
Probably K ₁ .	1. <i>Modiola</i> Phase, Km. Grey shales with ostracodal and oolitic limestones and a red hematitic limestone of <i>a</i> -type; quartzitic sandstone with modioliform lamellibranchs near the base.		
Total			<u>between 400 & 500</u>

The characters and thicknesses of Groups 1–3 and of 200 feet of Group 4 were determined in the banks of a dingle, north-east of Fairyhill House; the total thickness of the zone was estimated from its outcrop at Cheriton.

The base of the zone, as in most parts of the South-Western

¹ Mem. Geol. Surv. vol. i (1846) pp. 133–34.

² Swansea Memoir, pp. 22 & 30; also fig. 2 (p. 23).

³ The field-observations were made conjointly with my colleague, Mr. T. C. Cantrill. I gladly take this opportunity of thanking him for encouragement and advice at many stages throughout this work.

Province, is well-defined lithologically and, in a sense, palæontologically also, the unfossiliferous, 'continental' Upper Old Red Sandstone giving place, rapidly but conformably, to the fossiliferous marine Avonian. The junction, as exposed in the Fairryhill dingle and recorded, with details of the adjacent beds, in the West Gower Memoir (p. 8), is as follows. The top of the Upper Old Red Sandstone is a barren, red marl with limestone-nodules ('race').¹ The base of the Avonian is a grey, sandy limestone with limestone-fragments, some of which are red like the 'race' in the underlying marl, the rest being, doubtless, practically contemporaneous. In the matrix between the fragments are a few ostracods and ooliths, and the bed is followed immediately by the 'sandstone with modioliform lamellibranchs' previously mentioned. The junction of the Old Red with the Avonian is so sharp as to be clearly traceable in a microscope-slide [E. 5288, Geol. Surv. Colln.]; and though it partly owes its conspicuousness to the difference in colour between the beds above and below, this difference is but the outward sign of a fundamental change in the conditions of deposition, marine organisms appearing in the rocks immediately above. In Gower, it may be noted, the change (which is inferred to be conformable from a merging of the top of the Old Red into the Avonian) is almost abrupt, and has been accompanied by current-action, as is evident from the fragments at the junction; in some parts of the South-Western Province, on the contrary, as for instance in Pembrokeshire,² marine and continental conditions alternated for a space.

The junction of Km with Group 2 of K₁ and that of K₂ with Z do not appear to be exposed. The limestone-group (3) probably separates K₁ from K₂, for it is exactly similar, lithologically, to the limestone that affords the best boundary between these two sub-zones in Pembrokeshire.

Zaphrentis Zone, Z.—Cherts are found, though only now and then, above the group of limestones with cherts near the base of the zone.

Syringothyris Zone, C.—The *Caninia* Oolite is partly dolomitized at various horizons up to the top, the matrix between the ooliths being the material chiefly affected. In places, the rock has been partly silicified, the silica taking the form of numerous aggregates of microscopic quartz-crystals; unlike the dolomite, these replace ooliths and stop at the interstitial matrix.

The top of the Oolite is uneven, the irregularities reaching a depth of a foot or more. But the hollows are not in all cases simple depressions; some extend into the Oolite laterally and are overhung by roughly-horizontal tongues of it measuring up to a foot

¹ The red marl, to a thickness of 4 feet, rests upon unfossiliferous, pebbly, grey quartzite which is probably the top of the thick quartz-conglomerates that form the upper part of the Upper Old Red of Gower (West Gower Memoir, p. 5).

² 'Summary of Progress for 1908' Mem. Geol. Surv. 1909, pp. 35-36.

in length. Somewhat similar irregularities are known elsewhere in shallow-water deposits, and have been caused by distortion of originally evenly-bedded strata while still plastic, shortly after deposition. It is possible, therefore, that the *Caninia* Oolite, also, has been drawn out tangentially while in this condition, though by what agency is not apparent; in this case, however, it is equally possible that the irregularities are due to some slight subaërial corrosion prior to the deposition of C_2 .

The basal layers of the *Modiola* phase of C_2 , which follow the *Caninia* Oolite sharply and fill the hollows just described, are puckered in a fashion suggestive of parts of landscape-marble. At higher levels in the phase conglomeratic beds are rare in this district; and dolomite-mudstones appear to be restricted, as in parts of Pembrokeshire, to a buff-weathering, argillaceous group at the top. The fauna of the phase is devoid of standard forms, and a *Mitcheldeania*-like alga¹ (in minute fragments) is present; otherwise the fossils are those found in the Eastern District.

The standard limestones of C_2 differ considerably in character, though not in fauna—except in an absence or rarity of *Caninia*—from the same subzone in the Eastern District. The chief difference lies in an absence of contemporaneous dolomite in North-Western Gower, even the gasteropod-valves, which would be readily affected by ‘contemporaneous’ dolomitization, being wholly calcitic; further, the beds are more uniform in character, rather darker and less thickly-bedded than in the Eastern District. They thus approach the facies of the *Seminula* Zone of the outcrop north of the coalfield.

Note.—Along the east side of Burry Pill the zone has been extensively vein-dolomitized.

Seminula Zone, S.—The lower subzone, S_1 , is similar on the whole to the underlying standard limestones of C_2 , but includes, though rarely, thin calcite- or dolomite-mudstones and, possibly, a little dolomite that has ‘contemporaneously’ replaced limestone; chert is absent. The top of the subzone was not seen.

In the S_2 -oolites, as in the *Caninia* Oolite, some ooliths have been replaced by aggregates of microscopic quartz-crystals; but dolomite is wanting. In the *Modiola* phase which forms the upper part of the subzone, mudstones, including ‘chinastone-limestones’ and nodular beds, appear to be better developed than in the Eastern District, and dolomite is wanting. The change to the standard conditions of Zone D at the close of the phase was rapid.

Dibunophyllum Zone, D.—The chief divergences in this part of the sequence from the development in Eastern Gower are (1) an absence of dolomite, as in C_2 and most of S; (2) the presence of a thick oolite at the base of D; and (3) the presence of D_2 .

(1) The absence of dolomite is most noticeable where the material is known to be particularly susceptible to dolomitization; such

¹ By this expression is meant a calcareous alga with a minute, radially-columnar structure; such identification must suffice, until the revision of the genera promised by Prof. E. J. Garwood is published.

material is presented both by the valves of gasteropods and by the unrecrystallized part of pseudobreccias. The latter, with typical structure (see Pl. XXXVIII, fig. 2) and abundant foraminifera, are as widespread as in the Eastern District, but, like the gasteropod-valves, are in all cases devoid of dolomite. In connexion with pseudobreccias, it may be noted that one of the best exposures of the 'pitted' bedding-planes (p. 490) that characterize some of them may be observed in Tor-gro, Landimore. One such bedding-plane, dipping at a steep angle, which has been described by Dr. Strahan in the West Gower memoir, p. 12, is shown in Pl. XXXVIII, fig. 1. The incoming of *Cyrtina septosa* (Phill.), which varies in its horizontal distribution, is worth noting; in this district it first becomes abundant near the base of the pseudobreccias.

(2) The D₁-oolite, which is about 50 feet thick, is coarse-grained, like the band in the same position in Pembrokeshire. The lowest part is darker, more thinly-bedded and less oolitic than the rest; otherwise the contrast with the *Modiola* phase at the top of S₂, immediately below, is complete.

(3) D₂ has been recognized at Llanrhidian. In the top of the Main Limestone the highest bed exposed in the neighbourhood has yielded the following fauna¹:—

Choneti-Productus.

Papilionaceous *Chonetes* or *Daviesiella*.

Martinia glabra (Mart.).

Martinia lineata (Mart.).

Productus corrugatus M'Coy.

Productus edelburgensis-latissimus.

Spirifer bisulcatus Sow.

The rock is a dark and bituminous, crinoidal limestone, rich in brachiopods; unfortunately it is seen to a thickness of only a few feet (in low crags 260 yards east of the church). It is underlain by light-grey, nodular limestones, probably pseudobreccias, which have yielded merely giganteid *Productus* and may belong either to D₁ or to D₂. Downwards in the sequence other beds are not seen for some distance, and upwards nothing is known until the radiolarian cherts² at the base of P are reached: it is uncertain, therefore, whether D₂₋₃ is present in this district.

(3) South-Western District.

The structural features of this district, which lies between Rhossili and Port-Eynon Bays, are shown and described in the Geological Survey map and memoir.³ Here we need only mention that, owing to a syncline in the promontory south-west of Rhossili, much of the sequence Z-C₁ is found along both the north-western and the southern sides of the promontory, and that it is repeated yet again near the Inner Head by an anticline ranging west-north-west through the gut between the mainland and the Worms-Head islets. The western extremity of the Worms-Head islets was not examined.

¹ Identified by Dr. Vaughan and preserved in the Geological Survey Collection (E. D. 891-906).

² Poorly exposed in a lane a furlong east of the crags of D₂-limestone.

³ One-inch sheet 246 (Worms Head); West Gower Memoir, pp. 3, 12-16.

As in the case of North-Western Gower, attention will be paid chiefly to the differences between this district and those previously described.

Cleistopora Zone, K.—Probably equivalent to the Lower Limestone Shales. All the groups seen in North-Western Gower appear to be present, and in the following account are referred to under the same numbers as those on p. 496.

1. *Modiola* phase, Km. This group includes a 5-foot band of conglomerate (exposed north-west of Rhossili rocket-house) of red-brown quartz-pebbles, measuring up to half an inch in diameter, with larger pebbles of fossiliferous mudstone or shale, the whole in a matrix of calcareous sandstone which encloses, also, reddened and rounded crinoid-ossicles like those in limestones of *a*-type. The conglomerate is both overlain and underlain by ostracod-shales with limestones and sandstones, but probably lies near their base; the junction of those below with the underlying Upper Old Red Sandstone is not exposed. The thin limestones interbedded with the shales are chiefly of a type usual at this horizon, that is, most of them contain a mixed fauna of crinoids, brachiopods, ostracods, lamellibranchs, small *Bellerophon*-like gasteropods and *Orthoceras*, besides fragments of contemporaneous sediments, including phosphates.

2. As in Pembrokeshire, the standard part of K₁ commences with a group of quartzitic sandstones (Rhossili), and the parallelism is maintained in the shales immediately below Group 3, which, as exposed at the north-west foot of Hardings Down, are distinguished by the markedly nodular character of their interbedded limestones.

3. As in North-Western Gower (and Pembrokeshire); exposed at Hardings Down.

4. Not seen.

Zaphrentis Zone,¹ Z.—In South-Western Gower, Z consists largely of limestones, dolomites, though frequent, being subordinate. Shaly bands are found, but at considerable intervals, through much of the lower part of the zone: and cherts, with beekitized fossils, have a considerable upward range (beekite being abundant in the lowest beds—some part of Z₁—exposed in the anticline at Worms Head,² and cherts, with beekite, occurring for several hundred feet above).

As usual in this zone, the limestones are largely crinoidal and rather thinly-bedded, and the dolomites dark grey and finely crystalline. The fauna is tabulated on pp. 544–45.

¹ First recognized in this district by Dr. Gubbin & Dr. Vaughan, Proc. Bristol Nat. Soc. ser. 4, vol. i (1905) pp. 50–52, 54.

² The lowest beds in this anticline have been referred by Dr. Gubbin & Dr. Vaughan (*op. cit.* pp. 50 & 54) to Z₂, on insufficient evidence. Their horizon is probably lower than that of the lowest beds at Rhossili, rather than above as stated by those authors, and it is agreed that the Rhossili beds (*op. cit.* pp. 51 & 54) belong to Z₁. The beds at both places were once considered by myself to lie at the base of the zone (West Gower Memoir, p. 15), but a re-examination of the district, though leaving the age as Z₁ unquestioned, has thrown doubt on their basal position, especially as regards the beds at Rhossili.

The boundary between Z_1 and Z_2 was not located, but is evidently not marked by any striking lithological change; the top of Z_2 , also, is not conspicuous, as nodular and argillaceous beds, though most characteristic of Horizon γ above, commence in the highest part of Z_2 .

The most complete and accessible section lies between the core of the Worms-Head anticline, at a level somewhere in Z_1 (see preceding footnote), and the edge of the cliff at the Coastguard Station to the north-east, which exposes the top of the zone and a few yards of the overlying γ beds. Here at least 530 feet (a figure obtained largely by direct measurement) is exposed, but in this district the full thickness of the zone cannot be determined.

Horizon γ and *Syringothyris* Zone, C.—The chief points to be noticed in this part of the sequence are:—

- (1) Horizon γ is well developed;
- (2) The *Laminosa* Dolomites are but partly dolomitized, and contain cherts and beekitized fossils;
- (3) There is no *Modiola* phase at the base of C_2 ;
- (4) The facies of C_2 resembles that of the standard C_2 -limestones of the Eastern District, not that of the North-Western District.

(1) Horizon γ consists of thinly bedded, nodular limestones with soft argillaceous partings; this character is most marked in the upper part. Fossils, especially corals, abound; they are magnificently displayed on extensive, weathered surfaces at Tears Point—the best Avonian collecting-ground in Gower, but unfortunately discovered too late for its fauna to be recorded.

The base of Horizon γ , defined as the level at which large Caninids first become frequent, may be placed, to facilitate recognition, at a rather marked argillaceous bed which reaches the sea at Tears Point. The summit was placed, arbitrarily, but merely for the purpose of measurement, at the lowest of the dolomites which are the feature of the overlying *Laminosa* Dolomites, for in the absence, so far as could be seen, of *Chonetes* cf. *comoides* (Sow.) from the *Laminosa* Dolomites, there did not appear to be any important faunal difference between this group and Horizon γ .

(2) The lower part of the *Laminosa* Dolomites (for thickness, see the table on p. 502) consists of about equal proportions of dolomites of the usual character and of unaltered limestones; the limestones are rich in corals, and contain also some crinoid-cups. The upper part is normal in its dolomitization, but includes occasional chert-nodules and beekitized fossils. The change at the top to the *Caninia* Oolite is rapid, that is, the thickness of dolomitized oolite at the junction is small, and so too in North-Western Gower.

(3) Standard C_2 limestones follow the *Caninia* Oolite immediately. Their basal bed is firmly welded to the somewhat uneven top of the latter, and contains small masses of oolite (the outcome, doubtless, of a little contemporaneous erosion), which are mixed with broken brachiopods, corals, etc., proper to C_2 .

(4) The C_2 limestones are largely light-grey and thickly-bedded, and are less crinoidal than many parts of Z ; probably fewer dolomites and gasteropod-beds are intercalated than in C_2 of

Eastern Gower. The subzone passes up insensibly into the similar limestones which form the lower part of S_1 .

In the *Caninia* Oolite, it may be observed in passing, there are thick bands of fine-grained, non-oolitic material, some of which is foraminiferal, some comminuted-crinoidal; corals are more frequent near the top of the Oolite than at lower levels, those noticed being *Michelinia*, *Syringopora*, and a *Clisiophyllid*.

The thicknesses of the subdivisions of Horizon γ and C_1 vary somewhat, as may be seen from the accompanying table. The thickness of C_2 was obtained, with some uncertainty on account of faulting, only between Fall and Mewslade Bay; it probably lies between 280 and 350 feet.

Thicknesses in feet of Horizon γ and Subzone C_1 in South-Western Gower.

	γ	<i>Laminosa</i> Dolomites.			<i>Caninia</i> Oolite.	Total, $\gamma + C_1$.
		Lower part.	Upper part.	Total.		
Fall	120	80	60	140	160 (at least)	420
Inner Head...	135	115	90	205	130	470

The best exposures of Horizon γ and C are (1) the Inner Head islet, accessible for a limited time from the mainland by crossing the rocky isthmus at low tide:—the whole, except the top of C_2 ; (2) the cliff-top near Old Castle, Rhossili, where γ has yielded the fauna enumerated on p. 545, and the *Laminosa* Dolomites also are seen; (3) Tears Point and the cove called 'Fall' on the north:— γ and C_1 ; (4) the fine range of cliffs¹ between Fall and the bluff west of Overton Mere:— C_2 with much of S .

Seminula Zone,² S .—This zone has almost the same facies as in the Eastern District. The top appears, however, to be so lacking in rock-types characteristic of *Modiola* phases, that it is separated from the rest of S_2 , not on its own merits, but for purposes of comparison with the *Modiola* phase on this horizon in other areas.

The general sequence, which has been determined more exactly than in Eastern Gower, is as follows, in descending order:—

S_2 :—

3. *Modiola* phase. Evidenced merely by occasional 'pisolites.' The 'pisoliths' outwardly resemble those which are so abundant in the Eastern District, but include fragments of a *Mitcheldeania*-like alga. True calcite-mudstones, such as 'chinastone-limestones,' if

¹ Described by Dr. A. Strahan in the West Gower Memoir, pp. 12-13.

First recognized in this district by Dr. Gubbin & Dr. Vaughan, Proc. Bristol Nat. Soc. ser. 4, vol. i (1905) pp. 44-50, 54-56.

present, are rare; their place is, apparently, taken by limestones that are rather less fine-grained and contain an abundance of *Calcisphaera* (?) and foraminifera. These rocks and the 'pisolites' are interbedded with oolite like that of Group 2 below, their lower limit being unknown but probably quite indefinite. Upwards the change to Zone D with its characteristic fauna and sediments is comparatively rapid, the highest bed definitely assignable to S_2 being separated from the lowest undoubted D limestone—the oolite described on p. 504—by 24 feet of limestones of less determinate relations.

2. Chiefly light-grey, thickly-bedded oolites, many sparingly fossiliferous; some fine-grained limestones, composed of current-bedded organic detritus. At intervals occur dark, finely-crystalline dolomites, up to several feet thick, which, though in some cases markedly impersistent along the bedding, as already described by Dr. Strahan (West Gower Memoir, p. 13), are inferred, from their microscopic characters,¹ to owe their alteration to the influence of the Carboniferous sea. Near the base of the group the limestones contain sporadic, light-grey sponge-cherts, of concentric, alternately siliceous and calcareous zones. Thickness, including the *Modiola* phase above, between 820 and 870 feet.

1. Gasteropod-limestones,² dark, highly bituminous, and well-bedded; many of the gasteropods selectively dolomitized in the way described on p. 484; cherts and beekitized fossils. Thickness = 120 feet.

S_1 :—

2. Gasteropod-limestones, dark and continuous with those above, but on the whole more thinly-bedded, and separated from one another by soft shaly partings; highly fossiliferous (p. 547); beekite in fossils in the upper beds; some beds more or less dolomitized. Thickness = 105 feet.

1. Limestones, similar to the underlying C_2 limestones; less fossiliferous than the beds above; little dolomite and no chert or shale. Thickness = 115 feet.

The whole zone is exposed in the cliffs and shore near Port-Eynon Point, from the bluffs at the cliff-foot, west of Overton Mere,³ where S_1 is finely displayed, eastwards to Sedgers Bank, where D comes in. S_1 and S_2 enter largely into the cliffs on the west, also, of Overton Mere, as far as Mewslade Bay.

¹ Similar to those of contemporaneous dolomites described in the Swansea Memoir, pp. 16–20 & pl. i, fig. 4.

² Included by Dr. Gubbin in S_1 ; the absence of *Caninia bristolensis* Vaughan has led to my placing this group in S_2 .

³ Details from this locality, especially of the gasteropod-beds, are recorded by Dr. Gubbin (Proc. Bristol Nat. Soc. ser. 4, vol. i, 1905, pp. 44–50), who has, however, greatly underestimated the thickness of S_2 ; the 'black calcareous sandstone' to which he refers on p. 47 is the dolomite in Group 2 of S_2 , mentioned above.

Dibunophyllum Zone,¹ D.—The important feature of this zone in South-Western Gower is a strong development of D_2 , a subzone which is unknown in Eastern Gower though present in the North-Western District.

The facies of D_1 differs in no important respect from that observed in Eastern Gower. An oolite, 7 feet thick, in part of the coarse type usual at this horizon, lies at or near the base. Some of the beds, including some pseudobreccias,² are dolomitized, the gastropods showing the selective action previously mentioned.

In D_2 are placed the highest beds of the Main Limestone. The change of fauna in passing from D_1 to D_2 is not accompanied by any lithological change, so far as known, but pseudobreccias die out some distance above. The upper part of D_2 consists of various rock-types, including coarse oolite, coral-limestone, dark fine-grained bituminous limestone with black chert, dolomite, and pale limestone. Some of the latter occurs immediately below D_{2-3} , but is as pure as any other part of the Main Limestone, except for an abundance of pale chert. The change to D_{2-3} , faunal and lithological, is as rapid as in Eastern Gower.

D_{2-3} is equivalent to the Upper Limestone Shales, which are similar to the beds of the same name in Eastern Gower, except that they include, interbedded among the argillaceous limestones and shales, a few pure, pale limestones with chert, resembling some of the D_2 limestones. The correlation with D_{2-3} of Eastern Gower is discussed on pp. 551–52.

The top of the zone and its relations to the overlying P beds are not revealed.

The whole of D_1 and D_2 , 600 feet in all, and a thickness of at least 150 feet of D_{2-3} , the latter folded, and much reddened by Triassic infiltration, crop out on the shore of Port-Eynon Bay. Better exposures of much of D_1 and D_2 , with the faunas recorded on pp. 548–49, are obtained in quarries in the plateau above Port-Eynon Point; those in D_2 lie about a quarter of a mile south-south-west of Port-Eynon church.

Posidonomya Zone, P.—The development in this district is probably the same as in other parts of Gower. Radiolarian cherts are believed by Dr. Strahan³ to crop out near Overton; and shales, grey or raddled, with the fauna enumerated on p. 552, are exposed in an old 'paint-mine' at Port Eynon described by the same author (*loc. cit.*). Black, evenly-bedded sponge-cherts do not appear to be exposed in the mine *in situ*; but, to judge by fragments in the overlying Triassic conglomerate, they do occur in the series. A single band, a few inches thick, of shale studded with crinoid-ossicles and now decalcified, emphasizes the rarity of such material above the *Dibunophyllum* Zone.

¹ First recognized in this district by Dr. Gubbin & Dr. Vaughan, Proc. Bristol Nat. Soc. ser. 4, vol. i (1905) pp. 43–44 & 56.

² These I described in the West Gower Memoir, p. 16, as 'without dolomite,' whereas some only are undolomitized.

³ West Gower Memoir, p. 18.

TABLE I.—SYNOPSIS OF THE LITHOLOGICAL SEQUENCE IN DIFFERENT PARTS OF GOWER.

Zone, Subzone or Horizon.		North-Western District.		Eastern District.		South-Western District.	
Posidonomya Zone. P.		As in the Eastern district, apparently.	Unknown.	2. Chiefly dark; non-calcareous shales; some black sponge-cherts in the lower part. 1. Radiolarian cherts with shales; wavellite.	Thickness. At least 1500 feet. Probably less than 50 feet.	As in the Eastern district, apparently.	Unknown.
Dibunophyllum Zone. D.	D ₂₋₃	Not known.	Probably less than 300 feet.	'Black lias':—soft-weathering, argillaceous limestones and shales, with sponge-cherts and some dolomite; wavellite near the top.	Between 175 & 350 feet.	Much as in the Eastern district, but with some intercalated, non-argillaceous, cherty limestones.	At least 150 feet.
	D ₂ or D ₁₋₂	D ₂ :—Limestones.		D ₁₋₂ :—Limestones, including pseudobreccias; dolomite.	At most 120 feet.	D ₂ :—Coral-, oolitic, and other limestones; pseudobreccias; cherts; dolomite.	About 200 feet.
	D ₁	Much as in the Eastern district, but without dolomite.		Chiefly pseudobreccias, generally dolomitic; 'pitted' surfaces; thin coal with underclay.	Between 400 & 450 feet.	Much as in the Eastern district (coal not known).	About 400 feet.
Seminula Zone. S.	S ₂	2. <i>Modiola</i> Phase:—Calcite-mudstones; 'pisolites'; standard limestones. 1. Chiefly light-coloured oolites; no dolomite known.	At most 800 feet.	<i>Modiola</i> Phase:—Calcite-mudstones; 'pisolites'; standard limestones; dolomite.	About 1250 feet.	3. <i>Modiola</i> Phase:—Poor; a few 'pisolites' among standard limestones; dolomite. 1 & 2. Chiefly light oolite with zoned sponge-cherts near its base; some cherty gasteropod-limestones below; dolomite.	About 1000 feet.
	S ₁	Rather dark limestones, with a rare calcite- or dolomite-mudstone or 'contemporaneous' dolomite; no chert.		The development of S up to the <i>Modiola</i> Phase is much as in the South-Western district.		2. Gasteropod-limestones; shaly partings and a little chert; dolomite. 1. Thickly-bedded, pale limestones; little dolomite.	220 feet.
Syringothyrus Zone. C.	C ₂	2. Similar to S ₁ , but without mudstone or dolomite. 1. <i>Modiola</i> Phase:—Much as in the Eastern district.	Between 230 and 280 feet.	2. Much as in the South-Western district.	200 to 350 feet.	Similar to group 1 of S ₁ . (No <i>Modiola</i> Phase developed at the base.)	Probably between 280 & 350 feet.
	C ₁	2. <i>Caninia</i> Oolite. 1. <i>Laminosa</i> Dolomites.	45 feet.	1. <i>Modiola</i> Phase (wanting at Longland Bay):—Chiefly calcite-mudstones; dolomite.	16 to 20 feet.		
	γ	Not recognized.	100 feet.	2. <i>Caninia</i> Oolite. 1. <i>Laminosa</i> Dolomites; few undolomitized (crinoidal) limestones. Not recognized.	70 to 160 feet.	2. <i>Caninia</i> Oolite. 1. <i>Laminosa</i> Dolomites, the lower part half undolomitized; occasional cherts. Chiefly nodular, argillaceous limestones.	130 to 160 feet. 140 to 200 feet. 120 to 135 feet.
Zaphrentis Zone. Z.	Z ₂	Similar to the <i>Laminosa</i> Dolomites.	About 520 feet.		320 feet.	Limestones with dolomites; cherts in the lower part, to a higher horizon than elsewhere in Gower.	At least 530 feet.
	Z ₁ β	Much as in the Eastern district. Not recognized.		Similar to the <i>Laminosa</i> Dolomites. Dolomites, with crinoidal limestones in the lower part; cherts towards the base. Crinoidal limestones.	150 feet.	Not recognized.	
Cleisopora Zone. K.		Lower Limestone Shales:—2-4. Grey shales with thin limestones; a thick oolitic limestone, probably between K ₁ and K ₂ . 1. <i>Modiola</i> Phase, Km. Grey shales with ostracod-limestones; limestone of α-type. (Passage)	Between 400 and 500 feet.	Probably the Lower Limestone Shales, which are barely exposed.	Unknown.	Lower Limestone Shales:—2-4. Little exposed, but probably as in North-Western Gower. 1. <i>Modiola</i> Phase, Km. Chiefly grey ostracod-shales; a quartz-conglomerate.	Unknown. Unknown.
Devonian.		Upper		Old		Red Sandstone.	

(4) Tabulated Synopses of the Sequence.

Table I embodies the chief features of the developments in the three districts of Gower, together with the thicknesses, which are also shown graphically in fig. 4 (p. 506). In both table and figure the Eastern District is placed between the two others because, as will be seen, it has relations with both. The differences between the different developments, of which more will be said later, might not be regarded as of importance, were it not that each development could be matched elsewhere in the South-Western Province.

Table II embodies the features common to much or all of the area, and may be regarded as typical of much of the South-Western Province.

TABLE II.—SYNOPSIS OF THE LITHOLOGICAL SEQUENCE IN GOWER AS A WHOLE (GENERALIZED FROM TABLE I).

Zone or Subzone.	
P.	2. Chiefly dark non-calcareous shales. 1. Radiolarian cherts.
D. D ₂₋₃ ¹ D ₂ or D ₁₋₂ D	'Black lias':—soft-weathering, argillaceous limestones with shales and cherts. Limestones, including pseudobreccias. Chiefly pseudobreccias; thin coal ² or coals.
S. S ₂	2. <i>Modiola</i> phase, characterized especially by 'pisolites.' 1. Chiefly oolites.
C. C ₂ +S ₁	2. Limestones of various types. 1. <i>Modiola</i> phase, ³ characterized especially by abundant calcite- and dolomite-mudstones.
C. C ₁	2. <i>Caninia</i> Oolite. 1. <i>Laminosa</i> Dolomites (replacements of crinoidal limestones).
Z.	Crinoidal dolomites and limestones; cherts near the base.
K. ⁴	2. Grey shales with thin limestones; a thick oolitic limestone, probably between K ₁ & K ₂ . 1. <i>Modiola</i> phase, Km, characterized especially by limestones of <i>a</i> type.
Devonian.	Upper Old Red Sandstone.

Note:—K is equivalent, probably exactly, to the Lower Limestone Shales; the sequence, Z to D₂ inclusive, is termed 'Main Limestone' by the Geological Survey; D₂₋₃ is termed Upper Limestone Shales; and P is included with the Millstone Grit.

¹ Not known in the North-Western District.

² Not known in the South-Western District.

³ Not developed in the South-Western District.

⁴ K is best known in the North-Western District, but there is no reason to believe that the development is different elsewhere.

[506]

III. NOTE ON THE PSEUDOBRECCIAS OF THE D-ZONE
[E. E. L. D.].

The following observations relate to the pseudobreccias of Gower, where these rocks are peculiar to the *Dibunophyllum* Zone and at that level are extensively developed. While to some extent supplementing Mr. Tiddeman's description,¹ this note deals more particularly with the question of origin, which is of interest on account of the resemblance of the rocks to true breccias, and also because other nodular, foraminiferal limestones, such as some of the bands in the Chalk, may owe their structure to the same cause. It may be mentioned that the explanation of the structure of pseudobreccias, here advanced, has suggested itself to Mr. J. A. Howe also, in the course of a comprehensive study that he is making of these and other limestones.

The structure in question, the outward appearance of which may be gathered from Pl. XXXVIII, fig. 2, appears at first sight to be due to the incorporation of angular fragments of rather dark limestone, of various sizes up to a diameter of several inches, in a ground-mass which is limestone in some places, and dolomite in others, but is always lighter in colour and more argillaceous than the 'fragments,'² sometimes markedly so. The greatest contrast is presented where the 'ground-mass' is dolomite, as the latter forms a rather coarsely-crystalline, pale-grey mosaic.³ We may first, however, consider the undolomitized examples.

'Fragments' and 'ground-mass' vary in their relative proportions, but as a rule are approximately equal, and the latter occurs characteristically as meandrine tracts enclosing the former. In no pseudobreccia has it been possible to distinguish bands differing one from the other in the size or abundance of their included 'fragments.' The same fauna, a standard marine assemblage, is found in the 'fragments' as in the 'ground-mass.'

Microscopic examination shows that the matrix of the fossils in both 'fragments' and 'ground-mass' is a consolidated 'mud,' consisting essentially of finely-divided calcite and containing, as a rule, a notable quantity of foraminifera.⁴ Furthermore, such examination shows that the outlines of the 'fragments' are not sharp and well-defined like those of true fragments, but that their matrix shades off into the matrix of the 'ground-mass.' On this account largely, and also in view of other considerations which will be adduced, it is inferred that the rocks are not

¹ Swansea Memoir, p. 10.

² See analysis by Mr. E. G. Radley, Swansea Memoir, p. 13.

³ A dolomitized example is figured in the Swansea Memoir, pl. i, fig. 1.

⁴ The following have been identified in microscope-sections by Dr. R. L. Sherlock:—'Callencroft Quarry, Mumbles [E 5000]: very rich in foraminifera, chiefly Endothyra, such as *Endothyra bowmani* Phillips, *Endothyra* sp., and *Textularia* sp. (either *gibbosa* d'Orb. or *eximia* Eichw.).—Oxwich [E 5001]: *Endothyra* sp. (small form).'

really fragmental. Ultimately, doubtless, they owe their structure in part to their composition,¹ but the proximate cause of it alone concerns us here.

Before passing to this subject, however, it should be remarked that Dr. Hume connects the nodular structure of parts of the Chalk with current-action.² A complete comparison of nodular chalks with pseudobreccias has not yet been made, but it is known that some of the former closely resemble the latter. It is possible that in the clay with limestone-rubble, into which occasional pseudobreccias pass laterally (p. 490), the rubble, which resembles the 'fragments' of pseudobreccias proper, but is sharply separable from its clay-groundmass, has been more or less rearranged by currents, after originating in a patchy recrystallization (and segregation) of calcareous mud in the manner presently to be described. There is no reason, however, to believe that the 'fragments' in the pseudobreccias proper have also been washed up and redeposited; and it remains to be seen whether all nodular chalks show evidence of current-action, or whether some are not pseudobreccias.

As contradicting the view that the 'fragments' in the Gower pseudobreccias are derived, and have been incorporated in their present condition, may be adduced the fact that occasionally some are continuous one with the other and partly enclose 'ground-mass,' thus losing their resemblance to fragments. It may be suggested, however, that the 'fragments' in these rocks are true fragments of contemporaneous sediments, which, unre-crystallized at the time of their incorporation, have lost all trace of their limiting surfaces through later recrystallization of themselves and some of the surrounding ground-mass. Apart from the absence from the pseudobreccias of any sign of current-action in either a sorting or a banding of their constituents, no evidence of a truly fragmental origin has been detected in any of the numerous 'fragments' that have been examined, and the suggestion takes no account of the fact that, so far as known, the pseudobrecciated structure is confined to foraminiferal limestones.

To turn to the proximate cause of the pseudobrecciated structure, the features whereby the matrix in the 'fragments' differs from that in the 'ground-mass' are, in addition to its less argillaceous character, mentioned above: (1) that it is invariably rather more crystalline, though still fine-grained and obviously a slightly recrystallized mud; and (2) that it is often, possibly always, more foraminiferal. Both features are illustrated by Pl. XXXIX, in which, indeed, hardly any foraminifera are seen in the dense 'ground-mass.' (The 'fragments' are the light areas with irregular boundaries along the top, bottom and left-hand margins of the field.)

¹ Rocks similar in structure to the pseudobreccias of Gower are known in other regions, in the same and in other formations, but all appear to be foraminiferal. Many foraminiferal limestones, however, are not pseudobreccias, whence it is obvious that other conditions, besides a suitable composition, have been necessary for the production of the distinctive structure.

² Proc. Geol. Assoc. vol. xiii (1894-95) p. 230.

Confining our attention at present to the first feature, it is concluded that a muddy matrix has undergone partial recrystallization, as it were into irregular clots, and thus, by differentiation, quite possibly from a homogeneous state, given rise both to the 'fragments'—the recrystallized parts, and the 'ground-mass'—the less crystalline residue. This local recrystallization is inferred, for several reasons, to have taken place shortly after the deposition of the mud. Its peculiar patchiness, in which it differs strongly from late recrystallizations, and its widespread occurrence on the same horizon (D)¹ are most readily explicable on this view. Probably the best reason, however, is that, in Gower, as we shall see, many pseudobreccias have been dolomitized after recrystallization—but while still within the influence of the Avonian sea²; such a view postulates an early age for the recrystallization. The latter, therefore, appears to be analogous to the early recrystallization known to affect recent calcareous muds,³ but why it assumed a patchy character is unknown.

The other features of the pseudobreccias would result directly from a patchy and early recrystallization of a calcareous mud. Recrystallization implies deposition of calcite, and would probably be accompanied by segregation of that mineral and, in a still soft matrix, extrusion of impurity into the unrecrystallized parts. This action would account for the excess of argillaceous matter in the 'ground-mass.' Furthermore, delicate calcareous bodies like foraminifera would tend to disappear from the unrecrystallized parts, to balance, by their dissolution, the deposition of calcite in the recrystallizing areas.

But, whatever be the ultimate nature of the 'fragments,' their more coarsely crystalline structure in comparison with the 'ground-mass' has given rise to secondary features of much interest. They appear darker in hand-specimens, on account of their greater translucency. Again,—and this is of far more importance,—they offer a greater resistance to chemical change than the 'ground-mass,' in accordance with the general rule that in calcitic rocks such resistance is proportional, *cæteris paribus*, to the coarseness of the crystalline structure. The difference is shown in a variety of ways. The 'fragments,' which may be almost invisible on a fresh surface, often project boldly on weathered faces. Occasionally, it is found that the 'ground-mass' has recrystallized at some late date completely to coarse calcite-mosaic, but the 'fragments' included in this mosaic are still unaltered. The difference, however, between the two kinds of material is shown most conspicuously by the difference in their susceptibility to

¹ Pseudobreccias are frequent in D, not only in Gower but also in many other areas both within and without the South-Western Province.

² This view as to the age of the dolomitization is based not only on the original evidence cited in the Swansea Memoir, p. 14, but also on various facts which have come to light since.

³ See, for instance, E. W. Skeats, Bull. Mus. Comp. Zool. (Harvard) vol. xlii (1903) pp. 105–18, figs. 2 & 4–6.

dolomitization by the waters of the Avonian sea. (Dolomitization due to clearly-later percolation, it may be noted in passing, has affected both materials equally.) In fact, the difference between the 'ground-mass' and the 'fragments' in this respect is so great as to suggest that at the period of dolomitization the former was still unconsolidated and readily permeable, as well as more unstable. In many pseudobreccias and for long distances in them it has been completely altered, though, as a rule, the whole of the enclosed

Fig. 5.—Diagram showing the relation of the dolomite, in a partly dolomitized pseudobreccia, to unrecrystallized calcite (see p. 511). Natural size.

[*a, a* represent 'fragments,' consisting of recrystallized calcite; *b, b*, tracts of 'ground-mass,' consisting of unrecrystallized calcite; and *c, c*, dolomite. The limit of the dolomite is generally sharp against the 'fragments,' but against the unrecrystallized calcite it is too indefinite, as a rule, to be shown properly in a diagram.]

'fragments' have been but little affected. It has thus, by determining the paths of dolomitization, led to a marked accentuation of the pseudobrecciated structure. In many dolomitized pseudobreccias, as in the example figured in the Swansea Memoir, pl. i, fig. 1, the 'ground-mass' has been so completely replaced by dolomite-mosaic in which its proper structures have been lost, that the original nature of the material replaced by the dolomite would be open to doubt, were it not that in other such pseudobreccias dolomitization has been incomplete, and wholly absent from parts

of others again. The latter, one of which is represented diagrammatically by fig. 5 (p. 510), show clearly that the paths followed by the dolomitization have been the tracts of unrecrystallized 'ground-mass.' Pl. XXXIX, taken from the junction of a dolomitized with an undolomitized portion of a pseudobreccia, shows that the dolomite has developed freely in the dense 'ground-mass,' partly as isolated rhombohedra, partly as a band of clustering crystals, but that it has been excluded from the more crystalline 'fragments.' (The diagonal cross seen in many of the rhombohedra is probably due to inclusions of the argillaceous impurity in the 'ground-mass.')

Dolomitized pseudobreccias, such as that figured in the Swansea Memoir, resemble true breccias more closely than do the unaltered rocks, owing to the great contrast in them between the sharply-defined 'fragments' of dark, fine-grained limestone and the enveloping pale-grey, rather coarsely-crystalline dolomite. Also, their structure is more conspicuous, whether on unweathered faces, or in cliffs, where they often become cavernous through the dissolution of their 'fragments.' Consequently in Gower, where they abound along the southern coast, they have attracted more attention; but, as we have now seen, their dolomitization has not been the primary cause of their peculiar structure—it has merely brought into prominence an earlier recrystallization.

IV. NOTE ON LAGOON-PHASES AND THE ORIGIN OF RADIOLARIAN CHERTS [E. E. L. D.].

(1) Introduction and Definition.

By a lagoon-phase is meant a group of rocks, the characters and development of which show that it has been deposited in a coastal area, of wide extent both parallel and at right-angles to the coast, but so extremely shallow as to have been, in effect, isolated from the neighbouring though deeper parts of the sea, and thus to have become the site of peculiar types of sediment and fauna.¹

A lagoon-area as thus defined is shown in section, at right angles

¹ It has been thought better to adopt the word 'lagoon' and to speak of such an area as that defined above as a lagoon-area, rather than to coin a new expression, because the conditions obtaining in lagoons, as defined in the 'Century Dictionary' (London, 1899), must approximate closely to those which it is desired to connote. According to this authority, a lagoon is primarily 'an area of shallow water, or even of marshy land, bordering on the sea, and usually separated from the region of deeper water outside by a belt of sand or of sand-dunes, more or less changeable in position.' Though there is no evidence that subaërial barriers, such as dunes or coral-reefs, have existed in the case of the lagoon-areas with which we shall deal, possibly because the chances are against their having been preserved, exposed, and correctly interpreted, they are evidently not essential features of lagoons. Further, several formations which are lagoon-phases as above defined, such as the Solenhofen Slate, are described by Prof. J. Walther and others as having been deposited in shallow lagoons.

Fig. 6.—*Ideal section across a lagoon-area, at right angles to the coast.*
[The slopes throughout, both terrestrial and submarine, are exaggerated.]

to the trend of the coast, in fig. 6. Lagoon-phases are regarded as coastal deposits, and the lagoon-area, consequently, is shown in the figure as a coastal shelf, for reasons which appear later (pp. 525-28). The area is represented as bordered by rapidly deepening water, to accord with the general profile of recent coastal shelves.

That extremely-shallow marine areas, such as are here postulated, have formerly existed is beyond question, for the deposits in some cases may be interpreted without hesitation. Under the names of '*Modiola* and *Posidonomya* phases' some of these deposits, of Avonian age and calcareous facies, have already been described by Dr. Vaughan,¹ and the object of this note is, partly to bring out the distinctive characters of these phases, partly to extend the concept to rocks of cherty facies. Lagoon-phases, therefore, would include both *Modiola* and *Posidonomya* phases and equally shallow deposits of other ages and facies.

But, although lagoon-areas have undoubtedly existed, it is probable that their extreme shallowness and wide extent together would not sufficiently account for the general absence from them of strong currents, which, as we shall see, has been their distinguishing feature. Apparently the rise and fall of the tide, which, if considerable, would inhibit lagoon-conditions on an open coast, has been either so small as to be negligible, or has been rendered inoperative by subaerial barriers, which may have included thickets of vegetation such as those of mangrove-swamps, along the margins bordering deeper water.

¹ These are tabulated by him in Table II of his report on the faunal succession in the Avonian, Rep. Brit. Assoc. (Winnipeg) 1909, p. 190; reference will presently be made to his description of the development of these phases in the Avon section.

(2) The *Modiola* Phases—Calcareous Lagoon-Phases—
of Gower.

Introduction.—The Carboniferous succession in Gower includes four lagoon-phases, namely, Km at the base of K, the base of ($C_2 + S_1$),¹ the top of S_2 , and the radiolarian cherts at the base of P. The chief features of each of the first three, which are calcareous in facies, are set out in Tables III & V (pp. 514 & 526), the deposits characteristic of shallow waters being distinguished from intercalated ordinary sediments where, as is generally the case, the latter exist. The fourth phase, which differs from the others in being cherty, will be discussed separately, as an origin at a totally different depth has been ascribed to rocks elsewhere possessing closely similar characters. All four phases have been recognized at other places in the South-Western Province and some of them elsewhere also; the information in the tables concerning the Bristol area has been extracted from Dr. Vaughan's accounts,² and that about Pembrokeshire has been obtained during the progress of the Geological Survey.³ No attempt, however, will be made to give full particulars of the geographical extent of any phase.

Distinctive features of the *Modiola* phases.—We may now learn from Table III in what way *Modiola* phases differ from ordinary marine formations; from the interpretation of their distinctive features, which will follow, it will appear why they may be styled 'lagoon-phases.'

On the one hand, all the Gower *Modiola* phases exhibit, at one place or another, but not necessarily everywhere, unmistakable shallow-water characters which are shown to some extent, also, by ordinary shallow-water deposits (Table III, 1 & 2). In the first and third phases there are, likewise, many intercalations of ordinary rock-types (Table III, 4), due to the interruptions of lagoon-conditions by more normal ones mentioned on p. 518.

On the other hand, they all include, also, one or more rock-types (Table III, 3) which, in a fairly shallow-water marine series, such as the Avonian, are confined either to *Modiola* phases or to merely occasional beds in the shallow-water parts of the sequence. Some of these distinctive rock-types (namely, limestones of a type and purple and green shales) appear to point directly to deposition in shallow, isolated, coastal areas, and, therefore, will be considered first; the other group to be discussed here—mudstones and shales characterized by an exceedingly fine grain,—which is far more important than the first group on account of its more general occurrence, resembles, rather, deep-sea deposits.

¹ By this expression is meant the group consisting of subzones C_2 and S_1 .

² Q. J. G. S. vol. lxi (1905) p. 181 (cited thus: 'Bristol paper,' p. ...); and, in the case of the Avon section, Proc. Bristol Nat. Soc. ser. 4, vol. i (1906) pp. 74-168 (cited thus: 'Avon,' p. ...).

³ 'The Country around Haverfordwest' and 'The Country around Pembroke' (Mems. Geol. Surv.); in preparation.

TABLE III.—THE *MODIOLA* PHASES—CALCAREOUS LAGOON-PHASES—REPRESENTED IN GOWER.

	Km, the <i>Modiola</i> Phase at the base of K.	The <i>Modiola</i> Phase at the base of ($C_2 + S_1$).	The <i>Modiola</i> Phase at the top of S_2 .
1. Shallow-water characters: faunal and floral.	Some beds yield little, except—lamellibranchs, such as <i>Modiola</i> and <i>Sanguinolites</i> ? <i>Spirorbis</i> ?; abundant ostracods, such as <i>Cytherella</i> . ¹	Almost all the beds are unfossiliferous or yield only—lamellibranchs, e.g. <i>Sanguinolites</i> ?; <i>Spirorbis</i> -like annelids; abundant ostracods, e.g. <i>Cytherella</i> ; <i>Mitchellianella</i> or similar algae (Forest of Dean; North-Western Gower, etc.).	Some beds, such as the 'chinalstone-limestones', are unfossiliferous, or yield only <i>Spirorbis</i> -like annelids. Beds with many fragments of a <i>Mitchellianella</i> -like alga (South-Western Gower).
2. Shallow-water characters: lithological. ²	Some beds show a brecciation in which the fragments are but slightly separated (Avon, p. 100). Beds with fragments of contemporaneous sediments. Presence of pure oolite of very fine grain (Pembroke-shire, etc.). Packing of ostracod-valves within one another by long-continued drifting. ³	Some beds show a brecciation in which the fragments are but slightly separated (Gower, etc.). Beds with fragments of contemporaneous sediments. Presence of pure oolites of very fine grain. Presence of 'pisolite' (Eastern Gower).	Beds with fragments of contemporaneous sediments. Presence of 'pisolites' (Gower, etc.). Presence of pure oolites.
3. Deposits distinguishing the <i>Modiola</i> phases from other shallow-water marine formations.	Calcite-mudstones, slabby to nodular ('Avon', pp. 100, 103; etc.). Limestones of a type. Purple and green shales (Pembroke-shire).	Abundant calcite-mudstones and dolomite-mudstones; homogeneous (for instance, 'chinalstone-limestones') to finely-laminated; a little of the latter in North-Western Gower crumpled as in landscape-marble.	Calcite-mudstones, chiefly homogeneous ('chinalstone-limestones'). Limestones with the structure of landscape-marble ('Avon', pp. 91-93; Sodbury ⁵).
4. Marine sediments of standard types included within the <i>Modiola</i> phases.	Always present, so far as known; consisting of: (1) intercalated limestones and sandstones with an abundant brachiopod-and-crinoid fauna,—frequent. (2) fine conglomerate (South-Western Gower, etc.).	Very rare; for example, a thin limestone-parting with crinoids at Caswell Bay (Eastern Gower).	Always present, so far as known; consisting of:—(1) intercalated limestones with abundant corals and brachiopods (often of few species),—abundant. (2) grit-bands (Sodbury ⁵).

Note.—Features of general occurrence are not followed by the name of a locality.

1 This fauna in its entirety is known only from the Avon; at most localities some members are missing.
2 Some characters found in the *Modiola* phases, though to be expected in shallow-water deposits, may occur through a considerable range of depth. They include:—the presence of plant-fragments (in the first and second phases); of contemporaneous dolomites (in the second and third phases); and of quartz-sand (in the matrix of breccias in the second phase in Eastern Gower).
3 First noticed in Purbeck limestone, and explained, by Mr. F. Chapman, Proc. Geol. Assoc. vol. xix (1905-6) p. 284 & pl. v. fig. 2.
4 The complete fauna is known only near Pembroke Dock; at most localities some members are missing.
5 A. Vaughan, Bristol paper, p. 200.
6 The cause of this brecciation *in situ* (p. 486) has not been determined, but that of an apparently phenomenon in certain beds of *Modiola*-phase type in the Scottish Carboniferous is, Mr. R. G. Carruthers tells me, sun-cracking and consequent flaking; and it may be added that cracks, vertical or irregular, having in section the appearance of sun-cracks and filled with the matrix of the overlying limestone, have been found in a calcite-mudstone in the phase at the base of C_2 in Gower.

Limestones of *a* type.—Limestones of *a* type, named after the well-known Bryozoa Bed, Horizon α ,¹ of the Avon, consist largely of crinoid-ossicles and dendroid bryozoa, broken and rounded by rolling and reddened with hæmatite. Hæmatite may appear, also, in the matrix and in the coatings of ooliths, but the characteristic feature of the rocks is the relation of the iron-oxide to the fossils; it infills organisms, such as bryozoa, that possessed open cavities, and picks out the trabecular structure of the crinoid-ossicles.²

The significance of limestones of *a* type lies in this—that their distribution and lithological characters, details of which are reserved for another communication, all point to the conclusions that they have received their iron-content (apart from secondary enrichment) at the time of their deposition, and that the iron-compound has been, from the first, in the condition of oxide and probably, as now, in that of hæmatite. These views, though differing directly from those of Prof. Cayeux as to the origin of a Devonian (Eifelian) hæmatite from the Ardennes which has the characteristic structure of limestones of *a* type,³ are in full agreement with the conclusions of C. H. Smyth, Junr.,⁴ and others⁵ regarding the period of introduction and primary nature of the hæmatite in the well-known Clinton (Silurian) iron-ore of the United States, an ore that also contains a quantity of rolled bryozoan and crinoidal débris, infilled and picked out with hæmatite.

The contemporaneous deposition, in a sedimentary formation unassociated with igneous rocks, of a considerable quantity of an iron compound,—and especially of hæmatite, if that were the original condition of the iron-ore,—appears to be incompatible with open-sea conditions and to demand comparative isolation and proximity to land. Limestones of *a* type are, therefore, directly suggestive of deposition in isolated, coastal areas⁶; but, at the same time, their fauna, which consists largely of standard forms, and the broken and rolled state of their constituents, indicate conditions which, though shallow, depart from those—of abnormal fauna and gentle sedimentation, as we shall see—most characteristic of lagoon-phases.

Purple and green shales.—Purple and green shales that owe their colour to conditions of deposition are of much more restricted

¹ Described by A. Vaughan, Proc. Bristol Nat. Soc. ser. 4, vol. i (1906) p. 99.

² See a microphotograph of a limestone of *a* type in 'The Country around Newport' Mem. Geol. Surv. 2nd ed. (1909) pl. fig. 1.

³ 'Les Minerais de Fer Oolithique de France' fasc. i (1909) pp. 225-26: the structure in question is beautifully figured in this volume. I am greatly indebted to the courtesy of Prof. Cayeux for the use of some of his material.

⁴ Zeitschr. f. prakt. Geol. vol. ii (1894) pp. 304-13.

⁵ For instance, E. F. Burchard & E. C. Eckel in 'Iron Ores, &c. of the Birmingham District, Alabama' Bull. U.S. Geol. Surv. 400 (1910) pp. 28 & 40.

⁶ This view, also, agrees with that of the American observers mentioned above, who regard the Clinton ore as deposited in shallow bays and lagoons (see, for example, *op. supra cit.* p. 40).

occurrence in the Avonian than the preceding rock-type, being recorded only from Km in a few localities. Their unusual colour points to deposition in comparative isolation. Furthermore, the fact that in other formations they are often associated with continental deposits suggests proximity to land; but it remains to be seen whether they have been formed under other conditions of isolation as well.

Calcite-mudstones and similar rock-types.—The second of the groups of rock-types distinctive of *Modiola* phases is much more widespread than either of the preceding types; its members occur in all such phases as are known and at many places in each. They are compact¹ limestones, dolomites, and argillaceous rocks which, under the microscope, are seen to be composed, essentially, of exceedingly fine-grained, calcareous (that is, either calcitic or dolomitic), argillaceous, and siliceous material, and evidently represent, in most cases at least, consolidated muds² of various compositions. For this reason, the calcareous members will be spoken of as calcite-mudstones or dolomite-mudstones,³ in order to distinguish them from limestones or dolomites of standard types. The calcite-mudstones include the rocks known as 'chinastone-limestones,' so called because their compactness and white weathered surface impart to them a porcellaneous appearance. Most of the rocks are evenly bedded, but some have concretionary forms. The calcite- and dolomite-mudstones are either homogeneous and more or less conchoidal in fracture, or are laminated and present a slabby or platy fracture; the laminae, which, in places, are extremely thin, are either persistent or lenticular—wedge-bedded on a minute scale. The homogeneous and the laminated types are equally fine in grain. Similarly, the argillaceous rocks may be either clay-mudstones,—rocks to which the term 'mudstone' is ordinarily applied—, shales, or paper-shales. Under the microscope the bedding in the case of the laminated mudstones, calcareous or otherwise, is clearly shown, even in the absence of fossils, by the layers, which differ greatly one from the other in composition and appearance; but in sections of the non-laminated mudstones it cannot, as a rule, be made out without the help of fossils. Where largely calcareous, such mudstones might be taken for travertine; but that they have been deposited in the form of mud, whether detrital or chemically precipitated, is shown by the fact that, where the material has partly

¹ That is, amorphous-looking, and, in fact, often called 'amorphous' though the constituent grains possess crystal-structure.

² This statement is based on an examination, still in progress, of various limestones and dolomites that occur in *Modiola* phases. It remains to be seen whether it is universally true of the compact layers; in some cases, as, for example, that of landscape-marbles, an origin as travertine by direct accretion from solution has previously been suggested.

³ It is possible that the dolomite-mudstones are calcite-mudstones dolomitized without change of structure, but the balance of evidence at present available suggests that they have been deposited as dolomite-mud.

filled a hollow fossil, its surface against the residual space, originally empty, is flat—in fact, a bedding-plane.

Those mudstones in which calcite or dolomite is predominant are the most important, because they differ conspicuously from rocks formed under standard conditions. It is true that standard limestones frequently contain a considerable proportion of calcite-mud,—so far dolomite-mud has not been detected in them; included with it in such limestones, however, are the remains of open-sea organisms, for example, crinoids, brachiopods, and corals. Limestones which are simply hardened calcite-muds, or contain merely the forms enumerated opposite '1. Shallow-water characters' in Table III (p. 514), appear to be confined in marine series to the distinct groups of strata (*Modiola* phases) under discussion, or, if found among standard deposits, to an occasional occurrence in shallow-water formations.

For the origin of the calcareous portion of the muds we do not need to look beyond the detrition and disintegration of organisms, both within and without the area of actual deposition, and also, to some extent, of contemporaneously-formed¹ limestones (oolites, etc.) and dolomites, fragments of which are found in the interbedded breccias. It is possible, however, that some, or even much, of the mud has been chemically precipitated. But, in any case, the uniform fineness of grain,—which characterizes the intermingled terrigenous material also,—compels a conclusion of much importance, namely, that the sedimentation of the mudstones was exceedingly gentle.

Conditions of deposition of the *Modiola* phases.—Hence we see that the notable feature of the *Modiola* phases represented in Gower is in each an anomalous association of characters. On the one hand, some of the latter indicate deposition in shallow water: in fact, signs of marked current-action, such as the inclusion of contemporaneous fragments, are to be found, in most localities, at some part of each phase. On the other hand, the texture of the rocks affords repeated evidence of the gentlest sedimentation.

The explanation of the anomaly is to be sought in peculiar conditions of deposition. It is conceived that, as the area of deposition of each phase was shallow, and extended far, both parallel and at right angles to the Avonian coast (Table V, p. 526), under certain circumstances the outer part of the area would protect the inner from waves and strong currents, even in the absence of subaërial barriers along the outer margin,—and of the existence of such barriers, as previously mentioned, we have no evidence. The necessary attendant circumstances would be that tides were at a minimum—to which

¹ Some of the mud may have been derived in part from older limestones also, but to an extremely small extent, probably, as regards material deposited in Gower; the phase in $C_2 + S_1$ may, however, owe more of its material to such sources in those parts of the South-Western Province where it rests upon older Avonian limestones unconformably.

condition a steep slope of the sea-bottom down from the shallow area would be favourable—, and that the shallowness of the sea was extreme.

In other words, each phase was deposited in a shallow area of sufficient extent to constitute, under conditions of minimal tides and excessive shallowness, a great series of lagoons as defined in the footnote on p. 511. In such a lagoon-area the pools would catch the finest material, both that of local origin and the part that was brought into the area from various sources outside,¹ and there wafted about by gentle currents. Slightly stronger currents would lead to the washing-up and incorporation of fragments of contemporaneous sediments; and the area would be liable to invasion at times by waters depositing standard rock-types. Such invasions might result either from some deepening of the whole lagoon-area, or from the temporary demolition of a bar—whether subaërial or merely submarine. At other times, local emersion would probably occur and be followed by cracking of the mud. In this way the brecciation *in situ* may have arisen.

Further, as some such conditions of deposition as are here outlined alone explain satisfactorily the whole of those lithological features of the Gower *Modiola* phases that are discussed above, we conclude that the strata of these phases have been deposited under such conditions and are, accordingly, representative of 'lagoon-phases.'

Landscape-marbles and 'pisolites.'—Some remarkable rock-types, however, which appear to be confined to *Modiola* phases, do not receive complete explanation, although their features are not inconsistent with genesis in a lagoon-area; consequently their discussion is deferred. Of these rock-types the chief may be called 'landscape-marbles,' as they reproduce the essential features of the Rhætic bed of that name, itself part of a *Modiola* phase. Landscape-marble is conspicuous in the top of S_2 , the so-called 'concretionary beds,' in the neighbourhood of Bristol and elsewhere; and one of its features is developed on a minute scale at the base of C_2 in North-Western Gower. The so-called 'pisolites' of C_2 and S_2 must have also required for their formation special conditions as well as shallowness.

Faunal characteristics of the *Modiola* phases.—The lagoon-area concept bears the same relation to the faunal peculiarities of the phases as it does to the lithological features; it explains most of them, and accords with the rest. The mudstones, which must be regarded as deposited under conditions of greatest isolation, are also most restricted in their faunal contents, being in many cases unfossiliferous, and in most of the others yielding, whatever be their horizon, only a special, phasal fauna. Of this fauna the members of most general occurrence are certain lamellibranchs, especially *Modiola* and *Sanguinolites*, *Spirorbis*-like annelids, as also ostracods

¹ The colouring-material of the purple and green shales and of the limestones of α type almost certainly came directly from the land, but much other mud, including part of the calcite, may have entered from the seaward margin.

and *Calcisphaera* (?) in abundance; but at many places one or two members only are found. The lack of standard fossils in the mudstones is striking in the case of the phases that we are considering: for the latter occur in a formation, the Avonian, dominated by crinoids and other open-sea forms. A few mudstones, however, as, for instance, some at the top of S_2 , contain many individuals (though but few species) of brachiopods, especially *Seminula*, that lived under standard conditions also. Again, the proximity of the postulated lagoon-areas to the open sea with its abundant fauna, and their liability to invasion, account for the abundance of standard forms in the intercalations of ordinary rock-types, which are frequent in some phases (Km and the top of S_2). The S_2 'pisolites,' also, have been laid down in disturbed waters, and yield an ordinary marine fauna. It may be noted that, in the phase in $C_2 + S_1$, intercalations yielding open-sea forms are extremely rare.

The phasal fauna is essentially shallow-water, but its poverty and peculiar nature demand for full explanation something more than mere shallowness. The isolated position in which the mudstones must have been deposited—whether separated from deeper water simply by extensive shallows or by actual barriers—would be favourable either, under certain circumstances, to an increase of salinity, or, near a river, to a freshening of the waters. In either case much of the standard fauna would be excluded. Some ostracods, like their modern representatives, doubtless could endure, and even thrive under, such adverse conditions. The same appears to have been true of other organisms also. Thus, although lagoon-conditions would favour the accumulation of plankton, including *Calcisphaera* (?), simply on account of its ease of transport from the open sea outside the lagoon-area, the fact that some of the calcite-mudstones contain an abundance of *Calcisphaera* (?), but fewer or no foraminifera, whereas standard Avonian limestones (including those deposited in shallow water) yield these two groups of organisms in inverse ratio, suggests that *Calcisphaera* (?) thrive in the lagoons.

But, whether due to more than mere shallowness or not, the characteristic feature of the calcite-mudstones, the most distinctive rocks of the calcareous lagoon-phases of Gower, is barrenness or a peculiar fauna, rich in individuals but not in species, which, though but poorly represented in intervening standard deposits, is remarkably recurrent, as regards genera and wider groups, in different phases.

(3) The Radiolarian Phase—Cherty Lagoon-Phase—of Gower.

Modes of origin of radiolarian deposits.—The fourth lagoon-phase, that at the base of P, consists of radiolarian cherts with interbedded, laminated shales which have yielded a few lamellibranchs and plant-fragments. As, however, many radiolarian rocks have, in consequence of the classic work of Dr. G.

J. Hinde, been regarded as deep-sea deposits (on account of their exceedingly fine-grained texture as well as the abundance of radiolaria in some recent deep-sea deposits), before proceeding we must show why we conclude that the Gower cherts, which are similar in both fauna and texture, have been formed under diametrically different conditions. This is the more necessary, on account of the weight attaching to the opinion of Dr. Hinde, as well as the importance of the further conclusions based by other geologists on his work.

The way has been prepared by many observers, who have shown that radiolarian rocks have not necessarily been deposited at the depths at which the radiolarian oozes above-mentioned are found; in the case of some there is strong evidence of a shallow-water origin. Thus, at Garn-bica in Carmarthenshire, on the same horizon as the Gower cherts similar rocks have been found by Mr. T. C. Cantrill and described by Dr. G. J. Hinde¹ which, though of finer grain, if anything, than the Gower cherts, may be inferred to have been deposited in shallow waters, on account of their occurrence as a thin, conformable group between two shallow-water formations—the Rottenstone Beds at the top of the Carboniferous Limestone below and the Millstone Grit above.

As may be gathered from the conclusions of Dr. Hinde & Mr. Fox as to the conditions of deposition of the Lower Culm radiolarian rocks of Devon,² of Prof. David & Mr. Pittman in the case of the similar Devonian rocks of New South Wales,³ and of others, as well as from a remark of Prof. Watts in discussing the Australian example (*op. cit.* p. 64), the most characteristic feature, next to their fauna, of all radiolarian rocks is their freedom from any, except the finest, terrigenous material. This character is irreconcilable with an origin in shallow waters of an open sea. A clue, however, to conditions of semi-isolated though shallow deposition is afforded by the calcite-mudstones of the *Modiola* phases previously described. These rocks, though differing from them in composition, resemble radiolarian cherts in being, essentially, exceedingly fine-grained.⁴ Further, some contain an abundance of unicellular plankton—*Calcisphaera* (?). We may, therefore, find parallels for the peculiar features of radiolarian cherts among those of the calcite-mudstones of *Modiola* phases. These mudstones have, undoubtedly, been deposited in shallow waters, but the chief feature wherein they resemble radiolarian cherts, namely their fine grain, is anomalous in shallow-water deposits; as we have seen on pp. 517–18, it is necessary to suppose that the waters were under lagoon-conditions. Similarly, it is concluded that those radiolarian rocks that appear, from independent evidence, to have been formed in shallow waters, owe their

¹ 'The Country around Ammanford' Mem. Geol. Surv. 1907, pp. 72–76.

² Q. J. G. S. vol. li (1895) p. 662.

³ *Ibid.* vol. lv (1899) p. 36.

⁴ The terrigenous material in the mudstones is as fine-grained as the calcareous.

anomalous characters to deposition under lagoon-conditions.¹

In the application of this general principle to particular cases, before a rock-group that satisfies the conditions of exceedingly fine texture and shallow-water origin is regarded as a lagoon-phase, its geographical extent should, if possible, be determined, and for the following reason. For the maintenance of lagoon-conditions it appears to be necessary that the shallow area be of wide extent (see 'Conditions of Deposition of the *Modiola* Phases,' pp. 517-18) ; and certainly their size has been a feature of the areas of deposition of those lagoon-phases that are known.

The Gower radiolarian cherts a lagoon-phase.—In the case of the radiolarian-chert group of Gower and Carmarthenshire, a shallow-water origin is inferred from the stratigraphical evidence cited above, and accords with the presence, in interbedded shales, of lamellibranchs (p. 551) known elsewhere only from shallow-water formations. Also, the group has a wide geographic extent: it crops out in Carmarthenshire at a considerable distance from Gower, in a direction at right angles to the trend of the Avonian coast; and is known through much of South Pembrokeshire, in a direction from Gower parallel to that coast. Consequently, we conclude that it constitutes a lagoon-phase, and, in essentials, differs from *Modiola* phases merely in consisting of the remains of siliceous instead of calcareous organisms, in addition to fine-grained sediment. For contrast it will, occasionally, be convenient to speak of it as a 'radiolarian phase.'

Although this conclusion rests primarily on the evidence cited above, it is confirmed by other features of the radiolarian cherts on this horizon. The first is the fine, sharply-defined lamination which is conspicuous in these, as in other, radiolarian cherts and imparts to them a striped appearance. The laminae differ considerably one from the other: for instance, in the proportion and nature of their detrital material. Further, many of the laminae in the Gower cherts are strongly lenticular, that is, wedge-bedded. The sharply-varying composition and the impersistence of the laminae point to the constant play of gentle currents laden with various, but always fine, sediments. These features, therefore, are to be expected in deposits formed under the lagoon-conditions previously described, and, as a fact, are found in many, including the finest, rock-types of *Modiola* phases, but they do not accord with a deep-sea origin—the only logical alternative to an origin in lagoon-areas in the case of most radiolarian cherts.

¹ Radiolaria are found at the present day at all depths, in greatest abundance in the plankton. Though they swarm in open seas, the environment in some lagoon-areas, also, would be favourable to their increase if the salinity were not appreciably higher than that of average sea-water,—a condition which would be realized in the proximity of a river (p. 525). Their accumulation in lagoon-areas would be further promoted by their ease of transport from the open sea outside.

In one respect, however, the radiolarian cherts differ from all the *Modiola* phases of Gower,—they are devoid of included fragments of contemporaneous sediments.

On the other hand, further confirmatory evidence is yielded by the black Pendleside or *Posidonomya* Limestones of the Midlands and elsewhere. These rocks include the horizon of the Gower cherts and appear, from their peculiar characters, to have been deposited in lagoon-areas¹; their occurrence over a wide extent of country testifies to a wide prevalence of lagoon-conditions on that horizon. Some of them contain an abundance of bodies which, Mr. Howe suggests with much hesitation, may be radiolaria (*op. cit.* p. 400); and in North Devon similar limestones, there known as the 'Venn Limestones,' are associated with radiolarian cherts which, as we shall presently see, probably include the horizon of the radiolarian cherts of Gower.

The Culm radiolarian cherts also a lagoon-phase.—We may now tabulate (Table IV, p. 523) the features of the radiolarian phase, including some which have been observed in the Devon radiolarian cherts, for comparison with those of the Gower *Modiola* phases. The radiolarian nature of the Lower Culm cherts (the Coddan-Hill Beds) of Devon was long ago pointed out by Dr. Hinde & Mr. Fox,² who, however, for the reasons mentioned above, regarded them as deep-sea deposits, the associated Venn Limestones being considered as but somewhat shallower in origin.³ Dr. Wheelton Hind has more recently shown⁴ that these limestones, together with part, at least, of the cherts, are correlative with the Pendleside Series. The correlation of the radiolarian cherts and overlying beds of Gower with the same Series tends to confirm the view, suggested by De la Beche and strengthened by recent work of the Geological Survey,⁵ that the radiolarian cherts of the two areas occur at one horizon.⁶ (It is probable, of course, that the Devon cherts, which are much thicker than those of Gower, include other horizons also.) The close similarity⁷ of the cherts of Devon and Gower is such strong presumptive evidence of community of origin, that it is difficult to believe that those of the one area have been formed in comparatively deep water and those of the other in shallows. And my own observations (see Table IV, p. 523) in

¹ With this view Mr. J. A. Howe, who first adequately described the rocks, *Q. J. G. S.* vol. lvii (1901) p. 399, tells me that he agrees.

² *Q. J. G. S.* vol. li (1895) p. 609.

³ Of the shallow-water origin of the Venn Limestones there can, however, be no doubt.

⁴ *Geol. Mag.* dec. 5, vol. i (1904) p. 392.

⁵ See A. Strahan, *Swansea Memoir*, pp. 22–25.

⁶ It may be useful here to mention that the horizon in Gower with which the Coddan-Hill Beds are homotaxial is that of the radiolarian cherts, not of the underlying beds with trilobites and brachiopods referred to by Dr. Wheelton Hind (*Geol. Mag.* 1904, p. 402); the latter belong to D₂₋₃.

⁷ The similarity has been remarked by others, including Dr. G. J. Hinde (*Swansea Memoir*, p. 25), and extends to the fine lamination and minute wedge-bedding that I have adduced as evidence against a deep-sea origin for the Gower cherts.

TABLE IV.—THE RADIOLARIAN PHASE AT THE BASE OF P.

Characters unusual in deep-sea deposits or directly suggestive of an origin in shallow water:—	(1) Faunal. ¹	The sole macroscopic organisms of some interbedded shales are the lamellibranchs (unbroken) mentioned on p. 551, known elsewhere only from shallow-water formations (Gower).
	(2) Stratigraphical	Occurrence as a thin group conformably intercalated between two shallow-water formations,—the Rottenstone Beds and the Millstone Grit (Carmarthenshire ⁴). Presence of intercalated beds of 'culm' (coal), several inches thick (Barnstaple ²).
	and Lithological.	Lenticular, wedge-bedding of many laminæ, the latter sharply defined and differing considerably in composition one from the other.
Characters distinguishing the phase from ordinary shallow-water marine formations.		Fine grain of the terrigenous material. Abundance of radiolaria.
Ordinary sediments included within the phase.		Very rare, if existent: for, although some beds resemble ordinary shales and silts and, in Devon, some contain corals, etc., ³ radiolaria appear to be abundant throughout and no coarse sediment is known.
Relations to the beds:—(1) below		Conformably resting upon the Rottenstone Beds in Carmarthenshire. ⁴ Not known with certainty in Gower, Pembrokeshire, or North-Western Devon. ⁵
(2) above		Sharp, but conformable, change to the Millstone Grit in Carmarthenshire. ⁴ Not known with certainty in Gower, where the group gives place to shales, apparently without recurrences of radiolarian cherts. Not known with certainty in North-Western Devon.
Development in (1) Bristol area ... (2) Gower (3) outcrops south of the Pembrokeshire coalfield. (4) North-Western Devon.		Absent. Present throughout (and to the north in Carmarthenshire). Present throughout, except in the southernmost (Bosheston) outcrop. Present throughout ⁶ ; thickness, and probably duration also, much greater than in South Wales.

Note.—Features of general occurrence are not followed by the name of a locality.

¹ The fauna of the Devon development, as recorded by various authors, contains no forms which need be supposed to have lived at great depths. For lists see G. J. Hinde & H. Fox, *op. cit.*,—all orders; H. Woodward, *Geol. Mag.* 1902, p. 481,—trilobites; Wheelton Hind, *Geol. Mag.* 1904, p. 398,—all orders; A. Vaughan, *Geol. Mag.* 1904, p. 531,—brachiopods and corals.

² First brought to my notice by Mr. J. G. Hamling.

³ G. J. Hinde & H. Fox, *Q. J. G. S.* vol. li (1895) pp. 617, 643 *et seqq.*

⁴ T. C. Cantrill in 'The Country around Ammanford' *Mem. Geol. Surv.* 1907, p. 75; and in conversation.

⁵ The Devonian and Lower Culm of Fremington Pill, Barnstaple, said by De la Beche to pass one into the other ('Rep. Geol. Cornwall, Devon, & West Somerset' *Mem. Geol. Surv.* 1839, p. 103), appear at present to be confined to isolated exposures affording no evidence on the question.

⁶ G. J. Hinde & H. Fox, *op. supra cit.*

the Barnstaple district of North Devon,¹ so far as they go, suggest that the cherts cropping out there are lagoon-deposits rather than those of a fairly deep sea. Consequently, as we need consider only these two alternatives, it is concluded that the Carboniferous radiolarian cherts of Devon, which continue through a wide extent of the West Country, constitute a lagoon-phase. This conclusion is not impugned by any of the known features of these rocks in the rest of their outcrops. Various particulars are given in the comprehensive work of Hinde & Fox, and those that bear on the question are mentioned in Table IV (p. 523). There is, however, much to be learnt concerning this highly interesting group, and it is well that a detailed examination of the Lower Culm and Upper Devonian has been undertaken by my colleague, Mr. H. Dewey.

(4) Meaning of the Difference between the *Modiola* Phases and the Radiolarian Phase of Gower.

Although the radiolarian phase of Gower appears to have been deposited under the same conditions of depth as the *Modiola* phases, in facies it presents a great contrast. Each phase, however, reflects the character of the fauna of the sea adjacent to the lagoon-area in which it was formed, as the terrigenous material is, typically, similar in all; the difference resolves itself, therefore, into one between the facies of the contiguous open-sea fauna.

The open sea during the *Modiola* phases was rich in forms possessing calcareous skeletons and giving rise by detrition to much calcareous mud; some of this, doubtless, passed into the lagoon-areas with fine terrigenous material, to form, with the detritus of the indigenous fauna and, possibly, calcareous material chemically precipitated, the characteristic mudstones. But the radiolarian rocks were formed at a time (the commencement of P) when, in the South-Western Province, the sea underwent a great and persistent change in character. Calcareous organisms disappeared almost completely from that province, being chiefly represented throughout P and higher parts of the Carboniferous by thin-shelled goniatites and lamelli-branchs. Siliceous forms, however, assumed greater relative importance, both in the ground-living fauna,—sponge-spicules being noticeable in the radiolarian cherts and at some higher horizons themselves forming cherts,—and among the free-swimming groups, where radiolaria became conspicuous.

But, although it is evident that in our area the sea from the commencement of P onwards was inimical to so rich a calcareous fauna as it had, up to that time, supported, yet not to a siliceous one, we do not know what precisely was the change in its character; for apparently that effect on its fauna might result from any one of several changes, such as a diminution in its soluble calcium-

¹ I take this opportunity of thanking Mr. J. G. Hamling, F.G.S., most heartily for placing his extensive knowledge of this district at my disposal.

content, or a reduction of temperature.¹ Neither its muddiness nor its probable shallowness would account for the faunal change, as similar influences had previously had no such effect, even when acting in conjunction, as during Km. But, whatever may have been the nature of the change in the sea, its radical character and the wide extent of the area affected point to an important change in the physiography of the region. Further, there is reason to believe that the change in the sea accompanied a marked increase in the influence of the waters from a neighbouring river (p. 539). And that radiolaria may abound in an area under such influence appears to be shown by their occurrence in quantity in the Lower Culm of Eastern Thuringia and the Vogtland, which is regarded, from the abundance of land-plants that it contains and for other reasons, as having been deposited in a muddy sea near the mouths of rivers.²

(5) Topographical Position of Lagoon-Areas.

That the areas of deposition of the *Modiola* phases of Gower were situated, probably in each case, immediately between the coast of the Avonian sea, which lay to the north throughout the South-Western Province, and the open water may be inferred from the following details, partly outlined in Table V, 2, p. 526. In arriving at this conclusion, true overlap of a phase by conformable higher beds is accepted as evidence that the lagoon-area in which that phase has been deposited has been limited by a shore-line along the line of overlap; while lateral replacement by ordinary marine sediments is accepted as evidence of the deeper-water limit of the area. For this reason a lagoon-area has been defined (p. 512) as essentially coastal.

The lagoon-phase at the base of K in Pembrokeshire overlaps the conformable Upper Old Red Sandstone in a northerly direction, and is itself in turn overlapped by conformable higher beds, which then rest directly upon Lower Old Red Sandstone or upon still older rocks. The deeper-water limit, also, of the lagoon-area appears to be reached in that county: for, at Freshwater West in the southernmost outcrop, the Upper Old Red Sandstone is followed immediately by ordinary marine sediments in which no lagoon-deposits have yet been found.

The lagoon-phase at the base of $C_2 + S_1$ in Pembrokeshire succeeds the *Caninia* Oolite conformably south of West Williamston, but at that place an unconformity develops at its base. The proximity of the lagoon-area to a shore-line, thus suggested, appears to have persisted throughout the phase, for at Pendine

¹ L. W. Collet, 'Les Dépôts marins' Paris, 1908, pp. 10-22 & 219.

² J. Lehder, Neues Jahrb. Beilage-Band xxii (1906) pp. 79-80, 111.

TABLE V.—RELATIONS AND GEOGRAPHICAL EXTENT OF THE CARBONACEOUS LAGOON-PHASES REPRESENTED IN GOWER.

	<i>Km, the lagoon-phase at the base of K.</i>	<i>The lagoon-phase at the base of C₂ + S₁.</i>	<i>The lagoon-phase at the top of S₂.</i>
1. Relations to the beds :— (1) below.	Generally conformable on Upper Old Red Sandstone, but in places (Chew, Pem.) unconformable on Lower Old Red Sandstone.	Generally conformable on <i>Canaia</i> Oolite, but in places unconformable on <i>Canaia</i> Oolite (West Williamson, Pem. ³) or lower horizons (Pendine, Carm. ⁴ ; etc.).	Conformable and intercalated with standard S ₂ oolites.
(2) above.	Sharp change to standard deposits of K ₁ age ('Avon, p. 98; etc. ¹).	Abrupt change, ⁵ in places with some contemporaneous erosion, to standard deposits of C ₂ or S ₁ age.	Rapid change to standard deposits which form the base of D ₁ .
2. Development ⁶ in (1) the Bristol area and the Mendips,—eastern part of the South-Western Province.	Present throughout. ²	Thick, extending to the base of S ₁ , in the north (Avon, etc. ⁶). Thin, forming the base of C ₂ , at Weston-super-Mare. Absent in the south (Mendips).	Present throughout,—maximum at Sodbury. ⁷
(2) Gower,—central part of the South-Western Province.	Probably present throughout.	Thin, forming the base of C ₂ , in North-Western & Eastern Gower. Absent in South-Western Gower.	Present throughout, but barely developed in South-Western Gower.
(3) Outcrops south of the Pembrokehire coal-field,—western part of the South-Western Province.	Recognized throughout, except at Freshwater West on the southernmost outcrop.	Thin, forming the base of C ₂ , in northern outcrops (West Williamson; Tenby). Absent in outcrops south of Tenby.	Present throughout, but poorly developed in the southernmost (St. Gowan's Head) outcrop.

¹ The change is said by Dr. Vaughan (*loc. cit.*) to be 'much sharper in the Avon sequence than in any other part of the South-Western Province,' but has been found to be equally sharp in Pembrokehire.

² The Country around it is absent, and the lagoon-phase commences and ends at the base of S₂ (see pp. 527-28).

³ At Pendine C₂ and S₁ are present, and the lagoon-phase commences and ends at the base of S₂ (see pp. 527-28).

⁴ In the Avon district (Bristol paper, pp. 194-95; Avon paper, pp. 94-95) there is less contrast than at most places between this phase and the succeeding deposit; but evidently much of the whole of C₂ and S₁ there present has been formed under lagoon-conditions.

⁵ A. Vaughan, Bristol paper, p. 222 (Tyrtherton); p. 207 (Sodbury); p. 194 (Avon).

⁶ A. Vaughan, Bristol paper, p. 222 (Tyrtherton); p. 207 (Sodbury); p. 194 (Avon).

⁷ *Ibid.* p. 210.

⁸ At present we know too little of the development outside the areas considered to make its discussion profitable.

(Carmarthenshire),¹ on the other (the northern) side of the coal-basin, the whole of $C_2 + S_1$ has been overlapped by S_2 .

Though the above statement suffices for the purpose of this section, it is advisable here to consider a little further the interrelations of the developments at West Williamston and at Pendine. At the latter place limestones, apparently no younger than Z, are followed in order by (1) a limestone-conglomerate; (2) a few feet of lagoon-deposits—barren, black calcite-mudstones; and (3) good representatives of S_2 and D, commencing with the base of S_2 . There is evidence, in the composition of the conglomerate and in 'contemporaneous piping' of the limestones below, that the latter are separated from the conglomerate by a true unconformity: that is, that they were raised above sea-level and eroded subaërially prior to the deposition of the conglomerate. Here the unconformity is marked also by a considerable hiatus in the sequence; on the other hand, at West Williamston, the only place on the south side of the coal-basin where an unconformity in the sequence is evidenced, the break is slight and lies between C_1 and C_2 , in other words, on a horizon in the middle of the hiatus at Pendine. Nevertheless, a consideration² of the general development of the Avonian in the South-Western Province leads to the conclusion that the unconformities at both places have been caused by one series of earth-movements. The magnitude of the hiatus at Pendine, and the fact that the horizon of the uplift lies within this hiatus, are explicable

¹ 'The Country around Carmarthen' Mem. Geol. Surv. 1909, pp. 81-82.

² 'The Country around Haverfordwest,' Mem. Geol. Surv. (in the press).

on the supposition that beds below the plane of unconformity are overstepped between the latter place and Pendine and that beds above that plane are overlapped in the same direction (see fig. 7, p. 527). Both of these phenomena are to be expected, though the extent of the overlap above the unconformity is surprising. Further, the presence of lagoon-deposits at the base of the overlapping series at Pendine and some other places along the same outcrop, as well as at West Williamston, suggests that across the unexposed ground between the latter place and Pendine the unconformity is succeeded, as a rule, by lagoon-deposits. This is also indicated diagrammatically in fig. 7.

The deeper-water limit of all horizons of the phase is reached in the south: for they are represented by ordinary limestones in the Mendips, South-Western Gower, and Southern Pembrokeshire.

The lagoon-phase at the top of S_2 has not been followed out in detail northwards; neither has its complete seaward limit been reached, although in Gower and Pembrokeshire it is poorly developed in the southernmost outcrops, its phasal deposits being there largely replaced by ordinary limestones.

(6) Earth-Movements accompanying Lagoon-Conditions.

The deposition of a lagoon-phase involves the fulfilment of certain conditions. Some of these are enumerated in paragraph 1 (below). Again, certain conclusions may be drawn from the development in the South-Western Province of the *Modiola* phases present in Gower: as the latter are few, these conclusions are provisional; but as, at the same time, the evidence is consistent, it may be useful to note its trend. It will also be seen that the deposition of the Gower radiolarian phase appears to have followed an earth-movement similar to one which preceded the deposition of the *Modiola* phase at the base of C_2 .

1. The accumulation of extremely shallow-water deposits to any considerable thickness can have been conditioned only by a general depression of the sea-bottom relative to sea-level—positive movement—over the area of deposition. Further, the depression has been isostatic,—the term ‘isostatic’ being used simply in a descriptive sense, to denote the state of an area over which the rate of sedimentation is equal to the rate of depression. Lagoon-phases are, by definition, extremely shallow-water deposits; the accumulation of each has, therefore, been accompanied by isostatic depression, provided its thickness is greater than the range of depth of the water in which it has been deposited.

2. The internal constitution of the Gower *Modiola* phases (see Table III, 4, p. 514) suggests (i) that in some cases (Km and the top of S_2) isostasy has been interrupted; but that in the other (the base of $C_2 + S_1$) it has been accurately maintained, practically throughout the duration of the phase.

3. A consideration of the nature of the beds below and above and

of their relations to the phases (see Table V, 1, p. 526, and 'Variations in Depth, etc.' pp. 532 *et seqq.*) leads to the further conclusions:—(ii) The periods of isostasy recorded in the phases have followed either a movement of elevation (in the case of the phase at the base of $C_2 + S_1$); a continental period, during which depression has not been exactly counterbalanced by sedimentation (in the case of Km); or a shallowing, marine period (in the case of the phase at the top of S_2). (iii) Each period of isostasy has been ended by a sudden movement of depression.

4. Taking into account, also, the fact that the lagoon-areas have been coastal features, we may generalize the two last conclusions (par. 3) thus,—lagoon-areas have been developed during periods of steady depression, more or less prolonged, which have either marked the first stage of the submergence of a foundering coast, or followed intervals of upward movement interrupting a general downward movement of a sea-floor. (The one condition is probably a particular case of the other.) These periods of steady depression have preceded more rapid subsidences.

It follows, as a corollary, that lagoon-phases may be expected to accompany transgressions, as do Km and the phase at the base of $C_2 + S_1$, but also that they will be subject to overlap by the deeper-water deposits that have succeeded them.

5. The phase at the base of $C_2 + S_1$, of which our knowledge is most complete, persisted far longer in the northern than in the southern part of the Bristol area: for in the north it is succeeded by S_1 beds, whereas at Weston it is confined to the base of C_2 . The significance of this fact lies in the greater proximity of the former region to the Avonian coast. In the western area (Pembrokeshire and Pendine) also, the phase apparently persisted longer near the Avonian land than elsewhere (fig. 7, p. 527), although there the matter is complicated by overlap of higher horizons over lower. It appears, therefore, that the depression that closed the phase affected the seaward margin of the lagoon-area before the more landward parts. The depression was greatest in the south, both in the Bristol area and in the west.

No such landward lag is known in the case of the depression that ended the lagoon-phase at the top of S_2 .

The radiolarian phase.—The conclusions regarding earth-movements, detailed above, afford a means of testing the hypothesis that the radiolarian cherts at the base of P are similar in origin to the *Modiola* phases on which those conclusions are based.

The formation of the cherts was preceded in Gower by a shallow-marine period (D_1 to D_{2-3}). The following evidence suggests that this period was closed by elevation. At Ifton,¹ near Newport (Mon.), on the east, as well as at Haverfordwest² on the west, Millstone Grit, similar lithologically to the band that succeeds the

¹ Rep. Brit. Assoc. (Winnipeg) 1909, p. 478.

² O. T. Jones, 'Summary of Progress for 1906' Mem. Geol. Surv. 1907, p. 53.

radiolarian cherts in Carmarthenshire immediately and conformably, rests unconformably upon S_2 ; at Ifton cavities (channels and swallet-like passages) penetrating the limestone to a depth of at least 40 feet below its uneven upper surface have been filled with an original deposit of Millstone Grit. The undisturbed character of the Grit in the cavities shows that, here as at the Haverfordwest occurrence described by Prof. O. T. Jones, the limestone was raised above sea-level and 'piped' by subaërial erosion prior to the deposition of the Grit. No direct evidence as to the exact period of the elevation is available; but the fact that the change from D_{2-3} to P has been marked by a considerable physiographic change in the South-Western Province (p. 524) suggests that the movement took place at the close of D.¹ The relationship of the uplift to the hiatus between S_2 and the Millstone Grit would then be analogous to the relationship of the uplift at the end of C_1 to the hiatus between Z and S_2 at Pendine (fig. 7, p. 527). Outside the South-Western Province, evidence of emersion between D and P has been found in places by Dr. Sibly and Mr. C. B. Wedd, although, as a rule, there is no such break at this horizon.²

It appears, therefore, that the radiolarian cherts of the South-Western Province have been the first deposits formed after a movement of elevation which has left indubitable traces in that area. Consequently, tested in this way their suggested relationship to *Modiola* phases receives support.

(7) Lagoon-Phases outside the South-Western Province.

In conclusion, it may be mentioned that the conditions under which the lagoon-phases of Gower have been deposited have been paralleled elsewhere and at many other horizons. Where, in such cases, the sediments have a calcareous, or a calcareous and argillaceous, facies, as, for example, the Burdiehouse-Limestone Group of Scotland, the Upper Rhætic and Cotham-Marble Group,³ and the Solenhofen Slate, their origination in shallow areas or even lagoons has been generally recognized, despite the fine grain of some of their most characteristic rocks. It is of interest, therefore, that several resemble the phases of Gower, not only in this fine grain and, frequently, in a fine lamination, but also in their relation to earth-movements.

As regards the radiolarian phase at the base of P, though it is not suggested that all radiolarian deposits have had a similar origin, it is certain that some, at least, of those which appear, from various considerations, to have been deposited in shallow water

¹ It is not, however, suggested that the movement resulted in unconformity everywhere, for in Carmarthenshire the sequence both below and above the radiolarian cherts is unbroken; see T. C. Cantrill, 'The Country around Ammanford' Mem. Geol. Surv. 1907, pp. 72-76.

² Q. J. G. S. vol. lxiv (1908) pp. 63 & 81.

³ The Lower Rhætic appears to be less peculiar in its rock-types than the Upper.

possess features linking them so closely with the Gower cherts that they, also, may be inferred to have been formed under lagoon-conditions. As an example may be cited the Gondwana plant-beds with radiolaria, described by Dr. A. K. Coomáraswámy¹ as having been 'deposited in comparatively shallow water.' They resemble the Gower cherts in their fine grain and very thin bedding; and, like various lagoon-phases, they occur near the base of a transgressing series, for they lie a short distance above gneiss, from which they are separated by merely a small thickness of sandstone, etc.

The close connexion between the development of lagoon-areas and certain earth-movements, discussed in the preceding section, and their locus on coastal shelves are related, possibly, to the fact that some *Modiola* phases, such as the Burdiehouse-Limestone Group, have been accompanied by volcanic outbreaks, including the effusion of lavas which possess a pillowy structure,² though the available data are insufficient to show whether such outbreaks have been connected with *Modiola* phases to a greater extent than with the contemporaneous standard deposits.

In this connexion, however, it is important to note that radiolarian cherts in various parts of the world are frequently accompanied by volcanic rocks, especially lavas possessing a pillowy structure; and, as bearing on this aspect of the subject and the views which I have ventured to put forward, the following words of Prof. Charles Lapworth are significant. Speaking of a hypothetical area on which radiolarian cherts and associated igneous rocks might be supposed to have been deposited, he said³:—

'Consider, for example, such an area as originally forming part of a slightly submerged continental shelf, or coastal platform, with volcanic and archipelagic conditions, and overlooking a broad and deeper sea. Suppose, further, the platform itself flooded by rocks already metamorphosed, and that this platform remains covered by shallow waters for an extended period of geological time, while the sea-floor in front of it is continually deepening. Under these conditions, only such mechanical, volcanic, or organic material could become accumulated as rock-layers on the submerged platform, as by their original nature or rapid cementation were incapable of being swept off by the waves and currents of the shallow waters into the open and deepening sea beyond. Such rock-formations as would be accumulated on the platform would necessarily be thin, but would be lithologically varied and peculiar.'

This graphic passage, with certain modifications, embodies the general conclusions to which I have come as to the situation in which lagoon-phases, so 'varied and peculiar' in their features, have been deposited. The chief difference lies in my interpretation of some of those features; they point to the dominance of comparatively still water within the shallow area.

¹ Geol. Mag. 1902, p. 305; see especially p. 306.

² Sir Archibald Geikie, 'Central & West Fife & Kinross' Mem. Geol. Surv. 1900, p. 54.

³ Q. J. G. S. vol. lviii (1902) p. 440.

V. INTERPRETATION OF THE LITHOLOGICAL SEQUENCE [E. E. L. D.].

(1) Preliminary Remarks.

Apart from many minor variations in rock-type, a general sequence of facies (Table II, p. 505) is presented by the Avonian of Gower. In the following interpretation of this sequence it has been assumed that those true oolites—that is, rocks consisting chiefly of typical, concentric and radially-crystalline ooliths,—that are widespread have been deposited in shallow water. The assumption is justified by the current-bedding of such rocks and the frequency of fragments of contemporaneously-formed oolite in them, as well as by what is known both of the conditions determining the formation of ooliths¹ and of the situations in which ooliths are found at the present day. The oolites include some of the purest of the Avonian limestones, but the fact that limestones practically free from terrigenous material may be formed in shallow, coastal waters is so well known as to need no emphasis.

The differences between the three districts (see Table I, facing p. 505) will be discussed later.

(2) Variations in Depth and other Conditions of Deposition over Gower as a whole at Successive Times; Conclusions; Earth-Movements.

The variation of depth in time is represented diagrammatically by fig. 8 (p. 533). In this there is no true vertical scale; the depth-line indicates merely periods of extreme shallowness, and, by the direction of its slope, shallowing or deepening. For horizontal scale, the duration of a subdivision is arbitrarily and only approximately made proportional to its thickness.

In the following discussion, on which the figure is based, separate reference will not be made to the depth of the *Modiola* phases; in accordance with the views already put forward, they are represented as extremely shallow, but with some interruption in the case of Km and, to a greater extent, in that of the top of S₂ (see Table III, 4, p. 514).

(i) The change from Upper Old Red Sandstone to Km.—This was a change from 'continental' to marine conditions: it is represented in fig. 8 (p. 533) by the depth-line passing below sea-level. The change was sudden, and was unaccompanied, so far as known, by marine intercalations in the top of the Old Red or continental recurrences in the base of Km; but, despite a little current-action, deposition was practically continuous.

(ii) The change from Km to K₁.—Sharp increase of depth.

(iii) Conditions during K₂.—The sea was muddy, but sup-

¹ G. Linck, Neues Jahrb. Beilage-Band xvi (1903) p. 495; and L. W. Collet, 'Les Dépôts marins' Paris, 1908, pp. 282-83.

² On the supposition that K is equivalent to the Lower Limestone Shales (see p. 496).

Fig. 8.—Diagram showing the variation in depth of the Avonian Sea at successive times, deduced from the sequence in Gower.¹

¹ For explanation, see V, § (2), pp. 532 *et seqq.* The bracket against the *Modiola* phase at the top of *S₂* is unclosed on the left, as the phase commences indefinitely. In accordance with the conclusion on p. 521, the radiolarian phase at the base of *P* is represented as deposited in as shallow waters as those of the *Modiola* phases.

ported a rich calcareous fauna belonging to standard groups, chiefly crinoids and brachiopods; the remains are found in both the shales and the subordinate limestones. In Gower, as in the South-Western Province generally, the conditions, for some reason, were unfavourable to the formation of chert and dolomite. The thick limestone in the middle of the zone, which is widespread, consists partly of oolite, and may, therefore, have resulted from a temporary shallowing (fig. 8).

(iv) The change from *K* to *Z*.—Probably in Gower, and certainly over the western part of the South-Western Province, this change of fauna was accompanied by a rapid and long-enduring lithological change, from a dominantly argillaceous to a dominantly calcareous facies of sedimentation: for there the junction of the zones is the junction of the Lower Limestone Shales with the main limestone-mass of the Carboniferous Limestone Series. This widespread change in fauna and rock-type probably resulted from some important physiographic event elsewhere, but apparently it was not accompanied in Gower by change of depth, for the *Z* limestones closely resemble the thin limestones in *K*. It is noteworthy that 'contemporaneous' dolomitization and the formation of chert practically came in at the period of the change; chert is especially frequent near, but not below, the junction.

(v) Conditions during the two sequences: (1) *Z*, *C₁* and the *Modiola* phase at the base of *C₂*; and (2) *C₂* & *S*.—A close parallel may be drawn between these two sequences, especially in South-Western Gower where the lithological series is

most varied and most typical of the South-Western Province generally. In the following table the members of each sequence are placed in order of superposition, and lettered like the parallel member of the other sequence :—

(1) The sequence Z, C ₁ and the base of C ₂ :—	(2) The sequence C ₂ and S :—
<p><i>d.</i> <i>Modiola</i> phase at the base of C₂.</p> <p><i>c.</i> <i>Caninia</i> Oolite.</p> <p><i>b.</i> <i>Laminosa</i> Dolomites (dolomitized crinoidal limestones); in Eastern & North-Western Gower with similar dolomites of Z-age.</p> <p><i>a.</i> Crinoidal limestones, some dolomitized; lower part or all of Z.</p>	<p><i>d.</i> <i>Modiola</i> phase at the top of S₂.</p> <p><i>c.</i> S₂ oolites.</p> <p><i>b.</i> Partly dolomitized gasteropod-limestones, most conspicuous in S₁.</p> <p><i>a.</i> Various limestones, some dolomitized; C₂ with part of S₁.</p>

The chief differences between the two sequences are (1) dolomitization has been much less complete in S₁ than in the *Laminosa* Dolomites; (2) the S₂ oolites, on the contrary, are much thicker and include dolomites more frequently than the *Caninia* Oolite; (3) the S₂ oolites pass gradually into the overlying *Modiola* phase, whereas the *Caninia* Oolite is separated from the parallel phase by a sharp line and, in places, contemporaneous erosion. (In fig. 8, p. 533, the passage of the S₂ oolites into the *Modiola* phase is indicated by the overlapping of unclosed brackets.)

As regards interpretation, it is clear that in each sequence the establishment of the *Modiola* phase (*d*), that is, of lagoon-conditions, has been preceded by a considerable period during which shallowness of the sea, though less extreme than during the phase itself, has had a marked effect on the facies of sedimentation: this is shown by the presence of thick dolomites (*b*)¹ and oolites (*c*) below each phase. But it is uncertain what cause has led to the replacement, in each sequence, of dolomite- by oolite-formation, for the two sequences are sufficiently parallel to suggest that the cause has been similar in both. Though, in view of the frequency of contemporaneously-eroded fragments in the oolites (*c*), we may well suppose that the waters in which these rocks have been formed have been shallower than during the preceding dolomititic stages (*b*)—just as the latter have been shallower than the preceding stages (*a*),—it seems probable that oolite-formation has depended on some condition additional to shallowness. For, in the Avon, the incoming of S₂ is marked, as at many places in the South-Western Province, by the

¹ The view of Prof. E. W. Skeats, Q. J. G. S. vol. lxi (1905) pp. 133-38, and others, that within certain limits shallowness favours dolomitization, is adopted; it is supported by the relations observed (see below, p. 537) between dolomitization and the probable relative depths of the Avonian sea. Although S₁ thus bears evidence of deposition in shallower waters than those of C₂ in Gower, in the Avon it includes some limestones of deeper-water origin than any in C₂.

appearance, for the first time in S_1 of thick oolites; but, at the same time, S_1 appears, from the recurrence through it of lagoon-deposits, to have been deposited in waters even shallower at times than those of the S_2 oolites.

It may be noticed that, throughout the series Z-S in Gower, the junctions of the zones and subzones do not correspond with marked lithological changes, except in the cases of C_1 with C_2 and (possibly in places) of S_1 with S_2 .¹

(vi) The change from the *Modiola* phase at the top of S_2 to D_1 .—The rapid replacement of the *Modiola* phase by deposits of open-sea, though shallow-water, facies points to rapid but not great deepening; the ensuing deposits yield a D fauna, and constitute the basal beds of the D zone. The migration of the D fauna from the region where it had developed into our area appears to have coincided, therefore, with a deepening. The increase of depth was probably immaterial, however, for the preceding fauna was ousted both over areas where standard conditions had been dominant during S_2 , even to the end, as in South-Western Gower, and over those where lagoon-conditions prevailed towards the close of that time, as in Eastern and North-Western Gower.

(vii) Conditions during D_1 and D_2 or D_{1-2} .—No rock-sequence comparable with that above the *Modiola* phase at the base of C_2 exists in D. The initial deepening of the area of deposition appears to have been slight and brief: for, once or twice during D_1 thin coals and underlay have been deposited, pointing to but little depth at those times, though not necessarily to actual terrestrial conditions. These shallow periods are indicated by the peaks in the depth-curve in fig. 8 (p. 533). It is noticeable that, despite the shallowness, a lagoon-phase has not been established, the coal being intercalated among standard marine limestones.

(viii) The change from D_2 or D_{1-2} to D_{2-3} (Upper Limestone Shales).—The incoming of the D_{2-3} fauna accompanied a reappearance of shale-depositing waters: such waters had been practically absent from our area since K-times. The D_{2-3} deposits, however, were much more calcareous than those of K; and both chert and dolomite were formed in considerable amount.

(ix) The change from D_{2-3} to the radiolarian phase at the base of P.—In regard to depth of deposition, this change resulted from the substitution of comparatively shallow conditions by what appear to have been lagoon-conditions (p. 521). In regard to facies of deposition, it resulted from a replacement of the calcareous (or calcareo-argillaceous) conditions that had prevailed in Carboniferous times hitherto by the non-calcareous conditions—whether cherty, argillaceous or arenaceous, marine or non-marine—that prevailed in the South-Western Province for the rest of Carboniferous times.

(x) Conditions during P.—Little is known with certainty

¹ In South-Western Gower, where the series has been most closely examined, the S_2 oolites come in a short distance above the base of S_2 .

of the poorly-exposed sequence above the radiolarian cherts; the P-shales, which form the lower part of the Millstone Grit Shales of the Geological Survey, pass up insensibly through the higher part into the Coal Measures—the latter being, in part at least, a ‘continental’ deposit.

Conclusions.

(1) The time represented by the Gower succession discussed above may be divided into four periods, namely (i) $Km-C_1$; (ii) C_2-S_2 -oolites; (iii) top of S_2-D_{2-3} ; and (iv) P. During these periods the depth varied as follows:—

(i) $Km-C_1$. The area at first was under lagoon-conditions (Km); its depth increased sharply (K & Z), and later diminished, in all probability progressively but intermittently (Z to *Laminosa* Dolomites; *Laminosa* Dolomites to *Caninia* Oolite), ultimately attaining that of lagoon-conditions again by the time that the next period was initiated.¹

(ii) C_2-S_2 oolites. The area, with the exception of the South-Western district, at first was under lagoon-conditions (*Modiola* phase); its depth increased abruptly—to an amount during the rest of C_2 greater, probably, than at any other Avonian horizon, and later diminished, again, it would appear, progressively but intermittently (S_1 ; S_2), to that of lagoon-conditions by the time that the next period was initiated. The stage of oolite-formation was greatly prolonged, and the shallowness which marked it culminated in the lagoon-conditions that followed.

(iii) Top of S_2-D_{2-3} . The area at first was under lagoon-conditions (*Modiola* phase); its depth increased rapidly, though probably to no great amount, and later was at times very slight² (coal and underclay in D_1).

(iv) P. The area at first appears to have been under lagoon-conditions (radiolarian phase), but its subsequent history during P is uncertain. The ultimate replacement of marine waters by continental ones (Coal Measures) was probably gradual.

(2) The foregoing summary of the periods shows that during each of the first three the area has undergone a bathymetrical cycle, complete in the case of the first two and probably in that of the third also; in each of the three cases initial shallowness, that of lagoon-conditions, has been followed by rapid deepening and, later, by shallowing, which, unlike the deepening, has been prolonged. The shallowing has, however, been more regular in the first two than in the third period.

Earth-Movements.

(3) Both the deepening and the shallowing have probably resulted

¹ At some places in the South-Western Province, such as West Williamston, the shallowing culminated in emersion, the lagoon-conditions being established as the land again sank at the commencement of the next period.

² At some places in the South-Western Province (see pp. 529-30) this period probably ended in emersion.

from earth-movements. The shallowing may be due to accumulation of material, but, for several reasons, has more probably resulted from movements of elevation (negative movements). Thus, in two of the cycles it appears to have been progressive and intermittent; and in one (if not two) it has been completed in parts of the South-Western Province by emersion.

(4) If we knew the exact depth in feet at which each uniform group of the series had been deposited and the time occupied in the deposition, by adding, for each group in turn, depth to total thickness of deposit already formed, we could construct a curve with these totals as negative ordinates and total times as abscissæ. This curve would show the movement of the base of the Carboniferous throughout Avonian time. It would then be seen that the negative movements above mentioned have been mere incidents in a prolonged spell of depression (positive movement). Thus, although the incoming of oolites in S_2 has, doubtless, been accompanied by a certain amount of negative movement, their deposition to the thickness which they attain can have been conditioned only by a renewal, and for a long time, of depression. In this case, where the thickness of the deposit is enormous relatively to any possible range of depth during its formation, depression has been approximately isostatic (in the sense in which that term is used on p. 528).

(3) Relations between the Different Districts of Gower at Successive Times; Conclusions; Earth-Movements.

The following interpretation is placed on the differences and resemblances between the three districts, summarized in Table I (facing p. 505).

(i) Of K little is known, but during the deposition of Horizon β the facies of sedimentation was practically uniform over the whole area.

(ii) During the interval, Z and the lower half of the *Laminosa* Dolomites, dolomitization was much more extensive in Eastern and North-Western Gower than in South-Western Gower; this difference accords with the view of Prof. Skeats¹ and others that within certain limits shallowness favours dolomitization, for the Avonian sea is known from other evidence to have shallowed northwards in the South-Western Province.

(iii) The facies of sedimentation of the rest of C_1 was practically uniform over the whole area, though the ratio of the two rock-types formed, dolomite and oolite, varied.

(iv) The fact that lagoon-conditions were established at the base of C_2 in Eastern and North-Western Gower but not in South-Western Gower was doubtless due to the southward deepening of the Avonian sea. At this period, however, part of the Eastern

¹ Q. J. G. S. vol. lxi (1905) pp. 133-38.

District approximated to the conditions of South-Western Gower: for at Longland Bay the base of C_2 is not a *Modiola* phase, but a group of dolomites representing standard gasteropod- and other limestones.

(v) In the character of C_2 and S_1 Eastern Gower definitely links itself with South-Western Gower, and dissociates itself from North-Western Gower. In one respect, namely in an absence of 'contemporaneous' dolomite and chert from the last-named district, this difference, between Eastern and South-Western Gower on the one hand and North-Western Gower on the other, persists to the top of D. The difference is marked in the case of dolomitization, because outside North-Western Gower many horizons have been affected by this alteration. As a similar difference between southern and northern outcrops is almost as noticeable in South Pembrokeshire; and as, further, outcrops to the north of either Gower or South Pembrokeshire are largely if not entirely devoid of 'contemporaneous' dolomitization at horizons above C_1 , it is evident that a widespread influence inhibiting such dolomitization—to which alteration the conditions of deposition at many horizons would have been favourable—existed in the coastal waters of the Upper Avonian sea. In Gower this influence first made itself felt in C_2 ; its nature, however, is unknown.

(vi) During the interval, S_2 and D_1 , the facies of sedimentation in different parts of the area varied little except in dolomitization, as mentioned above; at the top of S_2 , lagoon-conditions were but occasionally established in South-Western Gower.

(vii) Though the greater part of the 'black lias,' D_{2-3} , of the Eastern District doubtless corresponds to the same group of the South-Western, some of it, Dr. Vaughan concludes (p. 552, table), has been deposited during the formation of part of the comparatively pure D_2 limestones which underlie D_{2-3} in the latter district. That is, argillaceous limestones and shales in Eastern Gower are the contemporaries of pure limestones in South-Western Gower. D_4 is present in North-Western Gower also, and reappears in force, as a mass of pure limestones, along the North Crop of the coalfield at Mynydd-y-gareg near Kidwelly,¹ 12 miles farther north still. The occurrence of the pure limestone facies so far north would point to the existence of a marked embayment in the Avonian coast hereabouts, on the supposition that D_{2-3} was deposited in waters essentially shallower than those of D_2 in its 'pure' development.² It is more probable that the difference between D_{2-3} and D_2 is primarily independent of depth. For, north-east of Mynydd-y-gareg, along the North Crop between that place and Garn-bwli (Carmarthenshire), the pure limestone facies of D_2 gives place laterally to dark sandy limestones with chert,³ that is, to

¹ 'The Country around Carmarthen' Mem. Geol. Surv. 1909, p. 79.

² This is the view of Dr. Vaughan, Rep. Brit. Assoc. (Winnipeg) 1909, p. 190.

³ T. C. Cantrill, 'The Country around Carmarthen' Mem. Geol. Surv. 1909, pp. 76-77. He tells me that this facies continues for a great distance east of Garn-bwli.

beds that differ lithologically from the 'black lias' at the same horizon in being sandy instead of argillaceous. This lateral passage is observable practically due north of the change from pure limestone to 'black lias' in Gower. In other words, sand was being deposited in the north contemporaneously with mud at a considerable distance to the south; while at a short distance to the west of both places the water was clear. Such conditions are most readily explained on the supposition that the region lay at the western margin of the sediment brought into the Avonian sea by a southward-flowing river.

(viii) The deposition of 'black lias' in South-Western Gower above D_2 , while it persisted in the Eastern District, points to an extension of the muddy waters described in the preceding paragraph.

(ix) This widespread and progressive incoming of argillaceous rocks and sponge-cherts towards the end of D probably foreshadowed the marked advent of shales and cherts, spicular or radiolarian, at the commencement of P. The P-shales, therefore, may, also, have derived much of their material in the area here described from a southward-flowing river. With this view their general character and the lateral passage¹ northwards of some part of them, or of the overlying shales, into the comparatively-coarse basal Millstone Grit of the North Crop are in full agreement.

Conclusions.

(1) In the facies of Z and the lower half of the *Laminosa* Dolomites the Eastern District resembles the North-Western, and both differ from South-Western Gower; but in that of the rest of C_1 there is no marked difference between the districts: while, in the development shown by C_2 , S, and D_1 , the Eastern District links itself with the South-Western and both differ from North-Western Gower.

(2) These resemblances and differences are probably attributable to the fact that the depth of the Avonian sea and the influence of the Avonian coast differed in the several districts; for similar differences are observable in Pembrokeshire also, in different outcrops bearing the same relations to the probable position of the Avonian coast as do the parallel districts of Gower.

(3) The evidence afforded by the thicknesses of the zones and subzones in the different districts (Table I, facing p. 505) leads to the same grouping as the evidence summarized in (1). Thus:—

Zone Z and the *Laminosa* Dolomites together are at least 50 per cent. thicker in the South-Western than in the Eastern and North-Western Districts;

The *Caninia* Oolite and C_2 , though variable, show no marked increase in any direction;

Zone S is 50 per cent. thicker in the South-Western and Eastern than in the North-Western District; and

Zone D is thicker, probably in even greater ratio, in the South-Western and Eastern than in the North-Western District.

¹ A. Strahan, Swansea Memoir, p. 29.

In passing, it may be remarked that the sequence, Z-D inclusive, is thicker collectively in South-Western Gower than at any other place in the South-Western Province where it has been measured; and that D is more completely developed in this district, also, than anywhere else in the same province.

Earth-Movements.

(4) The thicknesses, when combined with the depths at which the deposits have been formed, in the manner outlined on p. 537, paragraph (4), would enable us to compare the magnitudes in the different districts of each earth-movement that accompanied deposition. In our general ignorance of exact depths, this can be done only for the summations of the various movements between those horizons that appear to have been deposited at a uniform depth in all districts. Of such horizons the most useful are the lagoon-phases, on account of their well-defined character and the certainty that differences between different districts (as regards average depth of deposition of any phase) are negligible, the range of depth in a lagoon-area being small. To apply the method:—

The first datum-horizon is Km, which is probably developed throughout Gower. By the time that the next datum-horizon, the base of C_2 ,¹ was initiated, Km had been buried more deeply in the South-Western District than in North-Western or Eastern Gower, by an amount equal to the difference between the thickness of the deposits formed in the interval in the first district and that formed in the others; this difference, on the justifiable assumption that K was no thicker in the North-Western and Eastern districts than in the South-Western, was at least 300 feet (see Table I, facing p. 505). In the interval between Km and the base of C_2 , therefore, depression was greater, by at least 300 feet, in South-Western than in North-Western or Eastern Gower.

In the same way, by the time that the next datum-horizon, the top of S_2 , was reached, the base of C_2 had been buried at least 400 feet deeper in the South-Western (see, however, previous footnote) and Eastern Districts than in North-Western Gower. And, if the radiolarian cherts at the base of P are a lagoon-phase and may, therefore, be regarded as a datum-horizon, by the time that their deposition was commenced the top of S_2 had been still more deeply buried in South-Western and Eastern than in North-Western Gower.

(5) From the preceding conclusions it is seen that the earth-movements that accompanied deposition were differential and that the regions of deeper water were the more depressed.

On the assumption² that the differential movements were widespread tilts, not irregular faulting or folding, the axis about which

¹ In South-Western Gower the base of C_2 is not a lagoon-phase, but the fact that it is there represented by deeper-water deposits only emphasizes the conclusion that follows.

² That assumption is justified, in the case of the Upper Avonian movements, by the fact that throughout those movements the Eastern and South-Western Districts were depressed *pari passu*.

hinged the total movement during the first interval, Km-C₁, that is, the Lower Avonian (see § VI) movement, was approximately parallel to the line connecting the Eastern and North-Western facies. (This line is given with sufficient accuracy by the line of outcrop between the Eastern and the North-Western Districts). Similarly the axis during the two succeeding intervals, C₂-top of S₂, and D₁-D₂₋₃ (that is, the axis of the Upper Avonian movements), was approximately parallel to the line connecting the Eastern and South-Western facies. Although this line was subsequently distorted beyond determination by pre-Triassic movements, it was evidently not parallel with the former axis. The axis of the Upper Avonian movements differed in direction from that of the Lower Avonian movement.

Summary of §§ III, IV, & V.

III. Pseudobreccias resemble true breccias outwardly, but their structure has resulted from patchy recrystallization of calcite in foraminiferal calcareous 'mud.'

IV. Lagoon-phases are rock-groups which have been deposited in extremely-shallow, coastal waters; they are characterized by an exceedingly-fine grain and peculiar faunas.

The radiolarian cherts of Gower and of the Culm of the West of England are concluded to be lagoon-deposits.

Lagoon-phases have marked the commencement of cycles of earth-movement.

V. Three bathymetrical cycles can be traced in the formation of the Carboniferous Limestone Series (Zones K-D) of Gower.

These cycles resulted largely from earth-movements, in which depression, on the whole, by far exceeded uplift.

Depression was greater in the seaward than in the landward regions; and the direction of the hinge of the Lower Avonian movement differed from that of the Upper Avonian movements.

It is a pleasure to recall the assistance which I have received in various ways from my colleagues on the Geological Survey and from other geologists. To Dr. J. J. H. Teall I am indebted for permission to publish some observations made in the course of my official work in Pembrokeshire. Dr. A. Strahan I have to thank for placing at my disposal the information obtained during the geological survey of Gower. Reference has been made on previous pages to Mr. T. C. Cantrill, who worked with me for a time on the ground and has put his detailed knowledge of the Carboniferous of adjacent areas at my disposal; Prof. L. Cayeux; Mr. H. Dewey; Mr. T. C. Hall; Mr. J. G. Hamling; Mr. J. A. Howe; Dr. W. Pollard; and Dr. R. L. Sherlock. To all these workers my heartiest thanks are due.

VI. NOTES ON THE DELIMITATION OF THE UPPER FROM THE LOWER AVONIAN IN THE SOUTH-WESTERN PROVINCE.

(1) Stratigraphical Considerations [E. E. L. D.].

The earth-movements discussed in the previous section (pp. 536-37), and the faunal, lithological and stratigraphical changes which are connected with them, afford a basis for a delimitation of the divisions of the Avonian in the South-Western Province. In applying this principle, it is advisable to include a lagoon-phase in the base of a division rather than to associate it with the underlying rocks, because such phases, in general, mark the commencement of cycles of earth-movement.

For these reasons the base of the *Modiola* phase at the base of ($C_2 + S_1$), that is, the divisional line between C_1 and C_2 , is the appropriate horizon at which to separate, as is necessary, an Upper from a Lower Avonian.¹ It is the more suitable because it is sharply defined lithologically and, in places, is emphasized by unconformity with the underlying rocks. And this unconformity, there is reason to believe, is the commencement of the transgression that has brought the Upper Avonian to rest upon far older rocks in the northern part of the South Wales area.

In those places, however, which represent the deeper parts of the Avonian sea, not only does the unconformity disappear, but also, as the distance from the Avonian coast increases, the *Modiola* phase itself is lost, and ultimately the base of C_2 may become untraceable by lithological means. Faunal methods, however, are available, such as is afforded by the incoming of *Cyathophyllum* ϕ Vaughan, though, as will be seen from the following note by Dr. Vaughan, the extreme base of C_2 does not everywhere correspond to the maximum of the change of fauna in passing from the Lower to the Upper Avonian.

(2) The Faunal Relations of C_2 to the Beds below and above [A. V.].

Wherever, as at Burrington, etc., C_1 is composed of limestones of standard type, the C_1 fauna is a continuation of the γ facies, that is, an extension and amplification of the Lower Avonian fauna, and, locally, this fauna is persistent into the base of C_2 (as here defined on lithological grounds).

On the other hand, the main C_2 fauna exhibits the earliest establishment of genera and gentes which are especially characteristic of Upper Avonian time. For example:—

Corals:—

The gens of *Cyathophyllum* aff. *murchisoni*.

The genera *Campophyllum*, *Diphyphyllum*, and *Clisiophyllum*.

¹ The terms Upper and Lower Avonian are used provisionally, pending a general discussion of the Avonian of this and other areas, in some of which other terms, such as Viséan and Tournaisian, are already in use.

Brachiopods:—

The following gentes and groups:—

Productus corrugato-hemisphericus.
Productus punctatus.
Productus elegans.

Productus pustulosus.
Productus sublævis.
Spirifer aff. *bisulcatus*.

The genera *Actinoconchus*, *Seminula*, *Dielasma*.

Furthermore, the C_2 and S_1 faunas are linked by

Cyathophyllum sp.
Cyathophylloid Caninia.
Early *Carcinophylla*.
Megastomatoid *Michelinia*.
Productus corrugatus, mut. ϕ . (nearly
equivalent to *Productus* of S_1).

Chonetes cf. *comoides* (of great size).
Certain large species of *Bellerophon*
and *Euomphalus*.

(In fact, beyond the occurrence of *Lithostrotion* there is frequently no other distinction between C_2 and S_1 .)

Dr. Gosselet in 'L'Ardenne' 1888, chap. xxii, includes the C_2 fauna (the *Productus-sublævis* Beds) with the typical S fauna above in a single 'assise,' the 'Calcaire des Ardennes.' The transition from the *Productus-sublævis* fauna into the basal *Pr.-corrugatus* fauna is, in fact, exactly that which marks the passage of C_2 into S_1 in the South-Western Province generally.

Hence it will be entirely in accordance with the faunal evidence to draw the line of division between the Upper and the Lower Avonian at the bottom of the main part of C_2 . The divide will therefore lie within C_2 , as at present defined, above the lowest part of the standard limestones, which contains a recurrent C_1 fauna.

VII. FAUNAL LISTS [A. V.].

Note.—Here and in the Palæontological Section, repetition of full references is avoided by the following abbreviations:—

'Bristol Paper' = Q. J. G. S. vol. lxi (1905) pp. 181-305 & pls. xxii-xxix—
'Palæontological Sequence in the Carboniferous Limestone of the Bristol Area,' by A. Vaughan.

'Rush Paper' = Q. J. G. S. vol. lxii (1906) pp. 275-322 & pls. xxix-xxx—
'Carboniferous Rocks at Rush (County Dublin),' by C. A. Matley & A. Vaughan.

'Loughshinny Paper' = Q. J. G. S. vol. lxiv (1908) pp. 413-72 & pls. xlix-l
—'Carboniferous Rocks at Loughshinny (County Dublin),' by C. A. Matley & A. Vaughan.

'Davidson' = 'Monogr. Brit. Foss. Brach.' vol. ii (1858-63), Palæont. Soc.

Except in the case of D_{2-3} , as explained below, no attempt was made to draw up an exhaustive list at any level; immediately the zonal position had been placed beyond doubt, further search was abandoned.

LOWER AVONIAN.

CLEISTOPORA ZONE (K).

South-Western District:

Km:

Only the base of K is exposed near Rhossili.

Brachiopods:—

Chonetes cf. *crassistria* Vaughan¹ (non M'Coy).

Eumetria aff. *carbonaria* (Dav.) Vaughan.²

Camarotæchia cf. *mitcheldeanensis* Vaughan.

Lamellibranchs:—

Modioliform species, including *Modiola* cf. *lata*.

Gasteropods:—

A small species of *Bellerophon* is common.

Ostracods.

ZAPHRENTIS ZONE (Z).

Z₁:

Eastern District: Threecliff Bay (indicated by the numeral 4).

South-Western District: Rhossili to Worm's Head (indicated by 1).

Corals:—

Small simple *Zaphrentis*—1 and 4 (rare).

Brachiopods:—

Productus burlingtonensis Hall, Vaughan—1 and 4.

Small *Chonetes*: [*Ch.* cf. *crassistria* predominating]—1 and 4.

Leptæna—4.

Orthotetid—1 and 4.

Spirifer aff. *clathratus* M'Coy, Vaughan—1 and 4 (abundant)

A small *Syringothyris* (cf. *S. typa* Winchell)—4.

Reticularia cf. *lineata*³ (Martin) (Dav.)—1 (rare).

Athyris (*Cliothyris*) cf. *glabristria* (Phill.) (Vaughan)—4 (rare).

Camarotæchia mitcheldeanensis Vaughan—1.

Crinoids:—

Dichocrinus sp.—4. (Determined by Dr. F. A. Bather, F.R.S.)

¹ 'Bristol Paper,' pl. xxvi, fig. 2; also Q. J. G. S. vol. lxiii (1907) p. 458.

² 'Bristol Paper,' p. 302.

³ This species differs very considerably from Martin's figure in the irregular and undulatory concentric wrinkles and in the prominence of the thick underlayer of close-set tubes from which the fringed bands composing the outer layer are formed.

Z_2 :

South-Western District: Rhossili to Worm's Head (indicated by 1).

(This subzone is also exposed at Threecliff Bay in the Eastern District; the beds are, however, here dolomitized, and fossils are indeterminable.)

Corals:—

*Zaphrentis omaliusi*¹ Ed. & H., Carruthers.

*Zaphrentis konincki*² Ed. & H., Carruthers.

Caninia cornucopiæ Mich., Carruthers.

All three are common in 1.

Brachiopods:—

The usual Z_2 assemblage: no list was drawn up.

Horizon γ .

South-Western District: Rhossili to Worm's Head (denoted by 1).

Corals: same as under Z_2 , together with:—

Michelinia tenuisepta Ed. & H. and de Kon., *non* (Phill.).

Caninia cylindrica (Seouler in M'Coy) Vaughan.³

Brachiopods:—

Productus cf. *pustulosus*⁴ Phill. (and cf. *Pr. christiani* de Kon.).

Semireticulate *Productus*.

Leptæna.

Orthotetid.

Spirifer aff. *clathratus* M'Coy, Vaughan.

Spirifer cinctus de Kon.⁵

Syringothyris cuspidata (Martin).

Syringothyris cf. *laminosa* (M'Coy) (Dav. pars), Vaughan.⁶

¹ = *Z. aff. phillipsi* Ed. & H., Vaughan: as described in 'Bristol Paper,' p. 270.

² = *Z. aff. cornucopiæ* (Mich.) Ed. & H.: as identified in 'Bristol Paper.'

³ *Caninia cylindrica*, as employed in the 'Bristol Paper,' covered all the Caninias from the entrance of the genus at the top of Z_2 to the maximum of '*C. cylindrica*, mut. S₁,' in the Lower *Seminula* Zone. These would now be differentiated into *C. cornucopiæ*, *C. patula*, and *C. cylindrica* with variants of each; at the time of our work in Gower (in 1905), however, all these forms were lumped under *C. cylindrica*.

⁴ A common form at several points of the South-Western Province and first appearing in γ : maximum at the top of C₁ and in C₂. The umbonal region is strongly wrinkled and pustulose, but the skirt is often almost smooth (cf. *Productus christiani*, which is abundant in C₂ of Belgium).

⁵ *Spirifer cinctus* Fischer is an Upper D form and a near ally of *Sp. planicosta* (M'Coy): the form cited above is referred to as *Spirifer konincki* Dewalque by Belgian writers, and is an important zonal index of Z-C.

⁶ The form here indicated accords with *Syringothyris* in its general build and area, but is more closely allied to *Spiriferina* in its umbonal plates. All Carboniferous *Spiriferinas* have a primitive syrinx which, however, remains undeveloped and is buried in, and filled in by, the umbonal callus. In the species here recorded, the syrinx is quite obvious, although confined to the apical portion of the delthyrium.

SYRINGOTHYRIS ZONE (C).

Eastern District: Threecliff Bay (4) and Longland Bay (6).
South-Western District:—Cliff west of Overton.

C₁:

Composed of the *laminosa* dolomite and *Caninia*-Oolite.

Fossils are very scarce; narrow-tubed *Syringopora*, large-celled *Michelinia*, small *Zaphrentis*, large *Caninia*, and an early Clisio-phyllid were, however, recorded from various places in Eastern and South-Western Gower, as well as a band of *Chonetes* cf. *comoides* and of *Bellerophon* in the *Caninia* Oolite of Longland Bay.

C₂:

Lowest part of the standard limestones.

Zaphrentis omaliusi Ed. & H., Carruthers—4 and 6.

Zaphrentis konincki Ed. & H., Carruthers—4 and 6.

Caninia cylindrica (Sculer in M'Coy) Vaughan—abundant in 4.

Syringothyris cuspidata (Martin)—4 and 6.

Syringothyris cf. *laminosa* Vaughan—rare in 4 and 6.

Cyathophyllum ♂ Vaughan (small mut.¹)—4 and 6.

UPPER AVONIAN.²

C₂:

Main part of the standard limestones.

Corals:—

Syringopora cf. *geniculata* Phill., Nicholson—4 and 6.

Syringopora cf. *reticulata*—4.

*Michelinia grandis*³ M'Coy—4 and 6.

Zaphrentis konincki, mut. C₂⁴—6 (nearly at the top).

Caninia cylindrica (Sculer in M'Coy) Vaughan—4 and 6.

Cyathophyllum ♂ Vaughan—abundant in 4 and 6.

Diphyphyllid *Lithostrotion*—Overton.

Brachiopods:—

Chonetes cf. *comoides* (Sow.).

Rhipidomella aff. *micelini*—6.

Spirifer cf. *trigonalis*⁵—6.

Syringothyris cuspidata—4 and 6.

Seminula sp.⁶ [cf. *S. ambigua* (Sow.)]—6.

¹ *Cyathophyllum* ♂ is not established until C₂; the small mutation is, therefore, a descending mutation, that is, a previously established member of the gens. It obeys the general law—'earlier and smaller.'

² The reasons for starting the Upper Avonian at this level are set out in § VI.

³ = *Michelinia* cf. *megastoma* of my earlier lists.

⁴ *Zaphrentis konincki* is a Z₂ to γ index; this mutation is, therefore, an ascending mutation; that is, a subsequently-established member of the gens. It is larger, and has longer intermediates, in accordance with the general rule. [See my note in R. G. Carruthers's paper, Geol. Mag. dec. 5, vol. v (1908) p. 70.]

⁵ Small, sharply-ribbed form with a strong angular fold, somewhat produced.

⁶ Deeply sinuate form, recorded as *Seminula ambigua* by Dr. T. F. Sibly, from Weston and the Mendips.

⁵ & ⁶ These forms characterize the C-S level at widely separate localities in England; they are, therefore, important diagnostic forms.

Gasteropods:—

Large species of *Bellerophon* and *Euomphalus* are abundant.

Cephalopods:—

A large species of an *Orthocerate* is common.

SEMINULA ZONE (S).

S₁:

Eastern District: Threecliff Bay (4) and the cliff-section from Longland to Mumbles.

South-Western District: Overton (denoted by 2).

Corals:—

Michelinia grandis (M'Coy)—2.

Caninia (Cyathophylloid) *bristolensis*¹ Vaughan—2, 4 and 6.

Cyathophyllum ϕ Vaughan—2 and 6.

Lithostrotion cf. *irregulare* (Phill.) Ed. & H.—4, 6 (only at the base).²

Lithostrotion martini Ed. & H., and variants—2, 4, 6.

Diphyphyllid³—6 (only at the base).

*Heterophyllia*⁴ sp.—2.

Brachiopods:—

Productus aff. *semireticulatus*, mut. S₁ Vaughan—2, with fragments of a large *Productus*.

Productus cf. *elegans* M'Coy—2 (abundant).

Chonetes cf. *comoides* (Sow.)—(persists at Overton as long as *Caninia*).

Seminula ficoides Vaughan, and varieties—2, 6 (very abundant).

Gasteropods:—

Bellerophon, *Euomphalus*, and *Loxonema* abundant—2, 6.

Bryozoa:—

Fenestellids and *Heterotrypa* cf. *tumida* (Phill.) crowd the shaly partings at Overton (2).

Trilobites:—A Phillipsid is common—2.

[The resemblance of S₁ at Overton to S₁ of the Avon is very striking, for the shaly partings are crowded with the same species of bryozoa enmeshed in a network of spines broken from the index *Productus* and accompanied by the same small trilobite; while the limestones themselves contain at both places *Caninia bristolensis* in abundance.]

S₁ of Gower only differs from S₁ of Bristol in that, in Gower, *Cyathophyllum* ϕ persists into S₁ and *Seminula* is established later.

¹ Proc. Bristol Nat. Soc. ser. 3, vol. x (1903) p. 103 & pl. i, fig. 4; = *Caninia cylindrica*, mut. S₁ of 'Bristol Paper.'

² This species differs from *L. irregulare* of the D levels in its strikingly ramulose habit. The beds in which it occurs have a typical C-S fauna, and were formerly considered by me to mark the top of C₂; it is somewhat more convenient to consider them as the base of S₁. In any case, C₂ and S₁ are faunally continuous both in (2) and (6).

³ Diphyphyllid *Lithostrotions* are not uncommon in C-S, and again are very common in Upper D, but are apparently absent from S proper, where *Lithostrotion* reaches its maximum.

⁴ Collected by the late Dr. W. B. Gubbin (Bristol Nat. Soc. *op. jam cit.*).

S_2 :

Eastern District: Longland to Mumbles section (6); Bishopston Valley (8).

South-Western District: Overton (2).

The Lower (cherty) part abounds in silicified clusters of a large *Lithostrotion*—*L. affine* (Flem.) Ed. & H. *Seminula* and *Productus* fill recurrent bands. Gasteropods are numerous.

The Upper (oolitic) part contains a fauna characteristic of the S_2 subzone throughout the South-Western Province. Bands crowded with *Seminula*, *Lithostrotion*, or *Productus corrugato-hemisphericus* recur again and again.

Fauna of Upper S_2 :

Corals:—

Syringopora—6.

Alveolites—6.

Lithostrotion martini Ed. & H., and varieties.—2, 6, 8 (very abundant).

Carcinophyllum θ Vaughan—6, 8.

Brachiopods:—

Productus hemisphericus Sow.—2, 6, 8.

Productus corrugato-hemisphericus including '*P. cora*, mut. S_2 ,' Vaughan—2, 6, 8 (very abundant).

Papilionaceous *Chonetes*—6, 8.

Seminula ficoides Vaughan and varieties.—2, 6, 8 (very abundant).

Seminula cf. *ambigua* Sow., an S_2 form—2. (Collected by the late Dr. W. B. Gubbin; see 'Bristol Paper,' p. 235.)

Gasteropods:—*Bellerophon*—6.

DIBUNOPHYLLUM ZONE (D).

D_1 :

Eastern District: Pwll-du (5); Limeslade Bay (6), and Mumbles Head (6 a); southern end of Colts Hill (Oystermouth); Bishopston Valley (8).

South-Western District: Port Eynon (3).

Corals:—

Syringopora geniculata Ed. & H.—3.

Alveolites septosa (Flem.) Ed. & H.—5.

Zaphrentis ambigua Carruthers (?)—a unique specimen from (3).

Campophyllum aff. *murchisoni* Ed. & H.—6, 6 a, 8, 3.

Clisiophylloid *Campophyllum*—5, 3.

Cyathophyllum murchisoni Ed. & H.—5, 3, 8, 6 a (extremely abundant).

Lithostrotion martini Ed. & H., and vars.—5.

Lithostrotion junceum (Flem.).—6 a.

Koninekophyllid—6.

Diphyphylloid *Cyathophyllum*—3, 6 a.

Cyathophylloid *Aulophyllum*—8.

Carcinophyllum θ Vaughan and a variety convergent with *Dibunophyllum* θ —5, 6 a.

Dibunophyllum θ Vaughan, and *D. \phi* Vaughan—6, 6 a, 5, 3, 8.

Dibunophyllum aff. ψ Vaughan—3.

Brachiopods:—

Productus hemisphericus Sow.—5, 6 a, 8 (very abundant).

Productus giganteus (Mart.)—5, 3, 6 a, 8.

Chonetes (*Daviesiella*) aff. *comoides* (Sow) Vaughan—5, 8. (See p. 561.)

Athyris cf. *expansa* (Phill.), mut. D₁—6 a.

The above fauna is characteristic of D₁ throughout the South-Western Province.

D₁₋₂:

The uppermost beds of the Pwll-du section contain *Dibunophyllum* aff. ψ Vaughan and *Lithostroton portlocki* (Bronn) Ed. & H. (both of which are D₂ forms), in association with the normal D₁ assemblage.

D₂:

South-Western District: Well exposed in quarries at Port Eynon.

Fauna:—

Corals:—

A narrow *Syringopora*.

Michelinia tenuisepta (Phill.) (near the top).

Alveolites (?).

Zaphrentid and 'Calophyllid.'

Cyathophyllum muchisoni (?).

Lithostroton irregulare (Phill.) (abundant in the lower part).

Lithostroton junceum (Flem.) Ed. & H.

Petalaxis portlocki Ed. & H. (common near the top).

Dibunophyllum ψ Vaughan (not common).

Brachiopods:—

Latissimoid *Productus* (common).

Productus cf. *concinus* Sow. (broad form).

Productus cf. *muricatus* Phill.

Productus scabriculo-costatus (abundant at the very top).

Productus striato-corrugatus.

Papilionaceous *Chonetes* (small form).

Leptæna cf. *distorta* Sow.

Spirifer planicosta (McCoy).

Martinia cf. *ovalis* (Phill.).

Martinia glabra (Martin).

Bryozoa:—

Fenestella sp. nov.

Note.—Although *Lonsdalia* has not yet been recorded with certainty, the above fauna may be unhesitatingly regarded as characteristic of D₂.

The D₂₋₃ Phase:

Eastern District { Oyster-mouth. { Top of quarry on north of Colts Hill (a).
 { 'Black-lia' Quarry (b).
 { Bishopston (road-section in Rottenstones) (c).
South-Western District: Port Eynon, above D₂ series) (d).

Since this phase was unknown to me when writing the 'Bristol

Paper,' the fauna recorded below is, in the main, new to the South-Western Province.¹ I have consequently devoted the Palæontological Section, almost in its entirety, to descriptions of the most interesting components of this fauna. Reference must, therefore, be made to that section for explanations of the fossil names which compose the following list.

Corals:—

- Michelinia tenuisepta* (Phill.) (a, b).
Amplexus (?) sp. (b).
Zaphrentis enniskilleni Ed. & H. (a, b).
Zaphrentis oystermouthensis, sp. nov. (a, b); common.
Densiphyllum cf. *charlestonense* Thomson (b).
Caninia aff. *cornucopiæ*, mut. D₂₋₃ (a, b).
Caninia (?) sp. (a).
Cyathaxonid (cf. *Cyathaxonia* aff. *costata* M'Coy, Vaughan: figured in 'Rush Paper,' pl. xxix, fig. 5) (a).

Brachiopods:—

- Productus longispinus* Sow., var. (b, c, d).
Productus sulcatus Sow. (a, b, c).
Productus scabricuto-costatus (b, c).
Productus corrugatus M'Coy (b, c).
Productus aff. *hemisphericus* Sow. (b).
Productus edelburgensis Phill. (b, c).
Productus margaritaceus Phill. (b).
Productus punctatus (Mart.) (b).
Productus elegans M'Coy (b, c).
Chonetes sp. (b, c).
Papilionaceous *Chonetes* (small form) (b).
'*Orthotetes*' cf. *crenistris* (Phill.) (b, c).
Schizophoria cf. *keyserlingiana* (Dav.) (b); form intermediate between *Schizophoria* and *Rhipidomella* (b).
Spirifer bisulcatus Sow., var. *oystermouthensis*, nov. (a, b, c, d); very abundant.
Spirifer near *increhescens* Hall (b).
Spirifer wickensis, sp. nov. (b).
Spiriferina insculpta (Phill.) (b, c).
Reticularia lineata (Mart.), globular variety (b, c).
Martinia glabra (Mart.) (a, b, c, d).
Athyris (*Chiothyris*) *globularis* (Phill.), and a circular variety.
Camarophoria cf. *crumena* Dav. (non Mart.) (b).
*Camarotoechia*² aff. *pleurodon* (Phill.) (b).

Bryozoa:—

- A *Fenestella* and a Monticuliporoid seem both to be distinctive (b, c).

¹ A return visit, in company with Mr. W. H. Wickes, was therefore undertaken in order to make the list as exhaustive as possible.

² The rubbing-down of this, and of several typical specimens of *Rhynchonella pleurodon* Phill. from the Upper Devonian of Wetton, has shown that this species agrees with *Camarotoechia* in possessing a double and camerate septum in the brachial valve (see 'Bristol Paper,' p. 302); *Pugnax* has no septum in this valve.

Trilobites:

The types of *Griffithides barkei* H. Woodw. (c) (Geol. Mag., 1902, p. 484) and *Gr. glaber* H. Woodw. (b) ('Mon. Brit. Carb. Tril.' pt. ii, Pal. Soc. 1884, p. 40).

Note on the Bishopston Fauna.

The only noticeable differences that distinguish the Bishopston fauna from that of Oystermouth are the absence of Zaphrentids at Bishopston and the predominance there of *Spirifer trigonalis* (Mart.) over *Sp. bisulcatus* Sow. The large number of species common to the two localities demonstrates approximate equivalence of level, and also the Upper D age of both faunas.

Note on the Horizon of the D₂₋₃ Beds of Gower.¹

[Evidence and inference are here set out under localities.]

(i) Bishopston:—

- (1) The Rottenstones represent the top of the D₂₋₃ series;
- (2) The shales immediately above the Rottenstones, and interbedded with the radiolarian cherts, have yielded *Posidoniella laevis*, *Solenomorpha minor*, and *Glyphioceras bilingue*—fossils which Dr. Wheelton Hind identified and considers to indicate a level 'low down in the Pendleside Series.'

It is also important to note that, in shales considerably above the level of those in the Bishopston cutting, *Glyphioceras spirale* (Phill.) is recorded by the Geological Survey.²

(ii) The two quarries on Colts Hill, Oystermouth.—The Southern Quarry is in D₁, and there is a continuous series of some 120 feet of massive limestone between it and the base of D₂₋₃ in the Northern Quarry. This limestone is very poorly fossiliferous, although it contains a bed crowded with a latissimoid variant of *Productus hemisphericus* Sow. near the base of the Northern Quarry.

Allowing a certain weight to this form and to the small thickness of the limestone series between the quarries, it is evident that the base of D₂₋₃ cannot be of later date than D₂ age.

The fauna of the base of the D₂₋₃ series in the Colts Hill Quarry cannot be distinguished from that of the Oystermouth Quarry, except for the presence of '*Cyathaxonia*' in the first-named locality. This coral occurs with *Zaphrentis* nr. *oystermouthensis* in the *Cyathaxonia* Beds of the Rush and Loughshinny sequence in County Dublin, and I have already suggested ('Loughshinny Paper' p. 444) that these last-named beds are the equivalent of part of D₂₋₃ of Gower and of part of D₂ of the Bristol Area.

(iii) Port Eynon.—Succeeding the massive and typical D₂ limestones at Port Eynon are thinly-bedded limestones and shales of

¹ See also § II, p. 495.

² Swansea Memoir, p. 25.

'black lias' rock-facies, which are poorly exposed on the foreshore. From these beds we obtained *Productus longispinus* Sow., var., *Spirifer bisulcatus* Sow., var., and *Martinia glabra* (Mart.), var., identical in form with the same varieties from Oystermouth. Hence it is clear that the D_{2-3} fauna is, in part at least, later than that of the D_2 massive limestones.

It is also interesting to observe that *Productus scabriculo-costatus* does not come in at Port Eynon until the very top of the D_2 limestones, and that, in the Bristol Area on the east and at Kidwelly and Pendine on the west, the same form is first met with above the typical D_2 and immediately below the Millstone Grit.

(iv) From the shales at the 'paint-mine,' near Port Eynon, we obtained the following fossils (identified by Dr. Wheelton Hind):—

<i>Posidonomya becheri</i> (Bronn).		<i>Glyphioceras reticulatum</i> (Phill.).
<i>Posidoniella laevis</i> (Brown)?		<i>Glyphioceras bilingue</i> (Salter):

Dr. Hind suggests that, here, we are 'fairly low down in the Pendleside Series.' Since these shales are faulted against the limestone, it is impossible to ascertain their precise position stratigraphically; it can, however, be safely assumed that they lie above the 'black lias' of the foreshore at Port Eynon.

This record, taken in conjunction with the fossils cited from the shales at Bishopston, demonstrates that a Pendleside Phase occurs above the D_{2-3} Phase, just as it does above the *Cyathaxonia* Beds (D_3) of County Dublin.

Table correlating the Upper Avonian of Gower, Bristol, and Rush (County Dublin).

BRISTOL.	GOWER.		RUSH AND LOUGHSHINNY (Co. DUBLIN).
	Pendleside.		P.
'Millstone Grit.'	Port Eynon.	D_{2-3} .	
D_2 .		D_{2-3} .	D_3 (' <i>Cyathaxonia</i> ' Beds).
		D_{1-2} .	D_{1-2} (' <i>Dibunophyllum</i> ' Beds).
D_1 .		D_1 .	Unknown.
		Oystermouth.	

The correlation of the Bristol and Gower sequences with those of Northumberland, Yorkshire, North Wales, the Midlands, etc., is given in the British Association Report, 1909 (Winnipeg), on Lower Carboniferous Zones, p. 187.

VIII. PALÆONTOLOGICAL NOTES [A. V.].

(A) CORALS.

TABULATA.

Syringopora.

SYRINGOPORA cf. *GENICULATA* Phill., Nicholson, from C_2 (main part).

This is the dominant *Syringopora* of C_2 , Gower, and it there takes the place of *Syringopora* cf. *reticulata*, which is the commonest form in C_2 - S_1 of the Bristol area.

Description.—Broad and ramulose, with strongly thickened tubes and not numerous connectors. Very numerous, thick, septal spines project from the wall, and almost or quite penetrate the very thick lining of stereoplasm. (The abundance and strength of the septal spines causes a misleading resemblance, in cross sections, to *Lithostrotion junceum*.)

Tabulæ deeply dependent and distant.

Comparison.—*S. geniculata* Phill., Nicholson, 'Tabulate Corals' 1879, pl. x, figs. 4, 4 a, & 4 b, agrees closely with our species, from which it differs only in degree, namely, it has shorter septal spines, less widely-spaced tabulæ, and is far less ramulose.

Nicholson's conception of *S. geniculata* Phill. differs from the type-figure¹ in the scarcity of connectors, but agrees with Phillips's definition of the species in respect to the ramulose habit ('radiating, often flexuous, branching, round tubes'—Phill. *op. cit.* p. 201).

On the other hand, the accepted connotation of the specific name *geniculata* (compare Ed. & H. 'Monogr. Brit. Foss. Cor.' Pal. Soc. 1852, pl. xlv, figs. 2 & 2 a) disagrees with Phillips's figure in the absence of radiation and branching.

The type of Phillips's species being lost, there seems little hope of obtaining a clear conception of *S. geniculata* from his definition and figure.

ZAPHRENTIDS.

Zaphrentis.

ZAPHRENTIS OYSTERMOUTHENSIS, sp. nov. (Pl. XL, figs. 1 a-1 e.)

Compare *Zaphrentis* aff. *enniskilleni* Ed. & H., Vaughan in 'Rush Paper,' pl. xxix, fig. 2. See also 'Loughshiun Paper' p. 457.

Description.—Form conical and as a rule continuously cornute, 4 to 5 cm. in length. (Fig. 1 a illustrates the less common form, which is only bent at the tip.)

External surface practically smooth, except for fine annular striæ and flattened rugæ.

In the calyx :—Fossula on the concave side, very deep, and occupying more than half the diameter of the coral.

¹ 'Geol. Yorkshire' pt. ii (1836) pl. ii, fig. 1.

Major septa very strongly thickened and frequently coalescent; minor septa only developed in the calyx.

Horizontal sections.—In the young stage the horizontal section of the fossula is regularly and broadly oval, and bisected by a strong fossular septum (Pl. XL, fig. 1 b).

Fig. 9.—*Zaphrentis oystermouthensis*, sp. nov.: section below the calyx, magnified 2 diameters. (Specimen figured in Pl. XL, fig. 1 a.)

In the adult, where the septa have ceased to increase, the fossula is broadly open near the wall and narrows to the centre, being bounded by a single septum on either side (Pl. XL, fig. 1 e).

In intermediate stages the appearance of the fossula depends on whether the section intersects a pair of new septa or not; if it does so, the fossula is contracted in the middle, as in *Z. delanouei*; if it does not, the

appearance is much like that of the adult (compare Pl. XL, figs. 1 c & 1 d).

The septal plan has the following characters (see fig. 9, above):—

- (1) A strong thickening of the counter-septum;
- (2) Great variation in length and direction of the major septa;
- (3) A strong convexity of the longer septa (in the adult) to the generative planes (or 'fossular breaks'.)

From (2) and (3) results the characteristic palmate grouping of the septa and the dependent lateral boundaries of the counter series.

Tabulæ (see fig. 10, p. 555).—The tabulæ are broad, strongly depressed in the middle and along the fossula, and gently everted at the wall.

Horizon and localities.—This species is abundant in D_{2-3} of Oystermouth Quarry. A very similar form is rare in the 'Cyathaxonia Beds' of Rush.

Resemblances and differences.—*Zaphrentis delanouei* Ed. & H. is of much smaller average dimensions, and its dominant form is almost purely conical (only the tip being curved), whereas the usual form of *Z. oystermouthensis* is continuously conical.

In *Z. delanouei* the counter (anti-fossular) septa in the adult are regularly radial, uniformly thick, and approximately of equal length; whereas in *Z. oystermouthensis* they present a quadripinnate grouping, and are of very unequal thickness.

Comparison with *Z. enniskilleni* Ed. & H.—The type-specimen of *Z. enniskilleni* Ed. & H., represented in Pl. XL, fig. 2, shows how markedly *Z. oystermouthensis* differs in the greater irregularity and thickening of its septa, although agreeing in the general plan of septation and in the position of the fossula.

Evolution of *Z. oystermouthensis*.—Among the Z_1 forms, which may all be attributed to *Z. delanoui* Ed. & H., is one which agrees very closely with the young stage of *Z. oystermouthensis*

Fig. 10.—*Zaphrentis oystermouthensis*, *sp. nov.*: contoured plan of floor of calyx, magnified 2 diameters, and two sections on the same scale across the calyx. (Specimen figured in Pl. XL, fig. 1 a.)

depicted in fig. 1 b (Pl. XL); if this form represents the ancestral member of the gens of *Z. oystermouthensis*, variation has merely increased the size and emphasized the irregularity of the septa in direction and thickness.

Caninia.

CANINIA aff. CORNUCOPIE Mich., mut. D_{2-3} . (Pl. XL, figs. 3 a-3 c.)

I have already described the characters of this mutation under the generic title of *Amplexi-Zaphrentis*, in the 'Rush Paper' p. 315 & pl. xxix, fig. 7; and also in a note kindly inserted by Mr. Carruthers in his 'Revision of Carb. Corals' Geol. Mag. 1908, p. 169.

Consequently, it is only necessary here to point out the differences between the Upper Avonian form of the South-Western Province and the Lower Avonian species of the same region ($Z_2-\gamma$), an example of which is figured in Pl. XL, fig. 4. These differences are appreciable only in the adult, and by comparison of the average form at the two levels.

The D_{2-3} mutation :—The tabulæ are gently depressed and close-set; the fossular depression starts near the margin. Hence, in a horizontal section, the contours of the tabulæ, between the septa and in the fossular valley, are close-set and nearly parallel.

The thin vesicular wrapping is developed early, and is practically continuous over the whole of the cylindrical portion.

The γ species (= *Caninia cornucopie* Mich., emend. Carruthers) Pl. XL, fig. 4 :—The tabulæ are well spaced, and the fossular depression starts near the centre. Hence, in a horizontal section, the contours of the tabulæ, between the septa and in the fossula, are few and widely spaced.

The thin vesicular wrapping is developed late and, even in the adult, is seldom present over the greater part of the cylindrical portion.

Occurrence in the South-Western Province.—The γ species is doubtless common at most localities at the top of Z_2 and in C_1 , although Burrington (Mendips) and Stackpole Quay (Pembroke) are the only localities which have as yet been thoroughly searched.

At the top of D_2 and in D_{2-3} the upper mutation is usually to be found, and is often abundant. It is almost invariably cylindrical for the greater part of its length, and broader than the '*cornubovis*' variant from the Upper Tournaisian (specimens indistinguishable, externally, from that variant are, however, occasionally found).

In Gower, this mutation occurs in D_2 of Port Eynon and throughout D_{2-3} of North Colts Hill Quarry and of Oystermouth Quarry.

In the Bristol area it is a very rare fossil in D_2 . In the west of the South-Western Province—Horizon ϵ of Ragwen Point near Pendine—specimens are common, and at Bullslaughter Bay, near Boshes-ton (South Pembrokeshire), it is very abundant in D_2 in association with *Glyphioceras sphericum* (Martin) and a small *Cyathaxonia*.

CANINIA (?) sp. (Pl. XL, fig. 5.)

This coral is remarkable for the extent and slope of the tabulæ, for the shortness of the septa, and the practical absence of a fossula. The tabulæ are flat, and extend almost completely across the corallum; they slope very steeply downwards from the convex to the concave side.

Owing to their high dip, a cross-section cuts several tabulæ; and consequently the interseptal spaces, on the concave side, appear to be vesicular.

Above a tabula the septa are very short and amplexoid without any septal break; underneath a tabula, they are elongated on the concave side and short on the convex side with a small fossular gap. Minor septa are only developed in the calyx.

This species appears to be intermediate between *Amplexus* and *Caninia*—in fact an *Amplexus* above the tabulæ and a *Caninia* below: a *Caninia* in form, an *Amplexus* in calyx.

The few specimens that I have seen from Oystermouth were collected by Mr. Underhill from the base of D_{2-3} in North Colts Hill Quarry.

Mr. H. F. Barke, A.I.C., has collected from γ at Burrington a similar coral having the general form, the short septa, the broad steeply-sloping tabulæ, and the inconspicuous fossula of the species under description. The earlier coral is, however, Amplexoid throughout, and its minor septa are practically undeveloped in the calyx.

Amplexus.

AMPLEXUS (?) sp. (Pl. XL, figs. 6 a & 6 b.)

Form cylindro-conical, the junction of the cylindrical and conical portions being abruptly marked by an extremely strong growth-halt and by a change of axial direction.

The epitheca has the strong girth-ridges characteristic of *Amplexus nodulosus* Phill., but there are no spines.

The septa are long, widely spaced, and about 20 in number.

In the figured section (fig. 6 b) the septa extend much farther towards the centre than is the case in the small Amplexids so common at this level in northern localities (*A. nodulosus* Phill., etc.), and they again differ very remarkably in their inequality. It is, however, to be noted that sections differently situated with regard to the tabulæ exhibit a series of equal septa which fall short of the centre, and only differ from those of *A. nodulosus* in their greater length.

The tabulæ are strong, distant, tall arches, flattened or gently convex above and sharply bent down near the wall.

The species is rare in D_{2-3} of Oystermouth Quarry, where it takes the place of *A. nodulosus*, which is common at the same level in other localities.

This species differs from *A. nodulosus* Phill., de Kon.¹ in its septal plan (as explained above), in its few distant tabulæ, and in the small number of its girth-ridges.

CLISIOPHYLLIDS.

Dibunophyllum.

DIBUNOPHYLLUM θ Vaughan, and an early form of DIBUNOPHYLLUM ψ Vaughan. (Pl. XL, fig. 7.)

The figured specimen is from D_1 of Port Eynon; it shows a cross-section of two cylindrical *Dibunophylla*, side by side.

¹ 'Nouv. Rech. Anim. Foss. Terr. Carb. Belg.' 1872, p. 74 & pl. vi, fig. 5.

I. The right-hand specimen presents the typical characters of *Dibunophyllum* θ , namely :—

A circular, ill-bounded central area, which is completely bisected by a somewhat flexuous mesial plate; the general structure of the area is simply reticulate.

A loosely vesicular external area, with a very ill-defined inner boundary. Extremely short minor septa.

II. The left-hand specimen illustrates the early form of *Dibunophyllum* ψ , with which it agrees in the following characters :—

The cuspidate boundary of the central area.

The suggestion of Aspidophylloid characters: that is (1) the abrupt termination of the mesial plate within the central area; (2) the fact that the lamellæ start from the boundary of the area, and fall short of the mesial plate.

The close-set vesicles of the external area and the well-defined inner wall.

The elongate minor septa, which usually extend completely across the external area.

The differences are mainly of degree only, namely :—

The early form is cylindrical (convergent on *Dibunophyllum* θ); *Dibunophyllum* ψ is conical.

The sharp, cuspidate boundary of the central area and the Aspidophylloid characters are more strongly marked in *Dibunophyllum* ψ .

A purely vesicular peripheral area is practically absent from the D_1 form.

This early form of *Dibunophyllum* ψ is not uncommon at several localities in the South-Western Province.

CYATHAXONIDS.

‘CYATHAXONIA’ sp.

Compare ‘*Cyathaxonia*’ aff. *costata*, figured in Rush Paper, pl. xxix, fig. 5.

Two poorly preserved specimens, from the base of D_{2-3} in the North Colts Hill Quarry, were collected and kindly presented to me by Mr. Underhill; their matrix is rottenstone. Both specimens show the calyx, and one of them shows, at its lower end, the highly-vaulted under surface of a tabula.

Radiating from the wall are 26 septa, short, thick, and of equal length, except that the fossula is marked by a single short cardinal septum.

The floor of the calyx rises centrally into a tall spear-shaped columella, which is laterally compressed and crested like a helmet; the lateral surfaces of the columella are roughened by the external edges of the numerous close-set longitudinal lamellæ. By rubbing down the columella, the lamellæ are seen to radiate inwards towards the mesial plate which is the downward continuation of the crest.

After further rubbing down, through a very short distance, the section cuts an underlying tabula in a roughly concentric curve, and from this curve a second set of lamellæ radiate inwards towards the mesial plate.

The structure of the columella appears, therefore, to be closely similar to that of a Clisiophyllid, and, in particular, to that of *Carcinophyllum*:—the tall vaulted tabulæ, each buttressed by lamellæ and crested by the mesial plate, are arranged in a series of tents one below, and within, another.

The close approximation of the descending portion of successive tabular vaults forms a thick tubular wall enclosing the entire central cylinder throughout the length of the coral.

Comparison:—The specimen figured in the Rush Paper, and cited above, was derived from the *Cyathaxonia* Beds of Bradbourne in Derbyshire; similar forms, less well preserved, are common at the same level at Rush.

The structural plan is essentially similar to that of the Gower specimens.

By rubbing down the Bradbourne specimen through a complete tabular interval, it was demonstrated that the mesial plate is not continuous through the tabulæ, but that it starts afresh above each tabula as a crest, and extends to the under surface of the next tabula above; the lamellæ apparently exhibit the same phenomenon. The resemblance to the central structure of Clisiophyllids is, therefore, extremely close.

Specific differences from the Gower form are obvious:—In the Bradbourne specimen there is a narrow peripheral ring of vesicles and the lamellæ are much stronger and more closely set than in the Gower specimens. In both these characters there is a still closer approach to structures characteristic of *Carcinophyllum*.

(B) BRACHIOPODS.

PRODUCTIDS.

Productus.

PRODUCTUS SULCATUS Sow. (Pl. XLI, figs. 1 a & 1 b.)

Sowerby, 'Min. Conch.' vol. iv (1823) pl. cccix, fig. 2.

The holotype of the species is preserved at the British Museum (Natural History) in the Sowerby Collection; it was obtained from Derby. The same form is common in Yorkshire at the top of the Carboniferous Limestone.

The specimen here figured is from D₂₋₃ of the Oystermouth Quarry, where the species is not common.

Description:—The pedicle-valve has the following striking characters:—

Form cylindrical and of quadrate cross-section. Umbonal region parallel to the marginal, and therefore invisible in a 'back view'; the two regions are united by a high vault. Strong reticulation on the umbonal region. Coarse, tall, marginal ribs separated by deep furrows (whence the specific name). Cylindrical wings strongly projecting, and each ornamented by two rows of spines, respectively along and oblique to the hinge-line.

The holotype of *Productus sulcatus* Sow. has a shallow median sinus, whereas the Oystermouth specimen is merely flattened in the median region.

Comparison:—*Pr. costatus* J. de C. Sow., 'Min. Conch.' vol. vi (1829) pl. dlx, fig. 1, differs in possessing the following characters:—

Form broad and flattened. Both umbonal and marginal areas are visible in a 'back view,' and a vault is undeveloped. Coarse marginal ribs, which are remarkably flattened. Strong development of spines on the wing-ridges. Deep, narrow, median sinus.

Evolution:—The strength of the reticulation of the umbonal region, the thickening of the marginal ribs, and the development of spine-ridges are characters which commonly indicate phylogenetic old-age. It seems probable that *Pr. sulcatus* is derived from some such form as the specimen in the Sowerby Collection which is doubtfully identified with *Pr. antiquatus* Sow., 'Min. Conch.' vol. iv, pl. ccxvii, fig. 5; whereas *Pr. costatus* has probably resulted, by parallel development, from *Pr. antiquatus* Sow., *loc. cit.* fig. 1.¹

PRODUCTUS (MARGINIFERA) LONGISPINUS Sow., var. (Pl. XLI, figs. 2a & 2b.)

The predominant form of this species in D₂₋₃ of the Oystermouth Quarry agrees with *Productus setosus* Phill.,² var. *tissingtonensis* Sibly, Q. J. G. S. vol. lxiv (1908) p. 77 & pl. i, figs. 6a-6b.

The most striking character of this variant—namely, the large central and marginal rib in the pedicle-valve—is well shown in Dr. Sibly's figures. Two further points are illustrated in the plate which accompanies the present paper:—

Pl. XLI, fig. 2a—a partial cast of the interior of the brachial valve—shows a marginal groove, representing a ridge surrounding the muscular region; this is the essential character of the genus *Marginifera* of Waagen.

Pl. XLI, fig. 2b—a pedicle-valve—shows the sharply-raised, collar-like termination of the cylindrical wing.

(The structural exaggerations here noticed are probably no more than indications of phylogenetic old-age, for they occur also in other groups, as, for example, in *Pr. concinnus*.)

This variety is remarkably widespread in the upper part of the *Dibunophyllum* Zone throughout the British Isles.

¹ I cannot distinguish fig. 1 from the figure of *Pr. semireticulatus* (Mart.) in 'Petrificata Derbiensis' 1809, pl. xxxii, figs. 1 & 2. If, therefore, *Pr. antiquatus* Sow. is legitimately retained in addition to *Pr. semireticulatus* (Mart.), it must denote such a form as that of Sowerby, pl. ccxvii, fig. 5. On the other hand, if it be considered that Sowerby, by his query, rejected Martin's species as insufficiently described and figured, and proposed the new specific name *antiquatus* in its stead, fig. 1 becomes the holotype of *Pr. antiquatus* Sow., and *Pr. semireticulatus* (Mart.) dies out. In my use of *Pr. antiquatus* Sow. I have adopted the first-mentioned course.

² If *Productus setosus* Phill. ('Geol. Yorks.' pt. ii, 1836, pl. viii, fig. 9) be separated from *Pr. longispinus* Sow. (refigured in 'Davidson,' pl. xxxv, fig. 5), the Oystermouth form must be regarded as a variant of the small globose type of Sowerby's species, rather than of Phillips's large and sulcate type. On the other hand, *Pr. setosus* Phill., var. Phill. (*op. cit.* pl. viii, fig. 17) agrees closely in convexity and ribbing with our variant.

Chonetes.

CHONETES sp. (Pl. XLI, figs. 3 a & 3 b.)

Circumference rectangular, with nearly square cardinal angles. Ribbing fine and close-set.

In the pedicle-valve the form is flattened-convex, with the maximum rise at half-way across the valve. The strong growth-halt and the thickened marginal band below it cause a deep marginal depression in the cast. A shallow marginal median sinus and a corresponding median inflection of the circumference are occasionally developed.

This species is very common at certain levels in D₂₋₃ of Oystermouth and Bishopston in Gower. It also occurs abundantly farther west in the South-Western Province, at Ragwen Point near Pendine, at approximately the same level (namely, Horizon ϵ).

Daviesiella.

The essential character of *Daviesiella* Waagen as exhibited in the genotype *D. llangollensis* (Dav.) is the presence of an additional pair of adductors in the large valve ('Davidson' pl. lv, fig. 9—scars lettered C).

The large Productoid *Chonetes* from C—*Chonetes* cf. *comoides*—has no additional adductors; whereas, in the Productoid *Chonetes* from D—*Chonetes* (*Daviesiella*) aff. *comoides*—these scars are very conspicuous.

STROPHOMENIDS.

'ORTHOTETES' cf. CRENISTRIA (Phill.). (Pl. XLI, fig. 4.)

Description.—Hinge-line less than the width of the shell; area asymmetrical. Brachial valve strongly convex as a whole, but flattened in the neighbourhood of the beak.

The radial ornament is strong and markedly periodic.

The concentric fine, close-set growth-lines produce a delicate 'crenistriation' on the ribs and a reticulation of the narrow flat interspaces; there are a few well-marked growth-halts. Dental plates on a simple Strophomenoid plan.

Comparison.—In area and ornament this form approaches very closely to *O. crenistria*; by its marked convexity¹ and more strongly periodic ribbing it is easily distinguished.

This form is not uncommon in the Oystermouth beds (D₂₋₃).

¹ The figured specimen is from thin shales and is crushed flat; specimens showing the strong convexity of the brachial valve were, however, obtained.

SPIRIFERIDS.

Spirifer.

SPIRIFER BISULCATUS Sow., var. OYSTERMOUTHENSIS, nov. (Pl. XLI, figs. 5a & 5b.)

This variant differs from Davidson's figure¹ of the type of the species ('Davidson,' pl. vi, figs. 6-9) in having:—

- (1) Broad scaped ribs and a gently convex mesial fold, but little differentiated from the flanks;
- (2) A flatter and more transverse form; and
- (3) The cardinal region almost smooth.

It is extremely abundant in D₂₋₃ of Oystermouth, but rare in D₂₋₃ of Bishopston.

Comparisons.—*Spirifer calcaratus* McCoy² (Pl. XLI, fig. 5c) differs in its spine-like wings, unequally developed on the two sides. This species is common in D₂₋₃ of Pateley Bridge (Yorkshire).

Spirifer bisulcatus Sow. var., Q. J. G. S. vol. lxiv (1908) pl. 1, figs. 3a & 3b, differs in its small size and want of bilateral symmetry; it is common in Lower P of Loughshinny (Co. Dublin).

Ornament.—The intersection of radial and concentric ornament produces the characteristic 'twilling' shown in fig. 5b. This type of ornament is common to all the forms mentioned above.

SPIRIFER near INCREBESCENS Hall. (Pl. XLI, fig. 6.)

Compare *Sp. increbescens* in Hall & Clarke, 'Palæont. New York,' vol. viii, pt. ii (1894) pl. xxx, figs. 27-30 (specimen from Chester Limestone).

Compare also the type of *Sp. bisulcatus* Sow., as refigured by Davidson (pl. vi, figs. 6-9).

Our species only differs from *Sp. increbescens* in its smaller size (27 mm. wide as against 38 mm.), and probably also in the smaller degree of differentiation of the median fold from the flanks.

The shagreen-like ornament is also not so conspicuous as in fig. 30 of the American species.

The differences from the type of *Spirifer bisulcatus* Sow. are more numerous, namely:—

- (1) The smaller size.
- (2) The tall, angular median fold is composed of a narrow, central, grooved rib and two flat broad ribs, which form the steep slopes of the fold on either side; the valve-intersection is strongly deflected into a tall domed arch. In *Sp. bisulcatus* the median fold is gently and continuously convex from flank to flank, and the deflection is semicircular.

¹ There is considerable doubt whether the imperfect specimen in the Sowerby Collection at the British Museum (Natural History) is actually the holotype of *Spirifer bisulcatus* Sow.; if so, both Sowerby and Davidson have erroneously restored the mesial fold.

² 'Synopsis Carb. Foss. Ireland' 1844, pl. xxi, fig. 3; refigured in 'Davidson' pl. vii, fig. 4.

- (3) The slopes of the median fold pass gradually into the flanks, whereas, in *Sp. bisulcatus*, the fold is sharply separated from the flanks by a change in curvature and by two strongly marked furrows (to the presence of which¹ the species owes its specific name).

This species (or variant) is not uncommon in D_{2-3} near the top of Oystermouth Quarry. Its small, convex, *pinguis*-like appearance readily distinguishes it from the broad flattened *Sp. bisulcatus*, var. *oystermouthensis*, which is the predominant form.

SPIRIFER WICKENSIS, sp. nov. (Pl. XLI, fig. 7.)

Compare the following forms figured in 'Davidson':—

'*Sp. duplicicosta*,' var., pl. iv, fig. 4 (from Park Hill).

'*Sp. grandicostatus*,' pl. vii, figs. 8-11 (from Bolland).

Also compare:

Sp. cameratus Morton, var. figured in Hall & Clarke, 'Palæont. New York,' vol. viii, pt. ii (1894) pl. xxxii, fig. 12, from 'Coal-Measures.'²

See also *Sp. striatus*, var., in Q. J. G. S. vol. lxii (1906) p. 310 (from the Curkeen Limestone).

Description.—Cardinal angles often dissimilar, as in the Bolland specimen cited above.

Median fold bounded by two deep furrows.

Ribs few, but tall, angular, and separated by deep furrows; the striking and irregular manner in which they fork, so that the branches often diverge very widely, makes the species easy of recognition.

In so far as the type of ribbing is concerned, this is an old-age character that can be arrived at by structural change within distinct gentes. For example:—

- (1) The form in the Curkeen Limestone (Co. Dublin) is clearly a derivative from a striate stock, since it is connected with more normal members of that stock by intermediate forms.
- (2) The American variant is derived from a stock having the ribs in fascicles, and has been evolved by the blending of each fascicle into a single strong rib.
- (3) It is probable that in very many cases this type of ribbing has been arrived at from a variant of *Sp. bisulcatus* with coarse ribs, in which the characteristic forking is already initiated near the margin.

Hence this species is rather of the nature of a circulus than of a true species; it has, however, an important stratigraphical value, since it is widely distributed and is as yet unknown below D_2 .

In the Bristol area *Sp. wickensis* occurs abundantly in D_2 at Wick (whence the specific name) and, somewhat rarely, at Wrington and in the Avon section at the same level.

In Gower it is a rare associate of the D_{2-3} fauna, and has been found in Oystermouth Quarry.

¹ Davidson seems to have regarded the two grooves on the fold itself, to which the fold owes its tripartite character, as the reason for its specific name. Sowerby, however, states clearly that the character recorded in the name is the separation of the fold from the flanks by two deep furrows.

² Hall & Clarke assign to the 'Coal-Measures' British specimens obtained from Upper D (for instance, the types of the species:—*Spirifer striatus*, *Reticularia lineata*, *Martinia glabra*, *Productus longispinus*).

In Northumberland it has been found by Mr. Stanley Smith at the level of the Acre Limestone (probably of D_{2-3} age); it is there associated with a brachiopod fauna similar to that of Wick and with corals of Oystermouth facies.

The form from the Curkeen Limestone of County Dublin also occurs at approximately the same level.

Martinia.

MARTINIA GLABRA (Mart.), varietal form.

Compare *M. glabra*, mut. P., 'Loughshinny Paper' p. 468 & pl. 1, fig. 8.

This variety differs from Martin's type in its more equidimensional form and in the more flattened brachial valve. The cast of a pedicle-valve shows the absence of dental plates, and the presence of strong ridges which indicate deep furrows in the interior of the valve—characters which are distinctive of the genus, and are equally well exhibited in *M. ovalis* (Phill.).

Compared with the later mutation from P of Loughshinny, the muscular scars are very indistinct, whereas they are deeply impressed in the Loughshinny form.

This variety occurs throughout the Oystermouth Series (D_{2-3}); it is rare immediately above the massive limestone of North Colts Hill Quarry, abundant in Oystermouth Quarry, and rare again in the Bishopston cutting.

ATHYRIDS.

Athyris.

ATHYRIS (CLIOTHYRIS) GLOBULARIS (Phill.). (Pl. XLI, fig. 8 a.)

The holotype of Phillips's species is preserved at the British Museum (Natural History) in the Gilbertson Collection; it bears clear indication of the former presence of fringed expansions in the fine imprints which they have marked upon the valve. Hence this species may be regarded as a dwarf and late representative of the gens of *A. glabristria*, with which it agrees precisely in form and in the strongly developed beak.

The Oystermouth specimens only differ from Phillips's type of *A. globularis* in the preservation of portions of the actual fringes. These expansions are very close-set, and the 'fringe' starts immediately at the suture-line with the valve.

Specimens are common at one level in D_{2-3} of Oystermouth Quarry, and the species is widely spread at the top of the limestone-massif in the Midlands and South Yorkshire.

Comparison with *Athyris (Cliothyris) roissyni* (L'Éveillé), Vaughan (figured in this paper, Pl. XLI, fig. 8 b).—The Bristol species is abundant in Km and persists into Z. It differs from *A. globularis* in its non-transverse form, in the upright and small beak, the well-spaced principal expansions, and the existence of a narrow unfringed basal portion at the line of attachment of each expansion.

Comparison with *Athyris roissyi* (L'Éveillé).¹—The figures of L'Éveillé's type depict a shell of more or less transverse form with a large beak and perforation, a well-marked median fold and strong distant growth-halts. If we accept the tacit assumption, made by both L. G. de Koninck and T. Davidson, that the valves were ornamented with fringed expansions (neither mentioned in L'Éveillé's text nor indicated in his figures), L'Éveillé's species can be well matched by the Z_2 form of '*Cliothyris glabristria*' in the South-Western Province. This conclusion, if recognized,² would involve the following changes of nomenclature:—

'*A. glabristria*, mut. Z_2 ' becomes *A. roissyi* (L'Éveillé).

'*A. roissyi*' of Km and Z becomes a new species, on account of its difference of form and beak (see Pl. XLI, fig. 8 b).

Athyris globularis stands, since it is differentiated from the above forms by the closeness of its expansions and by the fact that they are completely fringed.

IX. SUMMARY OF PALÆONTOLOGICAL SECTIONS.

(a) The existence of a *Modiola* phase at the base suggests conformity to the underlying Old Red Sandstone; this conformity has been demonstrated by Mr. Dixon in North-Western Gower.

(b) The zones Z, C, S and D contain faunal assemblages identical with those of the same zones in the Bristol Area.

It is, however, interesting to note that small *Zaphrentes* of Lower Avonian aspect are abundant just above the *Caninia* Oolite, in association with *Ocyathophyllum* ϕ and *Michelinia grandis*.

(c) The faunal evidence suggests that the lower limit of the D_{2-3} phase lies within D_2 of Bristol, but that the upper limit of the phase lies above that subzone and within the 'Millstone Grit' of the Bristol sequence.

(d) The upper part of C_2 is so closely related faunally with the overlying Lower *Seminula* Zone, that it is impossible to refer these two subzones to different divisions of the Carboniferous Limestone.

(e) The most important coral and brachiopod groups discussed in the paper are:—

Zaphrentis oystermouthensis.

Productus sulcatus and *Productus costatus*.

Forms included in the group of *Spirifer bisulcatus*.

Athyris globularis and *A. roissyi*.

I have received much help from the following geologists, and to them I tender my heartiest thanks:—Mr. R. G. Carruthers, Dr. Wheelton Hind, Dr. F. A. Bather, Mr. J. W. Tutchet, and Mr. W. H. Wickes.

¹ Mém. Soc. Géol. France, vol. ii (1835) pl. ii, figs. 18–20.

² L. G. de Koninck was quite aware of the fact that *Athyris roissyi* had been wrongly identified; but he suggested that no change should be made, in order to avoid confusion. See Ann. Mus. Roy. Hist. Nat. Belg. vol. xiv, pt. 6 (1887) pp. 85. 86.

EXPLANATION OF PLATES XXXVIII-XLI.

PLATE XXXVIII.

- Fig. 1. 'Pitted' limestone in D_1 , Tor-gro (North-Western Gower). A steeply-dipping, weathered bedding-plane. The bag near the right lower corner affords a scale. The limestone is a pseudobreccia, and shows the usual rough surface. (See pp. 490 & 499.)
- Fig. 2. Undolomitized pseudobreccia in D_1 , slightly weathered, Tor-gro. The dark angular 'fragments' are well defined. Dolomite is absent, from both 'fragments' and 'ground-mass'; the amount of argillaceous impurity is very slight. (See pp. 507 *et seqq.*)

PLATE XXXIX.

- Partly dolomitized pseudobreccia in D_1 , Port Eynon (South-Western Gower). Microscope-section $\times 8$ diameters. The dense material is unaltered 'ground-mass'; the less cloudy, much more foraminiferous areas, at the top, bottom, and left, are the recrystallized 'fragments'; the rhomboidal crystals, many with a diagonal cross, in the 'ground-mass' are dolomite. (See pp. 507 *et seqq.*)

PLATE XL.

Upper Avonian corals from the *Cyathaxonia* Phase (D_{2-3}) of Oystermouth, Gower (together with certain forms for comparison).

- Figs. 1 *a*-1 *e*. *Zaphrentis oystermouthensis*, sp. nov. (p. 553). Fig. 1 *a*, external and calicular view, natural size. Base of D_{2-3} , North Colts Hill Quarry. Collected by Mr. Underhill. (For septal plan and curvature of tabulæ in this specimen, see text-figs. pp. 554 & 555.) Figs. 1 *b*-1 *e*. All from D_{2-3} , Oystermouth Quarry ('Black Lias'). Fig. 1 *b*, a very young form, $\times 2\frac{1}{2}$. Fig. 1 *c*, intermediate stage of the same specimen, $\times 1\frac{1}{4}$. Fig. 1 *d*, intermediate stage of another specimen showing differences due to the position of the section with regard to new septa. Fig. 1 *e*, adult stage, $\times 1\frac{1}{4}$.
- Fig. 2 (for comparison). *Zaphrentis enniskilleni* Ed. & H. (p. 554). Natural size. The holotype of the species. [*Olim* Coll. Geol. Soc., R. 5460.—Now in the Museum of Practical Geology.] Lough Gill (Co. Sligo).
- Figs.¹ 3 *a*-3 *c*. *Caninia* aff. *cornucopiæ* Mich., mut. D_{2-3} (p. 555). Three sections from the same specimen, D_{2-3} of Oystermouth Quarry. Fig. 3 *a*, young stage, $\times 2\cdot7$. Fig. 3 *b*, intermediate stage, $\times 1\frac{1}{4}$. Fig. 3 *c*, adult stage, $\times 1\frac{1}{4}$.
- Fig.¹ 4 (for comparison). *Caninia cornucopiæ* Mich., Carruthers (p. 556). Section, $\times 1\frac{1}{4}$. Stackpole Quay, Pembroke. Horizon γ .
- Fig. 5. *Caninia* (?) sp. (p. 556). Natural size. Base of D_{2-3} , North Colts Hill Quarry. Collected by Mr. Underhill.
- Figs. 6 *a* & 6 *b*. *Amplexus* sp. (p. 557), D_{2-3} of Oystermouth Quarry. Fig. 6 *a*, external view, $\times 1\frac{1}{4}$. Fig. 6 *b*, section, $\times 1\cdot2$.
- Fig. 7. *Dibunophyllum* θ Vaughan (to right) and *Dibunophyllum* aff. ψ Vaughan (to left), slightly reduced ($\times \cdot94$). D_1 of Port Eynon (p. 558).

PLATE XLI.

Upper Avonian brachiopods from the *Cyathaxonia* Phase (D_{2-3}) of Oystermouth and Bishopston, Gower (together with certain forms for comparison).

- Figs. 1 *a* & 1 *b*. *Productus sulcatus* Sow. (p. 559). Natural size. D_{2-3} of Oystermouth Quarry. Fig. 1 *a*, pedicle-valve, rostral view. Fig. 1 *b*, pedicle-valve, marginal view.

¹ In figs. 3 *b* and 3 *c* the fossula lies to the right of the plane of symmetry for the external surface; figs. 3 *a* and 4, in which the fossula lies in the plane of symmetry, have been set obliquely so as to facilitate comparison.

- Figs. 2 *a* & 2 *b*. *Productus longispinus* Sow., var. (p. 560). D₂₋₃ of Oystermouth Quarry. Fig. 2 *a*, cast of interior of brachial valve, × 1·5; Fig. 2 *b*, pedicle-valve (showing wing), × 1·6.
- Figs. 3 *a* & 3 *b*. *Chonetes* sp. (p. 561), natural size. Fig. 3 *a*, cast of pedicle-valve; D₂₋₃ (top), Bishopston. Fig. 3 *b*, pedicle-valve from D₂₋₃ of Oystermouth Quarry.
- Fig. 4. '*Orthotetes*' cf. *crenistris* (Phill.) (p. 561). Brachial valve and area, natural size. D₂₋₃ of Oystermouth Quarry. [The specimen is crushed, and does not show the strong convexity of the valve.]
- Figs. 5 *a* & 5 *b*. *Spirifer bisulcatus*, var. *oystermouthensis* nov. (p. 562). Fig. 5 *a*, pedicle-valve, natural size. D₂₋₃ of Oystermouth Quarry. Fig. 5 *b*, portion of a brachial valve, enlarged to show ornament. Same locality.
- Fig. 5 *c* (for comparison). *Spirifer calcaratus* M'Coy (p. 562), natural size; a pedicle-valve from D₂₋₃ of Lolley Scar Quarry, near Pateley Bridge (Yorkshire).
- Fig. 6. *Spirifer* nr. *increbescens* Hall (p. 562), natural size; a brachial valve from D₂₋₃ of Oystermouth Quarry.
- Fig. 7. *Spirifer wickensis*, sp. nov. (p. 563), natural size, a pedicle-valve from D₂₋₃ of Oystermouth Quarry.
- Fig. 8 *a*. *Athyris* (*Cliothyris*) *globularis* (Phill.) (p. 564), a pedicle-valve, × 1·5. D₂₋₃ of Oystermouth Quarry.
- Fig. 8 *b* (for comparison). *Athyris* (*Cliothyris*) *roissyi* of the 'Bristol Paper.' Beak and brachial valve, × 1·6 (p. 564). Km ('*Modiola* phase') of the Avon Section.

DISCUSSION.

Dr. A. STRAHAN observed that this region was one which called especially for combined stratigraphical and palæontological work. The Carboniferous Limestone Series presented a succession of stages in development. The fullest development was exhibited in the southernmost occurrences, as regarded both the sequence and the thickness of zones. A second stage, showing an incomplete sequence and considerable attenuation, with indications of near-shore origin, was presented along parts of the margin of the coalfield. Lastly, in the northernmost occurrences, near Abergavenny and at Pencerig-celch, the series dwindled away to 100 feet or less, while in Pembrokeshire it was wholly overstepped. The inference followed that the coast-line ran through Pembrokeshire, and not far north of Abergavenny. Traverses made at right angles to the coast-line, and comparisons of these various stages of development, could not fail to yield results of the greatest interest. The only other district where such a traverse was possible, lay on the northern side of this same Carboniferous land-area, where the Limestone Series could be followed continuously through Denbighshire to its disappearance in Shropshire. That the radiolarian cherts were of shallow-water origin seemed obvious, nor, so far as he was aware, was there evidence of deep water in any part of the Limestone Series: sedimentation or formation of organic deposits seemed to have kept pace with subsidence so nearly that the water was generally shallow. But he was unable to follow the Authors in their identification of lagoon-phases. He enquired whether further evidence had been obtained for identifying and defining the *Posidonomya* Zone. The change of fauna at the base of the radiolarian cherts

FIG. 1.—‘PITTED’ LIMESTONE IN D₁, TOR-GRO (NORTH-WESTERN GOWER).

T. C. Hall, Photo. (Geol. Surv. Colln.).

FIG. 2.—UNDOLOMITIZED PSEUDOBRECCIA IN D₁, SLIGHTLY WEATHERED, TOR-GRO.

T. C. Hall, Photo. (Geol. Surv. Colln.).

Bemrose Ltd., Collo., Derby.

QUART. JOURN. GEOL. SOC. VOL. LXVII, PL. XXXIX.

E.E.L.D., Photo.

Bemrose Ltd., Collo., Derby.

PARTLY DOLOMITIZED PSEUDOBRECCIA IN D1, PORT-EYNON
(SOUTH-WESTERN GOWER), x 8 DIAMETERS.

J. W. Tutcher, Photo.

Bemrose Ltd., Collo., Derby.

AVONIAN CORALS FROM D₂ AND D₂₋₃ GOWER.
 (* COMPARATIVE FORMS.)

J. W. Tutcher, Photo.

Bemrose Ltd., Collo., Derby.

was most pronounced, but the upward limit of a 'Posidonomya Zone' was extremely vague.

The PRESIDENT (Prof. WATTS) congratulated the Authors on the number of issues that they had raised, and on the discussion to which the paper had given rise. He agreed with Dr. Strahan that the South Wales limestone outcrops afford an unrivalled opportunity for the study of the broad variations occurring in a formation throughout its geographical extent: the extent and continuity of the South Wales sequence being far more favourable than the meagre outcrops north of the Midland barrier. He referred to the difficulty in finding modern parallels for the lagoon conditions postulated in the paper, which required wide extent of shallow water, abundance of organisms, and yet freedom from clastic material. Coral-reefs presented almost the only example of such conditions to be observed at the present day. He deprecated too much controversy on the application of Belgian subdivisions to British strata, until the exact extent and character of these subdivisions had been again checked in their country of origin.

Mr. R. H. TIDDEMAN said that he was much pleased with a paper which dealt with so many points of interest. He agreed with the Authors that many of the limestones of Gower had been formed in shallow waters; indeed, when mapping the ground, he thought it probable that the oolites had been formed under æolian conditions. Referring to the term 'lagoon' so much employed, he wished to ask Mr. Dixon whether he had seen anything in the area at all resembling a barrier-reef or reef-knolls, such as the speaker had found in the North of England: he himself had failed to see anything similar in Gower. He was satisfied with the identification of the 'Black Lias' of Oystermouth with the Bishopston white rottenstones, and mentioned that the former, when exposed for a short time to the weather, became covered with a white powder. He believed that the shales above the cherts represented the 'Bowland Shales' (of Phillips), a name which had priority over the Pendleside Shale of a later author.

Mr. A. L. LEACH remarked that the rocks formed under 'shallow-water' conditions appeared to be of great importance in the area described by the Authors, but the application of the definite term 'lagoon-phase' to the conditions under which these sediments were deposited seemed to need further justification. The Authors apparently did not claim a coral-reef origin for any of the rocks described, and the oceanic atoll-lagoons were therefore rightly excluded, as also those coastal lagoons which were bounded seawards by coral-reefs, since neither in Gower nor, so far as his observations went, in Pembrokeshire, were there any limestones of coral-reef origin. Lagoons bounded seawards by islets or by submarine banks were, he thought, usually associated with river-mouths, and were necessarily adjacent to land. Deposits formed in such lagoons would show, he thought, some unconformity with the boundary-deposits, and moreover terrigenous material would probably be present. He had not observed any deposits in the adjacent Pembrokeshire area that could be assigned to lagoons of this type.

The sudden change from the 'calcite-mudstones' of S_2 to the limestones with a deep-water fauna of D_1 , as seen at Lydstep in Pembrokeshire, was difficult to reconcile with the suggested lagoon-formation of the 'calcite-mudstones.'

Dr. T. F. SIBLY remarked that numerous interesting questions had been raised by this paper. Obviously, a careful study of the evidence would be necessary, in order to appreciate the Authors' conclusions with regard to 'complete cycles of intermittent change,' arrived at by comparing the sequences of lithological types in the three districts. The speaker, however, pointed out that, in view of the small extent of the area under consideration, and the distinctly shallow-water facies of the Avonian as a whole in this area, correlation of movements of the sea-floor in different parts of this area with similar disturbances in different parts of the much larger Bristol area, must be open to considerable error. He asked Mr. Dixon for a definition of the distinctive lithological facies of a 'lagoon-phase.'

It would have been instructive to hear a summary of the special features of the faunal succession in this area. The Avonian fauna appeared to have maintained, on the whole, its normal progression, despite remarkable vicissitudes. With regard to the very wide question of the limiting horizon between Lower and Upper Avonian, was this area, which clearly suffered a maximum of disturbance, a favourable one in which to deal with the question on a faunal basis? The description of the D_2 - D_3 phase of the *Dibunophyllum* Zone would doubtless furnish many important facts, and would be especially valuable to those interested in the difficult subject of the correlation of the highest Avonian beds in different areas.

Mr. J. H. COLLINS regretted that the Authors had not supplied a cross-section, and that no statements had been made as to the thicknesses of the various zones and their subdivisions.

Mr. COSMO JOHNS welcomed this account of the lithology and physiographic conditions of deposition of the Carboniferous Limestone in a most interesting district. The Authors' careful observations confirmed the evidence obtained in other areas, and demonstrated the shallowness of the Lower Carboniferous Sea. The non-appearance of dolomites at horizons where, from the evidence of shallowing, they would be expected, might be explained if a critical temperature rather than depth be the factor determining their occurrence. It would appear that the Authors, by 'lagoon-phase,' simply meant partly land-locked shallow waters, free from disturbance by currents. In such restricted areas the water might easily be appreciably warmer than in the outer sea, and thus dolomites for this reason might have a local development only. This suggestion could be put to experimental proof without much difficulty, and temperatures near 30° C. would, from such evidence as is available, seem most promising. He regretted the absence of Dr. Vaughan, in view of the important faunal considerations that arose out of the work done in the area.

Mr. G. W. YOUNG enquired whether the Authors had formulated any theory as to the cause of the frequent oscillation of the sea-

bottom which they believed had taken place; and also whether their suggestion that a considerable deepening occurred at the close of the S_2 lagoon-stage was compatible with the presence, in the succeeding deposit, of a 'thin coaly layer.' As this, in one portion of the district, was accompanied by an underclay, it seemed to present a difficulty.

Mr. A. WADE compared these beds with some which had been forming quite recently at the northern end of the Red Sea. Here, there was a very shallow platform fringing the Eastern Coast, and running out to sea for upwards of 20 miles. A series of limestones could be traced on the islands from the edge of the platform to the mainland. The changes which took place in the physical characters of these limestones might clear up the difficulty that the Authors had with the limestones of the *Seminula* Zone, which were dolomitic in the south-west of Gower but were not so in the north-west. They were, however, entirely oolitic in this direction. The limestones referred to in the Red Sea area changed from massive limestones (not oolitic, but to some extent dolomitic) on Jubal Island at the edge of the reef, to oolitic limestones interbanded with beds of very fine sediment on Gaysum Island, half-way between Jubal and the coast; while on the mainland, they consisted of oolitic limestones with many gasteropod-remains, interbedded with sands and coarse grits—very like the material of the Millstone Grit. The later oolites were evidently laid down in the estuary of some river.

He was interested to hear that Mr. Tiddeman had long considered that some of the oolites of Gower might have been laid down under æolian conditions, because some of the oolites in these recent Red-Sea beds were probably affected by similar conditions. It seemed, therefore, that the conditions which the Authors considered to have existed during the deposition of these beds in Gower, might find a close parallel in the conditions which recently existed, and probably still exist, in the northern part of the Red Sea.

Mr. DIXON, after thanking the Fellows for their kind reception of the paper, remarked, with reference to an observation by the President, on the usefulness of a classification of the Lower Carboniferous in facilitating comparison of developments in different districts; and added, in reply to Mr. Tiddeman, that the occasional occurrence of large marine fossils in even the most unfossiliferous oolite militated against the view that any of these rocks were of æolian origin.

Regarding the name to be applied to the shales above the Limestone Series in Gower, the Pendleside Series had good claims, in that it represented rocks to which a distinctive fauna had been ascribed; and further, in answer to an objection raised by Dr. Strahan, he remarked that the correlation depended not solely on the occurrence of *Posidonomya becheri* Bronn, but rather on that of *Glyphioceras spirale* Phill. In agreement with Mr. Tiddeman, he did not think that any of the Gower limestones possessed features characteristic of true coral-reefs.

In reply to Mr. Leach and others, he stated that the calcareous 'lagoon-phases'—the *Modiola* phases—were all distinguished in places by exceedingly fine-grained calcite-mudstones with an impoverished fauna of lamellibranchs (especially *Modiola*), ostracods, and *Spirorbis*.¹ A certain amount of terrigenous material, mud and fine sand, was found in them all.

He took exception to Dr. Sibly's remark that the Gower area had been exceptionally shallow; on the contrary, he considered it in many respects typical of the South-Western Province.

He was very glad to hear from Mr. Wade that the conditions which must have prevailed during lagoon-phases could be approximately paralleled at the present day along the coast of the Red Sea, as he had not had an opportunity of enquiring whether modern representatives of them were known. The suggestion of Mr. Cosmo Johns that a lowered sea-temperature might account for the absence of dolomite from the Upper Avonian of the North-Western district was welcomed, as the speaker had suspected that the influence of the waters that deposited the Pendleside Shales was beginning to make itself felt there at that time, and a low temperature would also explain the extermination of the previously-rich fauna from the area on the arrival of those waters in force.

In conclusion, he remarked that, although there appeared to be no reason why lagoon-conditions should not be established during any shallow-water period, it was a matter of observation that they immediately preceded sharp subsidence.

¹ Lack of time prevented the speaker from illustrating the differences between the lagoon-phase deposits and the standard limestones as being similar to those between the Rhatic—a typical lagoon-phase—and the open-sea Lias.