DUCTLESS GLANDS AND THE NERVOUS SYSTEM.

The Pituitary Body and its Disorders. By HARVEY CUSHING, M.D.
Philadelphia: J. P. Lippincott Co., 1912.

The Internal Secretory Organs. By Professor ARTUR BIEDL, of Vienna.
English Translation by LINDA FORSTER, and Preface by LEONARD
WILLIAMS, M.D. London: John Bale, Sons and Danielsson, 1912.

*Innere Sekretion, ihre physiologischen Grundlagen und ihre Bedeutung
für die Pathologie.* Von Professor ARTUR BIEDL. Zweite
neubearbeitete Auflage mit 131 Textfiguren und 20 mehrfarbigen
Abbildungen auf 8 Tafeln. Berlin und Wien: Urban und Schwar-
zenberg, 1913.

DUCTLESS glands are in the present day theory of medicine as vaccines in its practice. There is little honest knowledge of either, yet all features of health and disease may be lightly ascribed to the one, and every ailment finds its golden hope of cure by injections of the other.

The books discussed in these pages will do much to dislodge confusion and ignorance, for they give a fair statement of what is known of the ductless glands, and they are written by men who rank among the chief authorities living on the subjects with which they deal. Cushing has modelled his treatise in the form adopted by Pierre Marie, Addison, and the great writers of the past, a separate account of each of a series of clinical cases being supplemented by the author's personal interpretation of the phenomena, so far as that can be based upon physiological experiment and pathological analysis. It is a clear record of fact, and while the evidence is marshalled to lead to definite conclusions, yet it is so displayed that criticism can at once disentangle the proved from the supposed and stop short, if it cares, of agreement with the final judgment. Original work has the advantage of this directness and simplicity, for it gives the outlook of a single mind.

Biedl attempts a more spacious task. Writing from the physiological point of view and not as Cushing, alert for every point that may tell in the diagnosis and treatment of clinical disease, he assembles all the observations that have ever been made on the ductless glands, in comparative anatomy, in chemistry, pathology, and especially in the

laboratories of experimental physiology. The host of papers so called up is nearly 4,000 strong, and its catalogue alone occupies more than 100 pages, a quarter of the entire book. But each in the multitude speaks as loudly as his fellow, and it is hard to pick out the main theme amid the babel of voices. In his desire to be disinterested and fair, Biedl has gone almost too far and produced rather an index catalogue than what one would have desired, a reasoned exposition of the views of the Viennese school, whose workers, with Biedl at their head, have explored every part of this wide field of labour. Worst of all is the extreme deference in the body of the book to that tedious custom which in physiology couples every observation with some man's name, as though it were a personal opinion that is quoted rather than a scientific demonstration of a fact. The result is as baffling as the walls of a railway station are to the traveller who, seeking to know where he is, hunts in vain for the name of the town amid the patchwork of advertisements. Time must establish a saner practice in this respect and teach the wisdom of what was written by Sir Thomas Brown in preface to *Pseudodoxia Epidemica*, "For Knowledge is made by Oblivion, and to purchase a clear and warrantable body of Truth, we must forget and part with much we know."

The English translation is somewhat unlucky in the time of its publication. Appearing at the close of 1912, though made from the first edition of the German text in 1910, it barely antedates Biedl's second edition. The latter is a great improvement on the first, printed in much pleasanter form, enlarged to two volumes and nearly twice its original size, and freely illustrated with reproductions from the chief papers in pathology, comparative anatomy, and physiology, so that the reader loses much by being confined to the English of the first translation. But the translator, Linda Forster, has done the work well. Slight mistakes catch the eye here and there. "Darmanhangsdrüse" (p. 13) are not the glands of the appendix, and despite the statement of p. 16 iodothylin does not occur in the adrenals. "Autonomic" was the word employed by Langley to designate Gaskell's visceral nerves, and the translator brings this back again from the German into English with the needless change to "autonomous." Acquaintance with physiology would have saved this slip, which, like several others, shows ignorance of English terminology rather than difficulty with the German text.

A brief abstract of the main teaching of these two books may be of interest to readers of *BRAIN*, for the pathology of the ductless glands touches neurology closely at many points.

THE PITUITARY GLAND.

Roughly, each decade has marked a great stride forward in the knowledge of this gland; 1889 brought the association of acromegaly with disease of the gland by Pierre Marie and Marinesco, the French clinician believing the main fault to be the loss of the secreting gland substance. In 1900 came the clinical recognition by Babinski and Fröhlich of the fatty eunuchoid condition, "dystrophia adiposogenitalis," that may be related to an undoubted atrophy or destruction of the gland. In 1910 Cushing gave the Harvey Lecture which has been expanded into this present book.

Physiological experiments have in the meantime established the fact that from the posterior "nervous" part of the pituitary gland can be extracted an active principle which stimulates all plain muscle to contraction [10], the arteries of the lung as well as those of the systemic vessels (Dale [3]), and which excites some gland cells to prompt secretion, notably those of the kidney and mammary gland; but these observations at present have little value in the discussion of the clinical problem and their meaning is altogether obscure.

For the adrenal gland a law has been clearly proved that stimulation, whether it be of gland-cell or of muscle, by the active principle, adrenalin, can occur only in tissues innervated by sympathetic nerves, and that the reaction caused in them is almost the same as that to the nervous impulse. No such rule relating the action of pituitary extract applies to any part of the nervous system. At present, therefore, no guess can be hazarded as to why this substance is produced in the gland and excreted into the body; indeed, its stimulating property may be quite accidental to the gland extract, and in no way concerned with the true function of the cells.

The pituitary is an extremely complex mass of tissue. In front is the real glandular mass, the pars anterior, most richly vascularized, and containing at least two different types of cell, the one staining blue with hæmatoxylin and eosin and the other a distinctive red. It is not known whether the red eosinophile cell is morphologically and physiologically distinct from the blue (chromophobe), or whether it simply derives its colour from a special phase of secretory activity. Behind lies the pars nervosa, tissue of an indeterminate character, like neuroglia but with very few nucleated cells. The anterior crescentic mass envelopes the pars nervosa, and between the two is a cleft lined on either side by a deep glandular layer, the pars intermedia, which differs from the pars anterior in that its cells are not packed compactly one against the other

but resemble in their arrangement those of the thyroid gland, alveolar spaces being formed by distension with a colloid material.

The active principle, obtained by grinding with sand and boiling with saline solution, is yielded by the *pars nervosa* only. Eosinophile colloid material seems to be secreted in abundance from the true glandular cells, those of the *pars anterior* as well as those of the *pars intermedia*. Herring believes that the secretion passes, probably undergoing some change in the meantime, through the *pars nervosa* and thence by lymphatic clefts into the third cerebral ventricle and the system of the cerebrospinal fluid circulation [7]. This has not been fully proved to take place, whereas eosinophile colloid material can always be readily demonstrated in the rich capillary network of the glandular anterior part. Perhaps the colloid material is passed directly into the bloodstream; if such were the case the colloid would have nothing to do with the *pars nervosa* nor with the active principle obtained from the latter, of which alone the properties have been studied, and there would be no knowledge at all of the functions of the *pars glandularis*.

Cushing has recently endeavoured to show that the secretion from the pituitary is controlled by the cervical sympathetic nerve, but the experiments that he brings forward are of a very doubtful nature [2].

The clinical phenomena produced by disease of the gland are two-fold, those caused by direct pressure of its growth on neighbouring structures, such as the optic nerve, and those which are exhibited by the body generally as the result of too abundant or too scanty pituitary secretion. Symptoms of the second group, so far as is known, appear only in the slow changes of the form and growth of the body, whereby the normal type of manhood is perverted to the acromegalic giant or to a lethargic, eunuchoid creature overgrown with fat. This in itself is amazing enough, that a chemical secretion can slowly act with some mysterious power upon the texture of our bodies and alter its entire pattern.

But the pituitary has an abundant blood-supply, and its glandular cells pour out daily a considerable quantity of eosinophile material. Still greater must be the excretion of the thyroid from its larger bulk; while it has been proved by direct experiment that adrenalin is discharged into the blood from the medullary cells of the adrenal with almost every emotional and vasomotor reflex. Hence there may be other and speedier actions than those concerned in bodily growth. Other metabolic affairs are controlled by the secretions of the ductless glands; and from the pathological observations of the slow changes

wrought by glandular disease arises a fresh conception of physiological activity, that growth, metabolism, muscular tone, indeed much of the steady regulation of the life that burns within and gives us being, results from the daily interplay of these glandular secretions which each and all in their part restrain or accelerate the activities of the various tissues, and so maintain the normal type. This aspect of the question will be referred to again later in considering the meaning of pluriglandular insufficiency.

Cushing regards acromegaly as the index of excessive secretion by the pituitary; diminished secretion has its special clinical features of obesity with eunuchoid changes, and may be caused by adenomata or tumours destroying the pituitary directly. The same result may be caused indirectly as the result of pressure by cerebral growths. Acromegaly is then comparable with exophthalmic goitre, and the second condition with myxœdema.

In these glandular diseases affixes are needed with the noun to denote the state of plus or minus secretory activity. General use accepts hyper- and hypo-, although they are found to be difficult to distinguish from one another in speech, and in the case of the pituitary lead to a slight confusion with the hypophysis itself. There is need of a pair of words that shall be in sharper contrast one with another, and it therefore seems better to take the prepositions *super* and *sub*, which give the desired contrast.

Acromegaly is assumed by Cushing to indicate superpituitarism. The classical case of subpituitarism is that reported by Madelung, a girl aged 9 who was shot in the eye by a rifle bullet which lodged in the sella turcica. She failed to grow, and became exceedingly obese. Very similar is the result in young animals, according to Cushing (though this is denied by Horsley [8]), when the pituitary is experimentally damaged. He gives the following list of changes caused by subpituitarism in children or young adults: infantilism of the sexual organs, absence of pubic and axillary hairs, adiposity, especially of the subcutaneous fat, a dry skin, polyuria, depressed temperature, but a febrile rise of two or three degrees upon intramuscular injection of an extract of the anterior lobe of the pituitary gland, increased tolerance of carbohydrates, so that 200 or 300 gm. of glucose by the mouth do not cause glycosuria. In the adult these phenomena may occur independently (Fröhlich's type), or develop as the last stage of exhaustion in a gland which had in its earlier spendthrift excess produced acromegaly. Good results are said to have been obtained by feeding with pituitary gland

extract in these cases, though much larger doses are required than have generally been tried in England.¹

Especially interesting is Cushing's account of adenomata of the pituitary. These benign growths are common in the adrenal, where they cause no symptoms; in the thyroid, where they may cause pressure on the trachea and become the seat of cystic or hæmorrhagic degeneration, and they are not rare in the pituitary. Their cells never excrete with such activity as to cause respectively the features of exophthalmic goitre or acromegaly. But in the pituitary they induce early pressure symptoms, bitemporal headache by distension of the dura mater enveloping the sella turcica, blindness or temporal hemianopia by pressure on the optic chiasma, and often subpituitarism by pressure on the rest of the healthy gland. Surgical treatment must be early, to preserve sight and save the gland. Cushing performs a sellar decompression, driving a tunnel from a submucous incision beneath the upper lip, up through the septum nasi and the sphenoidal sinuses, to expose the gland by reflecting its dura mater. This allows the adenoma to grow downwards, and as a rule suffices. Sight returns in a few days with the escape of the optic nerve from pressure.

Cushing describes in all forty-seven clinical cases of pituitary disease, of which twenty were proved by the microscope to be due to adenomatous growths causing subpituitarism. Of these forty-seven, 10 per cent. were Jews. The question is not raised, but it would be interesting to learn whether Jews as a race are particularly liable to pituitary derangements.

THE ADRENAL GLANDS.

It was in respect of these organs that the riddle of the ductless glands was first proposed by Addison's clinical observations in the disease that has since borne his name; and physiological work of the last twenty years, aided by the arguments of developmental anatomy, has won far clearer knowledge of the adrenals than it has of the other ductless glands. In this instance physiology has outstripped clinical pathology, while the latter has been unable as yet to make use of the

¹ Cushing employs a dry powdered extract obtained after passage of the entire fresh bovine gland through a Buchner press. Of this 30 grains may be given daily, that is the equivalent each day of about 150 entire fresh pituitary glands. This corresponds roughly to 1,500 daily of Burroughs and Wellcome's tabloids, of which 10 contain the extractives of one entire gland. Cases of subpituitarism are very tolerant of sugar, from which Cushing deduces a test for the adequacy of the treatment, namely, that pituitary extract shall be pushed until sugar appears in the urine on a meal of 150 grm. of dextrose. This in one case required a daily consumption of 500 entire glands.

discoveries of the physiologist. But it is well to remember that even in this later work, as in all the advances of knowledge of the ductless glands, the first promptings came from clinical observations. George Oliver thought that he had observed constriction of the vessels at the wrist in man when suprarenal gland extract was taken by the mouth. Perhaps the observation was erroneous, but it inspired him to seek Schäfer's aid in testing its truth by direct records of the blood pressure in animals, and the result of their joint work opened up new country, which has since been intimately explored by the many investigators who followed in their footsteps.

Balfour recognized the chief landmark when he saw that the cells of the adrenal medulla are derived from the same source as those of the sympathetic ganglia. This, indeed, is the peculiar interest of the gland, that it is practically a part of the nervous system.

The nature of this relationship has been worked out chiefly by the Cambridge School of Physiology, where the brilliant paper by Gaskell [5] in 1886 lit an eager interest in the visceral nerves that year by year has given forth more light. Gaskell challenged the old teaching, founded on Bichat's academic generalizations, which regarded the vegetative or sympathetic nervous system as a self-sufficing machinery for regulating the affairs of the viscera and almost independent of the central nervous system. Nowadays it is difficult to realise how deeply rooted was this opinion thirty years ago, and how widely it influenced all thought of the time as an almost axiomatic truth. Gaskell restored the monarchy of the central nervous system, showing that all the knotted complexity of the visceral nerves could be unravelled into single constituent strands arising from the spinal cord and carrying the dominant impulses of its central control. His, too, was the first broad classification of the nerves efferent from the brain and spinal cord. He showed that, while the motor nerves to the striped muscles connected with the skeleton issue in all the anterior roots, those to the unstriped muscle of the blood-vessels and viscera and to gland-cells do not flow out in a continuous series of anterior roots down the length of the spinal cord, but appear in three divisions, which are separated one from the other by the nerve-roots of the fore and hind limb plexuses. These visceral nerves all bear ganglion-cell relays on their course from spinal cord to peripheral tissue; by that they are distinguished from the skeletal nerves, and to that is due the anatomical complexity of their distribution. Langley accomplished the difficult task of analysing all the finer details of these nervous tracts, and he has introduced a fresh

DIAGRAM OF THE NERVES EFFERENT FROM THE CENTRAL NERVOUS SYSTEM IN THE MAMMAL.

A, the non-ganglionated ordinary motor nerves to striped muscle, which are distributed segmentally. On the left side these are omitted for simplicity, and only the autonomic or ganglionated visceral nerves to plain muscle are indicated.

Of these, B is the cranio-cervical outflow in the vagus, &c.; C, the thoracico-lumbar or sympathetic proper; D, the sacral outflow, or pelvic visceral nerve to the bladder and colon. All these subdivisions contain both motor and inhibitory nerves.

C_1 is the sympathetic ganglion-cell; C_2 the paraganglion-cell secreting adrenalin, the chief mass of these being concentrated to form the medulla of the adrenal gland, though a few, even in adult life, may be found elsewhere in relation to the various sympathetic ganglia. The black rectangle innervated by the nerves from the cells C_1 represents the mass of plain muscle which is also stimulated by adrenalin, that is by the secretion of C_2 .

Afferent sensory nerves and their posterior root ganglia are all omitted from the diagram. Their course from the viscera is not clearly known.

nomenclature, according to which the entire group of ganglionated visceral nerves is styled the autonomic system; in this are the three subdivisions of the cranio-cervical, the thoracico-lumbar, and the sacral root outflows (see diagram). To the thoracico-lumbar outflow is restricted the old term of the sympathetic.

Now the entire mass of plain muscle united with thoracico-lumbar or sympathetic nerves shows a remarkable reaction to the active principle extracted from the medullary cells of the adrenal. It, and it alone reacts to adrenalin; and the reaction is identical with that to electrical stimulation of the sympathetic nerves. This holds true both for contraction and for relaxation, and for muscles so far asunder in physiological differentiation as those of the rhythmically pulsatile heart or of the tenaciously closed sphincter urethræ. Further, the medullary cells have been shown to discharge adrenalin into the blood-stream, and to do so as the direct result of excitation of the splanchnic nerves which supply the glands. This they are continually doing in the various vascular reflexes.

It is therefore clear that there are two closely related types of cells connected with the primary preganglionic sympathetic fibres coming from the spinal cord:—

(1) The sympathetic ganglion cell, which is distally united with the plain muscle by its axon process, and so provides a path for the nervous impulse.

(2) The medullary or paraganglion-cell, which is not in connection with the muscle but is equally innervated from the spinal cord, and secretes a chemical substance into the blood that can produce an identical excitation of the muscle through its myoneural junction.

These cells (1) and (2), are as brother and sister to one another, being in the closest kinship of morphological parentage. Life can be maintained though all the ganglion-cells (1) are removed; death results from cardio-vascular failure when the adrenal glands, that is the cells of the cortex and cells (2) have been excised. In the moribund animal, after removal of the adrenal glands, stimulation of the vaso-constrictor nerves is almost without effect, because the muscles of the blood-vessels have lost their tone. It may be that the cells (1) and (2) were once identical, and that the liberation of adrenalin was an essential part of the nervous impulse; but now the two processes are carried out separately by separate anatomical units. The paraganglion-cell (2) excretes the adrenalin that maintains the muscle at a fair level of tone for its routine activities; the nervous impulse (1) effects the rapid

alterations from this level that may be required suddenly in special actions of the animal. Roughly, one may compare the muscle to the string of a violin that must be tightened up to its appropriate pitch before the fingers can evoke the rapidly changing notes.

Beyond such rough analysis knowledge has not yet advanced. A secret lies within this relationship of gland-cell to the efficiency of a nervous impulse, that may prove of the utmost value in explaining the processes of nervous activity. Given the solution of it, the formula of interaction between hormone and nervous impulse may perhaps be found capable of application to other departments of the central nervous system than the sympathetic, and a simple material explanation be obtained for such conditions as spinal neurasthenia. To describe a patient with Addison's disease as suffering from cardio-vascular neurasthenia, or from myasthenia gravis of the circulatory muscles is no wide stretch of the fancy.

The adrenal is, therefore, the ductless gland that promises most in the future to neurologists who seek for deeper acquaintance with the physiological processes of nervous activity. And it has been equally generous to historians of the past. Those who inquire into that most fascinating problem, the origin of the vertebrates, have read in the story of the adrenals clear proof that the thoraco-lumbar or sympathetic system is, in virtue of this reaction, a group distinct from the other visceral nerves, a stratum, so to speak, laid down at a probably later epoch and composed of its own peculiar materials.

As will be seen in the discussion of the thyroid gland, Continental physiologists have accepted this teaching of the English school with regard to the visceral nerves—it is described with admirable clearness in the second edition of Biedl—but at Vienna it has been extended to yield illegitimate conclusions with regard to the functions of the other ductless glands.

The adrenal gland is a compound of cortex with medulla. Both fail in Addison's disease, so that the cardio-vascular sympathetic fault carries with it some expression of the lack of cortical secretion. What that may be is still unknown. The cortical cells do not belong immediately to the sympathetic group; they secrete a doubly refractive fat of complex composition like that of the myelin sheath of nerves, but nothing can be said with regard to its function. However, the presence of this lipid in abundance indicates a healthy gland. Cushing falls into the error of regarding it as a phenomenon of degeneration, and so an index that the adrenal as well as the pituitary was affected in some of the cases described by him.

Doubtless there are clinical conditions of superadrenalism due to excessive secretion, but they have not been identified, though frequent endeavours have been made to prove that persistently high blood-pressure should be ascribed to this cause.

On the other hand, overgrowth of the cortex in children, producing large adrenal tumours [1], is beyond doubt associated with sexual precocity, premature adolescence, and an early setting of the body, the very opposite of juvenile acromegaly. As a mysterious oddity may be quoted the fact that the human adrenal is distinguished from that of all other animals by a great overgrowth of the cortex in foetal life; this overgrowth does not occur in cases of hemicephaly, congenital fault that has led to absence of the cerebral hemispheres and reduced the child to the level of the brainless beasts [4].

Simple adenomata of the cortex are very frequent in the adrenal, as they are in the other ductless glands. They cause no pressure symptoms and no clinical phenomena.

THE THYROID GLAND.

In this gland clinical workers first recognised the three main forms of pathological disorder, that have since served as types for the study of changes in the other ductless glands. They are superthyroidism, or Graves's disease; subthyroidism, or myxœdema and cretinism; and adenomata causing pressure symptoms. The adenomata and the parenchymatous goitres are localized or general enlargements of gland tissue, which do not produce secretion proportional to the increase of bulk, and may perhaps be regarded as anatomical evidence that the gland has been striving under adverse conditions to do its proper physiological work.

But there are two chief problems still unsolved, despite much study in the laboratories of experimental research. In the first place arises the question, what is the nature and meaning of the thyroid secretion? Its absence hinders growth, its presence hastens metabolism; with either change curious and special nervous phenomena are seen. Secondly, analysis is perplexed by the double nature of the gland. All the ductless glands, pituitary, adrenal, thyroid, pancreas, and testicle, present this difficulty; two types of cells living in close neighbourhood, often under the same roof, and yet with no obvious reason why they should have chosen such a union.

The associate of the thyroid is the smaller group of parathyroid glands. To them, but without sufficient proof, has been ascribed the

train of nervous symptoms that appear when the thyroid is removed. The leading feature of this condition of tetany is an exaggerated excitability of the skeletal muscles through their motor nerves. Many workers have proved that calcium salts in general lower neuromuscular irritability, and the deduction led to the observation that after removal of the thyroid and parathyroids there is an excessive loss of calcium from the body, a so-called calcium diabetes [9]. This, however, does not seem to be the full explanation of the trouble, for while calcium salts allay tetany after removal of the thyroid, they do not serve to prolong life. Further, it has been urged, but without the least attempt at proof, that all forms of tetany are due to subparathyroidism; and that even myotonia congenita and paralysis agitans are caused by a slow, progressive diminution of parathyroid secretion, while myasthenia gravis is the result of excessive parathyroid activity. This is pretty imagining, but at present bears no relation to substantial fact.

Even with regard to the first question, what is the meaning of the thyroid secretion, little definite can be said. The view that it is required to neutralise certain poisons formed in the metabolism of the body has now few adherents. It is admitted that the thyroid secretes into the blood some substance which influences the heart and the nervous system and growth; that is, it produces one of Starling and Bayliss's hormones, belonging to the same group of simple though potent non-protein bodies as adrenalin, pituitrin, and secretin. Now the Viennese school has elaborated a doctrine with regard to the hormones that is essentially inspired by Ewald Hering's theory of nervous control. The first instances observed of the activity of these bodies were naturally those of stimulation, where a muscle was made to contract or a gland cell to secrete, as in the original discovery of the action of secretin on the pancreas. But adrenalin inhibits some muscles while it causes contraction of others. Hence is deduced the view that a hormone may be purely inhibitory, and, according to Hering's view of inhibition, is therefore a chemical agent which excites an assimilatory process in other organs—in other words, it causes growth. Out of this philosophical analysis of assimilatory and dissimilatory, or inhibiting and exciting hormones, and from a conjectural relationship of the ductless glands to other parts of the visceral nervous system corresponding to the proved association between the adrenals and the sympathetic, have arisen curiously intricate statements of function. They are chiefly taught by Falta and other clinical investigators in Vienna, and they are quoted repeatedly in Biedl's book.

Thus (p. 420) removal of the pancreas removes an inhibitory hormone which it is supposed to form, and consequently the thyroid finds the tissues freed from this constraint and can exert without hindrance its power to accelerate metabolism by its own dissimilatory hormone. Reference will be made again later to this attractive study of the balancing of the body within as well as without, the view that just as our equipoise in standing or in walking depends on the adjustment of nervous impulses, motor and also, as Sherrington has so brilliantly shown, inhibitory towards the many muscles concerned in the posture, so, too, the equilibrium of metabolism and the delicate adjustments of growth, that so marvellously round the whole into the symmetry of adult life, are evidence that the balance of power is fairly adjusted in the concert of the ductless glands, and that no hormone precludes his fellow from a place in the sun of vital growth.

The theory enunciated by Falta and Rudinger as to the actual function of the thyroid secretion is this. Thyroid and parathyroid oppose one another; the former stimulates the sympathetic nerves, the latter inhibits them. Pancreas is associated with parathyroid in retarding metabolism, while thyroid and adrenals augment it. Hence loss of the thyroid leads to:—

- (1) Loss of its secretion, and therefore slowed metabolism.
- (2) Absence of stimulation of the adrenals.
- (3) Escape of the pancreas from inhibition.
- (4) Decreased excitability of the nerve which is supposed to innervate the thyroid, that is the vagus, in all its area of distribution.

All this is visionary, and a discussion of it would be profitless at present. More facts are wanted.

PLURIGLANDULAR INSUFFICIENCY.

Increasing emphasis has been laid in the last few years on the need to look upon the ductless glands as parts of a connected whole, so that one gland can neither fall out by the wayside nor grow tenfold in bulk without the others showing consequent change. Careless clinical analysis is apt to accept this readily, and, beneath the ugly phrase of "polyglandular syndrome," to cover ignorance of the cause of pigmentation, of obesity, glycosuria, or gigantism by a general reference to some one or other in the ring of ductless glands, of which nothing is known with certainty.

But the original view of the matter has good justification. The ductless glands do bear some functional relationship one to another;

normal health and growth swing in the equal balance of their activities, and damage of one is likely to result in altered action of another, so that the ultimate bodily change shows the plus of B as well as the minus of A. To name a few common features, which suggest a common bond :—

(1) Carbohydrate metabolism is influenced by all. On ordinary diet sugar appears in the urine when the pancreas is damaged; in exophthalmic goitre, when the thyroid is over active; in acromegaly, that is presumably with superpituitarism; and lastly, by injection of adrenalin or stimulation of the splanchnic nerves to the adrenals. On the other hand, sugar tolerance is exhibited by patients with subglandular conditions of pituitary, thyroid, or adrenal. The correlation may be explained directly or indirectly. Thus in exophthalmic goitre the state of nervous fright may be exciting the adrenals through the splanchnics and so leading to glycosuria; that is, the fright is due to the thyroid, and the glycosuria is a secondary expression of this.

(2) Bodily growth and general metabolism. With the development of the testicle, especially of its interstitial cells, the individual is consolidated and longitudinal growth checked. This process is accelerated to an amazing degree when the cortex of the adrenal is hypertrophied; it is arrested in the dwarfish cretin with atrophied thyroid, and sexual ability fails alike with the acromegalic giant and with obese examples of subpituitarism.

(3) Nervous implications are less obvious. Tetany is a result of sudden thyroid or parathyroid loss. There is nothing corresponding to the clear connexion of the adrenal medulla with the sympathetic nervous impulses.

(4) Association between the glands themselves. The pituitary becomes hypertrophied when the thyroid is removed. Acromegaly frequently leads, in its later stages, to enlargement of the thyroid. Thyroid extract is found to be a useful remedy in acromegaly, even for the amblyopia of that disease.

(5) The double anatomical constitution of all the glands is a morphological chance that does not enter into the question. So, too, no stress can be laid upon the fact that an active principle of simple molecular constitution, which is not destroyed by boiling, can be extracted from the adrenal, the pituitary, and apparently from the thyroid and pancreas.

The adrenal cortex and the interstitial cells of the testicle secrete a special fatty substance; the thyroid and pituitary a special colloid. But the lipoid and the colloid are probably different substances in each case. Certainly that of the pituitary does not contain iodine.

(6) Morphological relationship. This argument, when it can be applied, is one of great weight to emphasize a likeness of function. But the work done in comparative physiology and that in morphology is accumulated in separate islands of thought, and few have sought to cross the estranging sea. Gaskell, on morphological grounds [6] classifies together the pituitary, thyroid, and adrenal cortex as being all modified from the coxal glands, that is the segmental excretory apparatus of some primitive arthropod type. This suggests that these ductless glands may be found to have physiological activities that are more closely related to each other than some workers would at present care to admit.

Indeed the question of the relationship to one another of the ductless glands is one of subtle intricacy, which needs the clearest thought and more proved facts for a right judgment to be made. Hypotheses serve to direct the search this way and that, and through them more evidence is brought to light and tested. But confusion will disorder all, if theory and analogy be permitted to develop a statement and this be too early accepted as conclusive. The teaching of the Viennese school, despite its brilliant suggestiveness, carries with it that dangerous fault, for it has been elaborated far beyond the facts. There is not a shadow of proof that the thyroid has any special and peculiar relation to the vagus nerve, like that of the adrenals to the sympathetic. Drugs, such as pilocarpine and atropine, cannot be classified with regard to their action upon different departments of the nervous system as rigidly as can adrenalin. Rightly to explain clinical observations is hard enough, even when one uses the proved truths of physiology and pathology; error is unavoidable, if the deduction is made from error. The apparent simplicity of a generalization will always recommend it to medicine, impatient for aid to comprehend the instant problems of disease. But the speculative nature of Falta's views should be properly appreciated, lest on them too much be built and much weary work of reconstruction be called for later, if the foundations fail.

The essential facts and the broad laws concerned with the activities of the visceral nervous system have been mainly ascertained by experimental work in England. As it chanced, this study of the visceral nerves was almost hindered in its progress by the necessity of devising at each step experiments to demonstrate the error of a general law which had been formulated in Vienna by von Basch, the law of crossed innervation of the longitudinal and circular coats of plain muscle in hollow viscera. The law was absolutely unproven, but it was

so engagingly simple that it was accepted without question as it stood, to be the working creed of many medical writers. Its removal cost much labour, and now the cleared ground is threatened afresh. Medicine owes no debt of gratitude to those who teach to her theories without proof.

Still these views, and particularly Biedl's book, will do good service, insomuch as they emphasize the problems to be solved. We see how widely and deeply the influence of the ductless glands penetrates through the busy factories of life. But the how and the why of it all is chiefly a mystery still. The reading of the riddle must increase man's power to help man.

T. R. ELLIOTT.

REFERENCES.

- [1] BULLOCK and SEQUEIRA. *Trans. Path. Soc. Lond.*, 1905, vol. lvi, p. 189.
  - [2] CUSHING, WEED and JACOBSON. *Johns Hopkins Hospital Bulletin*, February, 1913.
  - [3] DALE. *Biochemical Journ.*, 1909, vol. iv, p. 427.
  - [4] ELLIOTT and ARMOUR. *Journ. Path. and Bact.*, 1911, vol. xv, p. 481.
  - [5] GASKELL. *Journ. Physiol.*, 1886, vol. vii, p. 1.
  - [6] *Idem*. "The Origin of Vertebrates," London, 1908, p. 430.
  - [7] HERRING. *Quart. Journ. Exp. Physiol.*, 1908.
  - [8] HORSLEY and HANDELSMANN. *Brit. Med. Journ.*, November 4, 1911.
  - [9] MACCALLUM and VOIGHTLIN. *Johns Hopkins Hospital Bulletin*, March, 1908.
  - [10] SCHÄFER. Croonian Lecture, *Proc. Roy. Soc.*, 1909, vol. lxxxi, p. 442, and *Proc. Roy. Soc. Med.* (Neurol. Sect.), March, 1913.
-