Ovariectomy Followed by Ectopic Pregnancy in Stump.

By HAROLD CLIFFORD, M.B. (Lond.), F.R.C.S (Edin.), Manchester.

THE patient was a married woman, aged 26 years, who had had two living children, and possibly a miscarriage, for which she was curetted in May, 1913. In September, 1913, she had all the signs and symptoms pointing to an ectopic pregnancy on the right side, but on opening the abdomen the tumour was found to be a small cystic tumour of the right ovary, with no sign of pregnancy; this was removed together with the tube. The left appendages were apparently healthy. She made an uninterrupted recovery.

In October she menstruated for five days without pain; she missed the November period, however, but commenced to lose on December 20th, being then three weeks over her usual time; at the time she complained of abdominal pain and faintness. The pain became worse, especially on the left side, and caused her to faint completely on two occasions. About this time I saw her in the out-patient department of St. Mary's Hospitals, Manchester, and found she had a fresh swelling in the pelvis, roughly the size of a three months' pregnancy; she, however, denied the possibility of pregnancy, and as she was going downhill I decided to reopen the abdomen. On doing so I discovered an ectopic pregnancy, apparently in the stump of the previous ovariectomy, but owing to that operation it was difficult absolutely to define its relations to the Fallopian tube and round ligament; there was also a small ovarian cyst on the left side. I removed the cyst together with the uterus and ectopic sac, and the patient made a speedy recovery.

Dr. Shaw, who examined the specimen microscopically for me, found chorionic villi in the sac.