Datum	M. E. Z.	Luft	Stufenschätzungen
1910 Sept. 29	11 ^h 5 ^m	sehr gut	<i>b</i> 4 <i>RY</i> , <i>RY</i> kaum zu sehen
Okt. 5	10 50	»	<i>b</i> 4 <i>RY</i> , <i>RY</i> kaum zu sehen
Nov. 7	9 16	»	<i>a</i> 2 <i>RY</i> <i>b</i>
Dez. 21	8 35	gut	<i>c</i> 2 <i>RY</i> 3 <i>b</i>

SS Geminorum.

Vergleichsterne: *a* = BD + 22° 1178 Gr. = 8^m8
b = » + 22 1188 9.2
c = » + 22 1193 9.2

1910 Febr. 28	10 35	gut	<i>a</i> 1 <i>SS</i> 3 <i>b</i>
März 8	10 40	»	<i>a</i> <i>SS</i> 3 <i>b</i>
» 9	9 28	»	<i>a</i> <i>SS</i> 4 <i>b</i>
» 14	9 42	»	<i>a</i> <i>SS</i> 3 <i>b</i>
» 21	9 0	»	<i>a</i> 1 <i>SS</i> 3 <i>b</i>
» 30	9 51	»	<i>a</i> 2 <i>SS</i> 2 <i>b</i>
April 1	10 30	»	<i>a</i> 2 <i>SS</i> 2 <i>b</i>
» 11	9 36	sehr gut	<i>a</i> 1 <i>SS</i> 2 <i>b</i> , <i>SS</i> 2 <i>c</i>
» 28	10 13	»	<i>a</i> 1 <i>SS</i> 2 <i>b</i>

ST Geminorum.

Vergleichsterne: *a* = BD + 35° 1666 Gr. = 8^m3
c = » + 35 1669 8.7
d = » + 35 1664 9.3
b = 7^h 29^m5 + 34° 50' —

1910 Febr. 28	10 30	gut	<i>a</i> 4 <i>ST</i> 4 <i>b</i>
März 8	10 36	»	<i>a</i> 2 <i>ST</i> 5 <i>b</i>
» 9	9 24	»	<i>a</i> 2 <i>ST</i> <i>c</i>
» 14	9 39	»	<i>a</i> 2 <i>ST</i> <i>c</i> , <i>ST</i> 4 <i>b</i>
» 21	8 58	»	<i>a</i> 2 <i>ST</i> 5 <i>b</i> , <i>c</i> 1 <i>ST</i>
» 30	9 49	»	<i>a</i> 3 <i>ST</i> 3 <i>b</i> , <i>c</i> 3 <i>ST</i>
April 1	10 28	mäßig	<i>a</i> 3 <i>ST</i> 3 <i>b</i>
» 11	9 32	sehr gut	<i>a</i> 4 <i>ST</i> 3 <i>b</i> , <i>c</i> 3 <i>ST</i>
» 28	10 8	»	<i>d</i> 2 <i>ST</i> <i>b</i>
Mai 12	10 12	gut	<i>b</i> 2 <i>ST</i> , <i>ST</i> kaum zu sehen

Mundenheim, Pfalz, 1911 Mai.

Datum	M. E. Z.	Luft	Stufenschätzungen
-------	----------	------	-------------------

SV Tauri.

Vergleichsterne: *a* = BD + 28° 920 Gr. = 8^m7
b = » + 27 895 9.0
c = » + 28 917 9.0
d = » + 28 922 9.5

1910 Febr. 28	10 ^h 55 ^m	gut	<i>a</i> 2 <i>SV</i> <i>b</i>
März 8	10 50	»	<i>a</i> 2 <i>SV</i> <i>b</i>
» 9	9 18	»	<i>a</i> 2 <i>SV</i> <i>b</i>
» 14	9 34	»	<i>a</i> 2 <i>SV</i> <i>b</i>
» 21	8 53	»	<i>a</i> 2 <i>SV</i> , <i>b</i> 1 <i>SV</i> <i>c</i>
» 30	9 44	»	<i>b</i> 3 <i>SV</i> , <i>c</i> 3 <i>SV</i> <i>d</i>
April 1	10 24	»	<i>b</i> <i>SV</i> , <i>a</i> 2 <i>SV</i> 2 <i>d</i>
» 11	9 28	sehr gut	<i>a</i> 2 <i>SV</i> 2 <i>d</i> , <i>b</i> 1 <i>SV</i> <i>c</i>

Z Cancri.

Vergleichsterne: *a* = BD + 15° 1809 Gr. = 8^m5
b = » + 15 1802 9.0

1910 Febr. 28	11 20	gut	<i>a</i> 3 <i>Z</i> 2 <i>b</i>
März 8	11 13	»	<i>a</i> 3 <i>Z</i> 1 <i>b</i>
» 9	9 55	»	<i>a</i> 4 <i>Z</i> 2 <i>b</i>
» 14	10 4	»	<i>a</i> 3 <i>Z</i> 2 <i>b</i>
» 30	10 6	»	<i>a</i> 3 <i>Z</i> 2 <i>b</i>
April 1	10 46	»	<i>a</i> 3 <i>Z</i> 2 <i>b</i>
» 11	9 50	sehr gut	<i>a</i> 3 <i>Z</i> 2 <i>b</i>
» 28	10 28	gut	<i>a</i> 4 <i>Z</i> 2 <i>b</i>
Mai 12	10 19	»	<i>a</i> 3 <i>Z</i> 2 <i>b</i>

Ferner waren folgende Variable an sämtlichen Beobachtungstagen des Jahres 1910 (s. Klammer) unsichtbar, also < 12^m:

TV Herculis (Mai 12, Juli 28, Sept. 10, 25, 28, Okt. 5).

SZ Persei (Febr. 28, März 8, 9, 14, 21, 30, April 11, Okt. 6, Nov. 7, Dez. 21).

RT Cephei (Mai 12, 13, Juli 24, Sept. 6, 15, Okt. 6, Nov. 7, Dez. 21).

SY Draconis (März 30, April 10, 30, Mai 12, Sept. 6, 29, Okt. 5, Nov. 7, Dez. 21).

Max Mündler.

Horary number of meteors visible on every night of the year. By W. F. Denning.

The following table exhibits the hourly frequency of meteors. The numbers are deduced from data obtained at Bristol and relate to the mean of a. m. and p. m. observations and are for one observer watching a clear, moonless sky uninterruptedly. The results on which the table is founded were gathered during the years 1866 to 1911. There being no leap-year in 1900 and the great majority of the observations being taken before that year the dates may be regarded as one too early. Thus the Perseids at the present epoch will supply about 69 per hour on Aug. 11 not on Aug. 10 as formerly, for the maximum of that display is now on Aug. 11 or 12.

In various years the number of meteors appears to vary not only on individual nights but generally during a whole year. I saw many more meteors in 1877 than in

1887 but it is true my place of observation was better in the former year.

My materials though pretty abundant are not sufficiently full for an exhaustive investigation and perfectly reliable results. However I believe the table will furnish a useful guide and be found approximately correct. It must be remembered that no two observers will obtain accordant observations. *Zenzioli* in the beautiful skies of Italy and *Tupman* in the Mediterranean saw a surprisingly large number of meteors while *Schmidt* at Athens and others appear to have recorded comparatively few, so that horary rates computed from the experiences of different observers must necessarily vary considerably. The values here given are fairly correct for my eye and for the circumstances under which I pursued the observations.

I have taken no account of the great meteoric showers which I witnessed on 1866 Nov. 13, 1872 Nov. 27 and 1885 Nov. 27 as the thousands of meteors appearing in such very exceptional displays would swell the numbers far beyond ordinary experience. The idea in forming the table was to show the horary number of meteors to be expected on ordinary nights of the year when the conditions are favourable for viewing these phenomena. In or near large towns where artificial illumination affects the sky, the frequency of meteors is not nearly so well marked as in country districts where darker skies enable much smaller objects to be observed.

	Jan.	Feb.	Ma.	Apr.	May	Jun.	July	Aug.	Sept.	Oct.	Nov.	Dec.
1	18	8	7	7	7	6	8	20	15	13	12	9
2	28	7	6	8	6	5	7	19	14	14	12	9
3	12	7	7	8	6	6	8	19	13	14	13	10
4	9	6	8	7	7	6	9	21	12	14	14	11
5	10	6	7	7	6	7	9	22	12	12	12	12
6	9	6	6	8	5	6	8	24	14	13	13	13
7	8	7	7	6	5	6	8	27	13	12	13	12
8	7	7	7	7	6	6	9	34	12	14	14	14
9	7	7	7	7	6	7	9	48	12	13	14	16
10	7	6	8	8	6	6	10	69	13	14	15	22

Bristol, 1911 June 24.

	Jan.	Feb.	Ma.	Apr.	May	Jun.	July	Aug.	Sept.	Oct.	Nov.	Dec.
11	8	6	8	8	6	6	10	48	12	14	16	23
12	8	7	7	7	6	7	11	30	13	15	17	19
13	8	7	8	7	7	6	11	22	14	16	20	16
14	9	7	9	8	6	6	12	20	15	17	21	13
15	8	8	8	7	6	7	12	18	15	21	18	10
16	9	8	9	6	5	7	13	17	15	20	17	10
17	9	8	10	6	5	8	13	16	14	21	16	9
18	9	8	9	7	5	7	14	15	13	21	15	8
19	9	9	8	8	6	6	15	15	14	20	14	9
20	10	9	8	9	6	8	16	16	15	19	15	8
21	9	9	7	10	6	8	17	17	17	18	16	10
22	8	8	7	8	7	6	18	19	16	17	15	10
23	7	8	8	7	7	7	19	19	15	16	16	8
24	6	8	8	6	6	7	20	20	14	16	14	8
25	6	8	8	6	6	7	21	21	13	15	15	9
26	7	7	7	5	6	7	23	20	14	14	14	8
27	7	7	7	6	5	7	25	19	15	15	15	9
28	8	7	8	5	6	6	27	20	13	16	14	9
29	7		8	6	6	6	26	19	14	15	13	10
30	8		9	7	7	7	23	18	15	14	12	9
31	7		8		8		21	16		14		10

W. F. Denning.

On the orbit of $\Sigma 2525$. By W. Doberck.

$\Sigma 2525$ was noted single by *Schiaparelli* in 1887, 1888, and 1889 and also by *H. Struve* in 1888, but in 1886 *Schiaparelli* noted the companion in $208^{\circ}7$ at a distance of $0''.22$, a position approximately agreeing with his previous observations. The following orbit gives for 1886.92 $\theta = 113^{\circ}7$, $\rho = 0''.06$. *Schiaparelli* probably mistook some small speck of light for the companion. The motion is retrograde. The anomalies are taken negative before and positive after the peri-astron passage. The hypothetical parallax is $0''.024$. Precession has been neglected. Abbrevi-

ations of names are explained in A. N. 4346: Gls means *Glaisher*.

$$\begin{aligned}\Omega &= 9^{\circ} 0' & P &= 243^y 9 \\ \lambda &= 274^{\circ} 23' & T &= 1887.09 \\ \gamma &= 132^{\circ} 42' & a &= 0''.952 \\ e &= 0.9175\end{aligned}$$

$$\rho \cos \theta = +0''.0264 + 0''.3770 \sin(E - 4^{\circ} 23')$$

$$\rho \sin \theta = -0.5940 + 0.6486 \sin(E + 86^{\circ} 30')$$

$$\pm z = +0.6401 + 0.6981 \sin(E - 88^{\circ} 15')$$

Ephemeris.

Year	θ	ρ	z	Year	θ	ρ	z	Year	θ	ρ	z
1912.50	306.0	0''.686	0''.6623	1920.50	300.6	0''.788	0''.7932	1928.50	296.4	0''.876	0''.9032
1914.50	304.5	0.713	0.6973	1922.50	299.5	0.810	0.8225	1930.50	295.5	0.895	0.9278
1916.50	303.1	0.739	0.7306	1924.50	298.4	0.833	0.8505	1932.50	294.6	0.912	0.9514
1918.50	301.8	0.764	0.7626	1926.50	297.4	0.855	0.8774	1934.50	293.8	0.930	0.9743

Obs.	1800 +	θ_c	ρ_c	$\theta_o - \theta_c$	$\rho_o - \rho_c$
Σ	30.43	255.6	1''.04	+0.3	+0''.29
Σ	36.14	253.6	0.99	+1.9	+0.31
Da	40.62	251.8	0.96	+3.7	+0.29
$O\Sigma$	.70	»	»	-1.6	+0.17
Gls	41.70	251.4	0.95	-0.7	+0.08
Mä	42.41	251.1	0.94	-0.1	-0.12
Gls	.73	251.0	»	-0.5	-0.12
»	43.61	250.6	0.93	+4.4	0.00

Obs.	1800 +	θ_c	ρ_c	$\theta_o - \theta_c$	$\rho_o - \rho_c$
Mä	43.69	250.6	0''.93	+3.4	+0''.02
Gls	44.60	250.2	0.92	-2.1	0.00
$O\Sigma$	54.63	245.5	0.82	+1.6	+0.18
Se	56.61	244.3	0.79	+2.8	+0.06
A	58.39	243.3	0.77	+15.4	-
A	64.74	238.8	0.68	+2.5	-0.13
En	65.48	238.3	0.67	+15.3	+0.06
A	.58	238.2	»	+2.4	-0.04