has shown that arsenic is eliminated after intravenous injection for four days, and in cases of intramuscular injection arsenic is still present up to the twelfth day. Between the first and second injections of salvarsan the patients have been ordered a mercurial inunction, this being continued after the second injection for 20 days in each month for a period of six months. The preparation prescribed is "crème de mercure" (Gordon), as it is easily absorbed and is more cleanly than the official ointment.

None of the cases so treated has had a relapse or has shown symptoms of syphilis, but although at present their Wassermann reaction is negative it is impossible until further time has elapsed to regard them as cures.

We have had no serious accidents to report. In one case soon after the second injection there was an eruption of herpes round the chin and lips and on the tongue, which yielded to the ordinary treatment. Some of the cases presented slight induration at the site of the injection which soon disappeared.

In women we observed that the regular period following the injection appeared a week earlier than usual, and was sometimes profuse in character. In one patient with severe ulceration of the leg iritis occurred, but it was not due to the treatment. The dose administered has varied from 0.4 to 0.6 gramme, according to the condition of the patient.

The injections were done in three ways:—(1) *Subcutaneous*.—This method was quickly discarded as it proved too painful. (2) *Intramuscular*.—(a) In neutral solution; this method also proved painful and caused a good deal of induration. (b) In the form of an oily emulsion; this proved more satisfactory. (3) *Intravenous*.—This method if properly performed is much the best way of administering salvarsan, except in those exceptional cases where it is impossible to find a suitable vein.

The apparatus we used in the intravenous method consisted of a glass cylinder of a capacity of 300 c.c. To this is attached indiarubber tubing fitted with two glass windows. The whole having been sterilised by boiling, a cannula and needle (previously sterilised in oil) are then attached. The apparatus having been filled with saline solution, it is impossible for air to enter at any time during the operation. The salvarsan solution must be prepared *immediately* before use and kept at a temperature approximating blood heat during administration. Chemically pure sodium chloride and caustic soda must be employed for the solutions. The technique is as follows.

Into a narrow-necked graduated glass cylinder of a capacity of 250 c.c., containing about 40 sterile glass beads, 40 c.c. of distilled water at a temperature of 104° F. are measured. The salvarsan, 0.6 gramme, is then gradually added, the tube being vigorously shaken to aid solution. The solution must not show (even when examined by transmitted light) any drop-like or gelatinous particles. To this are added 20 drops of a 15 per cent. caustic soda solution. A precipitate forms which redissolves on the addition of a few more drops of caustic soda solution. The clear yellow liquid is then filled up to 240 c.c. with a 0.5 per cent. saline solution prepared from sterile freshly distilled water. Each 40 c.c. of the solution now contain 0.1 gramme of salvarsan, 200 c.c. contain 0.5 gramme, so the dose required may be accurately ascertained.

An elastic band having been placed on the upper part of the arm the most prominent vein is selected at the bend of the elbow. The site of the puncture is then painted with iodine solution. The apparatus being filled with saline solution care must be taken that all air is expelled. The needle is then run into the vein, when blood will be seen to flow back by the glass window. The elastic bandage is then removed and the saline solution allowed to pass into the vein. If this takes place readily with no swelling at the site of the injection, the salvarsan solution is then slowly added to the saline until the required amount is given. Saline solution is again added until the liquid entering the vein is quite clear.

If swelling occurs at the site of injection it is caused by the needle having pierced the side of the vein; in that case it is advisable to remove the needle and select another vein. The presence of salvarsan solution in the tissues will cause painful induration and sloughing might occur. It may happen that the liquid does not flow freely, the point of the needle having come in contact with the wall of the vein;

this is obviated by either raising or depressing the point of the needle.

There should be no local reaction. A general reaction nearly always occurs and varies in intensity according to the condition of the patient and the amount of salvarsan given. During the injection the cheeks generally become pink and some patients complain of a metallic taste in the mouth. An hour or two after the injection there is likely to be a rigor and the temperature may rise to 101°–102° F., with, perhaps, headache and vomiting. This condition quickly passes and the patient feels well soon afterwards. During the last few months we have observed that cases of vomiting have occurred less frequently. Whether this is due to an improvement in technique or the salvarsan supplied by Ehrlich being purer we cannot say. It is hoped by the particular combination of salvarsan and mercury to attain a means of preventing the development of tabes and paralysis.

There can be little doubt that Professor Ehrlich by his recent work has added a most valuable drug to our Pharmacopœia.

CLASSIFICATION IN GYNÆCOLOGY.¹

By W. E. FOTHERGILL, M.A., B.Sc., M.D. EDIN.,
HONORARY ASSISTANT GYNÆCOLOGICAL SURGEON, ROYAL INFIRMARY
AND ST. MARY'S HOSPITALS, MANCHESTER; CLINICAL LECTURER
IN GYNÆCOLOGY, UNIVERSITY OF MANCHESTER.

CLASSIFICATION marks the transition of any subject of inquiry from being a mere collection of observations and theories into being a science. In other words, no subject becomes a science until it is sufficiently well known to admit of the logical and orderly arrangement of its details. If this test be admitted, gynæcology is still a very young science, if, indeed, it has yet reached the scientific stage. As an art it has, of course, reached a high level.

The first attempts to give an orderly account of the diseases of women in this country seem to have been made in the later third of the nineteenth century. Thus the third edition of West's "Lectures," published in 1864, and the third edition of Grailey Hewitt's book, published in 1872, both contain traces of arrangement into diseases of the vulva, vagina, uterus and other organs. In 1877 Lawson Tait published a small book with a definite anatomical classification, and the same arrangement was used by Barnes, whose second edition appeared in 1878. From this date until the present time most books on the diseases of women have been arranged on the same plan, though some are conspicuously free from any attempt at logical classification. The object of the present paper is to examine the nature and results of this anatomical classification, and to find out if it is the best that can be made now, as it doubtless was when it came into use.

Though it may have been useful as a "preliminary effort," it is not only an artificial division, but a bad one of its kind, and it has been made worse by faulty application. For a number of conditions which are not conditions peculiar to women have been brought in. Such are the various skin diseases described under the headings "Vulva" and "Mons Veneris"; symptoms like pruritus, and affections of the bladder, urethra and rectum. Naturally the gynæcologist has to treat them; but when he is trying to give a scientific account of diseases of the female reproductive organs he might at least relegate other matters to an appendix. Conditions do not always appear in the appropriate anatomical division. Thus cystocele and rectocele are often called diseases of the vagina, prolapse and retroversion being placed amongst diseases of the uterus, while all four are really abnormalities of the pelvic connective tissue. These are all avoidable errors in the application of the system.

But the division itself is to blame for splitting up and multiplying diseases. Gonorrhœa, for example, is one condition; but in books it is described in different sections under vulva, urethra, vagina, cervix, uterus, tubes and ovaries. Thus facts which should be together are scattered abroad. Further, local pathological processes are made into separate diseases, and names are given to them such as

¹ A paper read before the North of England Obstetrical and Gynæcological Society on Feb. 17th, 1912.

vaginitis, endocervicitis, endometritis, metritis, parametritis, perimetritis, salpingitis, ovaritis. These are described, each with its symptoms, signs, diagnosis, prognosis and treatment. "It is a dull and obtuse mind," said Calvin, "that divides in order to distinguish; but it is a still worse which distinguishes in order to divide." The older gynæcological writers may not have distinguished between ovaritis and salpingitis with the conscious intention of dividing them, but their writings have effectively separated these processes in the minds of their followers. Nor can the use of the anatomical system be said to be successful in practice. Senior students and young graduates constantly remark that while they can make useful progress in medicine, surgery and midwifery, they are in despair about the diseases of women. Interest, intellect and industry are not wanting; the method is alone at fault. The subject presents no unique mystery. It is badly taught simply because writers and teachers persist in using the anatomical arrangement. How can the student grasp the connexion between vulvitis and salpingitis when they are separated by hundreds of pages in text-books and by the lapse of weeks in lecture-rooms? How can he help regarding ovaritis as a disease when it is described as one? How can he escape the conclusion that prolapse is a uterine trouble when it is placed among diseases of the uterus? Various other bases might be taken for the main divisions of a new system, but what is required is a natural arrangement.

The word "cancer" at once calls to mind a number of resemblances between all cases of cancer; but the term "disease of the uterus" does not suggest to the mind any resemblances between diseases of the uterus except that of place. The word "infective" conveys a number of facts concerning all cases of infection; but the word "vagina" tells nothing about any case of disease of the vagina except the site of the lesion. In short, the knowledge that a condition affects the broad ligament, the uterus or the ovary conveys much less information than the knowledge that the condition is an injury, a tumour or a malformation. In general terms, the important feature of a disease is not its site, but the way in which it departs from health—namely, its pathology. The word pathology must not be taken in its narrow sense of morbid anatomy, but must include function as well as structure, morbid processes as well as lesions. Does it not follow that the main divisions of any natural classification of diseases should be based on pathological resemblances and differences, minor features, such as the position of lesions, being used only in the subdivisions of the system?

Attempts to bring into use pathological main groups of diseases of women have been made in this country by four authors: David Berry Hart, J. W. Ballantyne, W. Blair Bell, and the present writer. In 1893 Hart read a paper to the Edinburgh Obstetrical Society on the Pathological Classification of the Diseases of Women, with a Plea for the Revision of Current Views.² He suggested 11 headings as follows: (1) Congestive (vascular ruptures included); (2) Hypertrophy; (3) Atrophy; (4) Traumatism; (5) Simple Inflammations; (6) Micro-organismal (includes pathological versions); (7) Herniæ (prolapse, &c.); (8) New Growths; (9) Displacements (inversion only); (10) Developmental Errors; and (11) Neuroses.

Ballantyne published his "Essentials of Gynæcology" in 1905. His classification of the diseases of women was in 10 headings. He put hypertrophic and atrophic states together, although they are opposite in nature. He included extra-uterine pregnancy, which is midwifery; and also sterility and menstrual anomalies, which are symptoms rather than primary conditions. Hart and Ballantyne both include "neuroses," which would generally be regarded as disorders of the nervous system, rather than of the female reproductive organs.

In his "Principles of Gynæcology," published in 1910, Dr. Blair Bell also uses ten headings. One of these is "Disorders of the Normal Physiological Conditions in Relation to Menstruation," and this, like Ballantyne's "Menstrual Anomalies," deals with symptoms, not with primary conditions. Another heading is "Disorders of the Normal Conditions in Relation to Conception," which, again, is midwifery. Various conditions which are not primary disorders of the

female reproductive organs are included under the heading, "Allied Morbid Conditions."

Thus Ballantyne's and Blair Bell's arrangements are not faithful to their main basis of division, and they poach on obstetrical ground. Hart's plan, if the heading "neuroses" be omitted, is free from these defects; but it suffers with the others from the fact that the student cannot easily keep in mental view so many as ten groups of conditions.

For some years I have been using six main divisions both in clinical teaching and in my own systematic work.³ They were suggested by the arrangement of specimens introduced by Professor Lorrain Smith into the pathological museum of the University of Manchester, and thus have the sanction of professional pathologists. They are: (1) Developmental Errors; (2) Vascular Changes; (3) Mechanical Conditions; (4) Results of Infection; (5) Progressive Conditions; and (6) Retrogressive Conditions.

1. "Developmental Errors" must, for gynæcological purposes, contain two sub-groups: (a) congenital errors; and (b) errors in development at puberty, for the reproductive organs have two phases of development, one embryonic and foetal, the other occupying the years immediately preceding and following puberty.

2. "Vascular Changes" includes active and passive hyperæmia as recognised at first by symptoms and later by hypertrophy and fibrosis respectively. This group also includes varicosity of the veins in the pelvis and hæmorrhage into the pelvic organs, the pelvic cellular tissue, and the peritoneal cavity.

3. "Mechanical Lesions" is a large and obvious group containing injuries, displacements, and herniæ.

4. "Results of Infection" is a main division which should be subdivided according to the organisms present in each case. In this group the few cases of primary pelvic parasitic disease may be included without gross impropriety.

5. "Progressive Conditions" includes overgrowths, cysts and new growths. Perhaps this group should be divided. But it is hard to draw the line between hypertrophy and adenoma of the endometrium or between innocent and malignant new growths.

6. "Retrogressive Conditions" occur mainly in connexion with the menopause, but the reproductive organs share in the general changes of senility.

Further, there are errors in the retrogressive changes which are characteristic of normal menstruation, ovulation and involution. I have found that the use of these main groups has certain advantages. The division is logical and is based on the most important resemblances and differences between morbid conditions—namely, the pathological. It is familiar to students of medicine, and the headings are so few that they can easily be remembered. If the student is able to place a case in one of these groups, he has made a definite step towards a complete diagnosis and has gained some useful information about the patient. But it is, of course, a division of conditions, not of patients, for a woman can have two or more conditions at the same time (infection and new growth, for example).

Such a division has the further advantage that it does not magnify symptoms and pathological processes into diseases. Nor does it lead to confusion between primary conditions affecting the reproductive organs and general diseases such as phthisis, rickets, anæmia, and myxœdema, which often lead to the appearance of gynæcological symptoms. Such symptoms of general diseases may well, like diseases of the female bladder, urethra, and rectum, be mentioned by the gynæcologist in teaching; but it is better that they should be kept out of any classification of diseases of the female reproductive organs which aims at being scientific and logical.

Those who have written text-books and arranged courses of lectures according to the anatomical plan can hardly be expected to approve of the changes now advocated, though the use of pathological main groups saves much space and prevents much repetition. Again, students taught to use a pathological main division, unless well informed and agile in mind, might easily fail to please examiners of the anatomical generation. Thus mental inertia and the fear of the examiner tend to perpetuate the anatomical grouping. But it cannot last much longer, and the writer hopes to see the day when it will not be used in any course of lectures delivered in this country nor in any book which holds a place on the British market.

Manchester.

² Transactions of the Edinburgh Obstetrical Society, vol. xix. and vol. ii. Edinburgh Hospital Reports, 1894.

³ Manual of Diseases of Women. William Green and Sons. 1910.