may not find it easy to come to terms with a doctrine which rests upon the dogma—often beautifully illustrated but never substantiated by the facts—that perception depends upon organic movement; which ignores two or three competent accounts of introspection; and which uses the term “stimulus” almost as irresponsibly and variously as Dewey himself. The application to sensation of Professor Washburn’s notion of attention as “suspended reaction” suggests a fruitful tempering of the behaviorist’s doctrine.

The perennial problem of the relation of sensation to “sense data” and to physical objects is discussed by Broad (1) in his article on phenomenalism. The article is a criticism of B. Russell’s position on knowledge of the external world. It has much more to say of the “sensory data of knowledge” than of the sensations of analytical psychology. Articles by Nunn (4), and Moore and Stout (3) are of the same sort. Robinson (5) traces the history of the doctrine of sensation and perception from Condillac to Maine de Biran, who gives it a voluntaristic turn by basing the perceptual functions upon effort and will. Spiller (6) finds no present evidence that the different forms of sensations are separately represented in the internal capsule, though he thinks that they probably are separated in the parietal lobe, each sensory area lying adjacent to a corresponding motor area.

REFERENCES

1. BROAD, C. D. Phenomenalism. *Proc. of Aristot. Soc.*, 1914-15, 15, 227-251.
2. DE LAGUNA, G. A. Sensation and Perception. *J. of Phil., Psychol., &c.*, 1916, 13, 533-547, 617-630.
3. MOORE, G. E., & STOUT, G. F. The Status of Sense-data. *Proc. of Aristot. Soc.*, 1914, 14, 355-406.
4. NUNN, T. P. *Proc. of Aristot. Soc.*, 1915-16, 16, 156-178.
5. ROBINSON, A. The Philosophy of Maine de Biran; the Way Out of Sensationalism. *Proc. of Aristot. Soc.*, 1914-15, 15, 252-270.
6. SPILLER, W. G. Remarks on the Central Representation of Sensation. *J. of Nerv. & Ment. Dis.*, 1915, 42, 399-418.

VISION—GENERAL PHENOMENA

By EDWIN B. HOLT

Harvard University

During the past year the most important work on this subject has been done not in Germany nor in Europe, but in America, and not in the colleges or universities, but in those now rather numerous

laboratories which have been established, and which are maintained, by companies that are engaged in illuminating, photographic, and other similar industries. The introspective and "purely psychological" points of view figure on the whole less prominently, and in their stead we find a more competent physical and mathematical handling of the problems than has hitherto been the rule.

One of the most comprehensive papers of the year is that of Troland on "apparent brightness" (33). He defines *light* as "radiation 'evaluated according to its capacity to produce' apparent brightness"; for, "strictly speaking, light is not the stimulus to vision, but is vision itself." The author introduces a semi-physiological light unit, the *photon*, which "is that intensity of illumination upon the retina of the eye which accompanies the direct fixation, with adequate accommodation, of a stimulus of small area, the photometric brightness of which, as determined by the standard flicker comparison and a normal subject, is one candle per square meter, when the area of the externally effective pupil, considered as lying in the nodal plane of the eye, is one square millimeter. The intensity of a visual stimulus, expressed in photons may be called its *physiological intensity*." Such a definition raises the question as to what is a normal or average eye. The answer to this question will doubtless have to come from organic chemistry: meanwhile several practical rules for discovering the observers who have "an average eye" are given by Crittenden and Richtmyer (8), who emphasize "the fact that for accurate heterochromatic measurements a systematic choice of observers is essential."

Troland further points out six stages in the (optical and) physiological process which intervenes between the impact of radiation on the outer surface of the eye and "the generation of the consciousness of light. . . . With regard to the intra-ocular adventures of this radiation we know a little; concerning retinal response, very little; and concerning the visual brain process practically nothing." As to the first, indeed, there are nearly thirty factors which must be known in order to deduce the quantitative constitution of the retinal image, and the majority of these "have not been measured even roughly." Yet it is the retinal image which is the actual physiological stimulus. There is room for exact investigations.

The author believes that retinal stimulation is "photochemical, depends on molecular or atomic resonance, and probably involves changes in the degree of ionization of substances contained in the receptors." There are two independent retinal processes, thoes

of the rods and of the cones, respectively. "The selective response of the eye to radiation of different wave-lengths (visibility curve) depends principally upon the photochemical selectivity of the retina." Both rods and cones may be electrolytic concentration cells. "Adaptation, or the general change in sensitivity to radiation of all wave-lengths, is conditioned by a change in the *concentration* of the light-sensitive substances in the retina." The retinal selectivity, or relative brightness of any radiation, probably depends on two factors—rod sensitivity and cone sensitivity. The former was measured some time ago by von Kries and others, while the latter has recently been measured by Ives and again by Nutting. The "visibility curve" of Nutting and Ives represents the relative sensitiveness of the cones to luminous radiations of different wave-lengths but of equal energy. It has one maximum, which is in the green, and it sinks rather rapidly towards the red and the violet ends of the spectrum. The curve is almost symmetrical, and Troland finds that when it is corrected for the *selective transmission* of the ocular media, macula lutea, etc., it is almost perfectly symmetrical, and approximates closely to the simple probability function.

Both Troland and Houstoun (16) have used this curve as the basis for some general remarks on color-vision. The former argues, as in substance also the latter, that "this striking symmetry of the cone visibility curve leads almost inevitably to the inference that *luminosity is due to a single photochemical process in the retina* [i. e., in the cones], and is not to be attributed to a summation of the response intensities of three or more color processes. . . . A more reasonable view lies in the original assumptions of Hering's theory, which identified luminosity with whiteness, and postulated a single, independent process for white, having a maximum in the middle of the spectrum." This view is further borne out by the fact that the following phenomena seem to be governed by intensity alone and not at all by wave-length:—the time required to produce a just-noticeable negative after-image, the time required for a stimulus to produce its maximum apparent brightness, critical flicker frequency, visual acuity, luminosity contrast and the summation of brightnesses (Ives).

Starting from the same visibility curve Houstoun (16) advances a "theory of color vision" which "does not depend on primary color sensations." "The most obvious explanation [of the curve] is to suppose that there exists in the eye a very great number of

vibrators, with a free period in the green, and that these execute forced vibrations under the influence of the light wave. The amplitude of the forced vibrations is a maximum when the free period of the vibrators coincides with the period of the incident light." These vibrators set up in the nerves waves which are transmitted to the brain. Trichromatism receives the following explanation: "The color-perceiving center [in the brain] is so badly developed that, so far as it is concerned, the curve [of the wave-lengths which reach the brain] is sufficiently specified by three points on it, provided that these points are distributed over the spectrum." The author does not "think it necessary to assume two different mechanisms," *i. e.*, rods and cones. "One system of vibrators will suffice." The word "assume" used in this connection illustrates how even to-day the physicist is apt to be ignorant of physiology, the psychologist of physics, and so forth. Von Kries (25) gives measurements of the relative amounts to which rods and cones participate in vision at various light-intensities: this is practically a correlation of the rod and the cone visibility curves previously mentioned. Under illuminations of the field varying from .01 to 25 candle meters, the proportionate part played by the rods in the seeing of gray paper ranges from .8 or .9 down to about .2. The intensity of illumination at which rods and cones participate equally varies, with different observers, from 1 to 5 candle meters. The computation involves an ingenious piece of reasoning.

There are two other papers of general scope. Ladd-Franklin (26), in a discussion of Parsons' book on "Color Vision," touches on several points of systematic interest, particularly trichromatism *vs.* tetrachromatism, and the relations between the Helmholtz, Hering, and Ladd-Franklin theories. And Goldschmidt (15), in a paper on "*Eigenlicht*," undertakes to summarize the various theories that have been advanced in explanation of the whole range of "subjective visual phenomena," and this interpreted so liberally as to include saturation, brightness, contrast, etc. These theories refer the phenomena to one or more of the seven possible sources named, which range from the dioptric apparatus to the Wundtian laws of apperception. The presentation is somewhat diffuse and academic.

Langfeld (27), in a paper which is perhaps on the border line between vision and thought-process, describes some experiments on imagery in which the subject was required to suppress some very

strong associations as for instance certain numbers in a consecutive running through of parts of the number series. The appearance and disappearance of imagery is significant. Imagery frequently "means" inhibition, and frequently it appears as a cue and the inhibitory process follows. Even very inadequate imagery can initiate and successfully guide motor response. After some practice of the *Aufgabe* imagery tended to disappear, but "when there was difficulty the image tended to appear again." Such observations, it seems to the reviewer, should be utilized in tracing the connection between immediate vision and the "higher processes."

From the introspective point of view Titchener and James Ward have argued as to the sensory character of black. In 1905 Ward expressed the opinion that black is not a "sensation"; that it is comparable to auditory silence. Titchener (31) finds that Ward does not distinguish nicely between the strictly introspective findings on "black," and such extraneous considerations as the conditions of eye stimulation and theories of vision. In his reply (41), Ward insists that black is not a "positive sensation; though a 'body-color,' i. e., a secondary quality in the epistemological sense, we must allow it to be." Titchener (32) publishes a rejoinder. The arguments are somewhat finely-spun, by both sides. It all comes down, doubtless, to the question as to whether one sees black as a sensory quality, and here probably the great majority of persons would agree with Titchener. Alspach (2) studies, by three methods, the question of "simplicity *vs.* complexity of color hues," and concludes that "however 'like' a hue may be to neighboring hues, it apparently cannot be resolved by purely psychological methods into these hues as components." It would be a mistake to accept this conclusion without an examination of the tables presented. There were many judgments of "complex" and many of "unitary," and it does not appear to the reviewer why either type of judgment should be arbitrarily minimized.

Among other qualitative papers is one of unusual interest by Swindle (30), on positive after-images of long duration. "In successive and simultaneous color induction, any color induces first itself and last of all its antagonistic color." A second positive may succeed the latter, and the duration of one or more of these phases can be remarkably prolonged by inhibiting all muscular activity, and by very brief and carefully timed renewals of the stimulation. In this way the author has obtained positive after-images of hallucinatory intensity and several minutes' duration.

In one case a positive after-image was renewed in great distinctness after forty minutes spent in a well-illuminated room. This observation is somewhat like those of G. H. Miles (1915). Swindle suggests that such after-images are utilized by spiritualistic mediums. He further describes some experiments in which an owl and a cockatoo were observed trying to adjust themselves with reference to an after-image, *e. g.*, they tried to hop on to an after-image of the perch. It was evident that the after-image appeared to move with movements of the bird's head and body. These observations are in method novel, and admirably illustrate the behavioristic point of view.

Cook and Kunkel (7) report that the simultaneous and successive contrast color for red is bluer than the true complementary (color-mixture); and for yellow and blue is redder than the complementary. The "anomaly" is enhanced when brightness contrast is eliminated. Almack and Arps (1) have studied the effect of a preceding colored stimulation on the recognition limen for color. In general, if the after-effect of the preceding stimulation is similar to the second stimulus, the limen for the latter is thereby lowered. But an inducing stimulus of any color will to some extent lower the color limen for a succeeding stimulus, and this even when the after-effect of the former is of a color complementary to the latter. Troland (40) describes an interesting "reversal" effect. One half of a circular field is exposed for 30 seconds, and then the second half is exposed as well. The longer-exposed half of course appears the less bright. If 3 seconds later the intensity of the whole field is reduced to one tenth, and then after 45 seconds more is restored to its original intensity, the longer-exposed half will appear *brighter* than the half that has been less exposed. This seems to reverse the law of fatigue. The effect can be obtained under a considerable variety of conditions, and "is present most strongly in the long-wave end of the spectrum, and with decrease in the wave-length higher and higher intensities are required to bring it out." The same author (37) has repeated, with several additional and obviously necessary precautions, the experiment with which Hering (1915) claimed to have demonstrated the Purkinje phenomenon to be fleetingly obtainable in foveal vision. With the additional precautions it was found "impossible to detect the slightest trace of a Purkinje effect when [foveal] fixation was accurately maintained." Weiss (42) describes an interesting mat woven from strips of colored papers, which is designed for demon-

strating the Purkinje phenomenon. A small white spot is fixed in the centre; and in twilight vision this is not visible foveally, while the adjacent colored squares are less distinct than those that are seen peripherally. "The foveal area tends to be filled in by imagery of the surrounding surfaces" (like the blind-spot). At that intensity of illumination for which the colors of the squares become just visible, the mat appears "to be made of mosaics which seem to be at different distances" from the observer.

Of the more exactly quantitative papers, several deal with the flicker-photometer. Ives and Kingsbury (17) add a second instalment to their (1914) theory of the flicker-photometer, and in this discuss "unsymmetrical conditions," *i. e.*, unequal periods of light and dark stimulation. This theory (for which see the earlier paper) is important, and while the authors do not claim that it is as yet entirely satisfactory quantitatively, yet "it does handle the principal phenomena qualitatively in a very striking manner." The flicker-photometer may be used in two ways. In one of these the color to be measured is alternated with black, and the speed adjusted until flicker just disappears. The speed so found is a measure of the intensity of the color. This method is "very insensitive." According to the other way, which is "very sensitive," the color to be measured is alternated with a variable and measurable white. The latter is adjusted to approximate equality with the former, and the speed then found which will just not give complete fusion; this speed is now kept. The white light is now again and more carefully adjusted, until the minimum of flicker is obtained. The intensity of the white is now equal to, and a measure of, the intensity of the colored light. "The high sensibility of the flicker photometer is shown to be due to the very rapid increase in the critical frequency of disappearance of flicker on each side of the equality setting." When the compared lights are exposed "for unequal periods, the less exposed color will be under-rated. This form of flicker photometer is less sensitive than the ordinary equal exposure arrangement."

Troland (35) says: "The flicker-photometer frequency of any two visual stimuli may be defined as the rate of alternation (cycles per second, with equal intervals) of the two in the same photometric field, which is just sufficiently rapid to eliminate flicker, when the ratio of their intensities is such as to give a minimum of flicker at a slightly lower rate." Both Troland (33, 35) and Ives (22) find this critical frequency to increase with the hue difference between

the lights compared. Troland finds that the flicker frequency curve for a white standard compared with various spectral colors shows one minimum, at about 575μ , and maxima towards each end of the spectrum. "A considerable alteration of the color tone of the standard changes the position of the minimum only slightly, so that it seems very probable that the curve depends upon the *inherent saturation* of the spectral colors at equal luminosity. . . . The addition of a white light to a saturated spectral color decreases its flicker photometer frequency very slowly." Ives (22), from in part similar measurements, is led to a different conclusion. He finds the critical frequency to be intrinsically a direct function of the "number of hue steps" between the compared lights. "If the comparison lamp is red, the minimum speed is in the red, if blue, in the blue, if yellow-white, in the yellow, but with a much higher speed at the blue than at the red end, while for a true white (5000° black body) the curve is symmetrical about a minimum in the green." And Troland (38) finds that in heterochromatic comparisons of brightness which are not based on apparent equality, but on just-noticeable inequality of brightness between the two colors, "the brightness discrimination threshold is a minimum for a minimum of color difference between the two compared stimuli," and "on the average it has a value four or five times its minimum, for a maximum of color difference."

Troland (39) further reports the critical frequency to be a direct function of the intensity of the compared lights. "If the logarithms of the frequencies for any one color are plotted against the logarithms of the intensities, the resulting curve is approximately a straight line for intensities between 40 and 1560 photons." In the Ives-Kingsbury theory of the flicker-photometer Ives (21) points out that "there figure two fractions, the brightness discrimination fraction, and the hue discrimination fraction. . . . The ratio of hue fraction to brightness fraction is about ten times larger for intermittent than for steady vision. *Herein lies the reason for the success of the flicker photometer.*" Crittenden and Richtmyer (8) deem the flicker-photometer distinctly more reliable than the equality-of-brightness photometers, whether the color differences between the lights to be measured are great or small. In (19) and (17) Ives discusses another factor, "diffusivity," which figures importantly in the Ives-Kingsbury theory. This seems to be so far an element in the theoretical equation, rather than an actually identified physiological process. For a given speed of the flicker-

photometer, an abrupt transition between the two illuminations enhances the sensitivity. But with such a transition mechanical imperfections in the apparatus can introduce serious errors. For some purposes a gradual transition is preferable. Ives (20) describes a flicker-photometer in which by means of polarizing prisms the sharp dividing edge is eliminated: one phase disappears as the other simultaneously appears, both decrease and increase exactly following the sine function.

Ferree and Rand (9) describe a method of heterochromatic photometry in which the stimulus to be measured is viewed peripherally (25 degrees) while the other field is fixated directly. Johnson (23) points out several objections to the method as described, which seem to the reviewer to be well-founded.

Troland (33, p. 952) proposes the term *minuthesis* for the "depression of a sensation under the influence of a stimulus, to replace the misleading word, 'fatigue.'" In (34) he gives measurements of the time required to produce a just discernible minuthesis for different colors at the same intensity. He also gives measurements (34, 36) of the time required for the retina to reach a condition just indistinguishable from that of equilibrium. For, "given a sufficiently long exposure, the visual system reaches a *state of equilibrium* with respect to any stimulus which may be acting upon it." In this state the eye yields a sensation which is not subject to further minuthesis.

Cobb (4) reports on the effect on vision of brightness of surroundings. He finds that brightness discrimination, and to a less extent visual acuity, are reduced when the surroundings have a considerably higher brightness than that of the test-field. Bright surroundings, *if not brighter* than the object to be looked at, are probably better for the seeing of detail than are dark surroundings. Brightness discrimination is best when the surrounding field and the test-field are of the same brightness. Cobb (5) also describes an improved apparatus for the study of this and similar problems. Nutting (28, of which 29 is an abstract) gives compact data on: (1) threshold brightness for different degrees of brightness adaptation; (2) "the brightness that would just appear uncomfortably bright [glare] with the retina adapted to any given brightness"; (3) the limen of brightness difference for different degrees of brightness adaptation (*cf.* Cobb, 4). A new term, "discrimination factor," "is defined as the field brightness [general brightness adaptation] divided by the just noticeable difference, B/dB . This

quantity is a direct measure of the power to distinguish details except when large color differences are present. Visual acuity so called, a mere sharpness of definition, is a minor factor when contrasts are slight." A brief but comprehensive table of measurements of this factor (for field brightnesses varying from 0.0000001 to 10,000.0 millilamberts) is given.

Ferree and Rand (12) add a fifth contribution to their series dealing with conditions of lighting. A considerable number of tables is given, which deal mainly with the properties of pendent reflectors, and will be of interest chiefly to those who are studying the question of illumination. The authors have summarized their general results in (13) and (14) where, however, little is added to the substance of papers which were reviewed in the *BULLETIN* in 1916. The authors find that it is highly desirable to have the field of vision uniformly illuminated (*cf.* Cobb, 4, and Nutting, 28), the light well diffused, and no extremes of surface brightness. Caldwell (3) finds "for a given observer and a given value of illumination a rather definite preferred relation between direct and indirect illumination." This preference is for rather "more indirect than direct illumination." This is in general agreement with findings of Ferree and Rand. Also "a notable capacity for retaining a concept of an illumination value and of reconstructing it after an interval of time, is demonstrated."

Of papers dealing more especially with methods and apparatus, one of the most generally valuable is by Cobb (6). The paper is written for "investigators in those branches of biological science which deal with light-effects," and describes clearly some of the fundamental physical conceptions which must be understood by any one who wishes to manipulate light-stimuli intelligently. Some of the heads treated are: Point and extended sources, regular diffuse and mixed reflection, the three analogous types of transmission, reflectors and transmitters as secondary light-sources, images as visual objects, the measurement of brightness, and the relation of light to energy.

Ferree and Rand (10) describe an ingenious form of campimeter in which the stimulus is spectral light of known wave-length and intensity, and can be presented at any point (up to about 90 degrees of eccentricity) of any meridian of the retina. The same authors (11) suggest, as a substitute for the artificial pupil, focusing the stimulus light upon the pupil of the eye, there "forming an image of the analyzing slit of the spectroscope." The essential features

of this method have been previously described by Cobb (*Amer. J. of Physiol.*, 1911, 29). Ives (18) describes an improved form of his visual acuity testing device (first described in 1910). The two single-line gratings of the earlier form are now replaced by cross-line gratings, which when rotated with respect to each other present a pattern of uniformly distributed squares which expand or contract as the gratings are turned. Kirschmann (24) recommends gelatine light-filters for the production of "monochromatic" light. The main advantage is that large areas can be illuminated. No special formulae for monochromatic gelatine dyes (the crux of the whole matter) are mentioned. Woodworth (43) gives an equation for calculating the conjugate foci of spherical lenses. The equation that is now commonly given in text-books is approximately accurate only for very thin lenses.

REFERENCES

1. ALMACK, N. & ARPS, G. F. On Color Induction with Reference to Color Recognition. *J. of Exp. Psychol.*, 1916, 1, 426-453.
2. ALSPACH, E. M. Simplicity vs. Complexity of Color Hues. *Amer. J. of Psychol.*, 1916, 27, 273-282.
3. CALDWELL, F. C. On the Effect of Brightness of Light Source on General Illumination. *Trans. of the Illum. Eng. Soc.*, 1916, 11, 1042-1049.
4. COBB, P. W. Vision and the Brightness of Surroundings. *Trans. of the Illum. Eng. Soc.*, 1916, 11, 372-398.
5. COBB, P. W. The Effect on Foveal Vision of Bright Surroundings—III. *J. of Exp. Psychol.*, 1916, 1, 419-425.
6. COBB, P. W. Photometric Considerations pertaining to Visual Stimuli. *Psychol. Rev.*, 1916, 23, 71-88.
7. COOK, H. D. & KUNKEL, F. M. The Qualitative Relation between Complementary and Contrast Colors. *Psychol. Monog.*, 1916, 22, 1-39.
8. CRITTENDEN, E. C. & RICHTMYER, F. K. An "Average Eye" for Heterochromatic Photometry, and a Comparison of a Flicker and an Equality-of-Brightness Photometer. *Trans. of the Illum. Eng. Soc.*, 1916, 11, 331-366.
9. FERREE, C. E. & RAND, G. A New Method of Heterochromatic Photometry. *J. of Exp. Psychol.*, 1916, 1, 1-12.
10. FERREE, C. E. & RAND, G. A Spectroscopic Apparatus for the Investigation of the Color Sensitivity of the Retina, Central and Peripheral. *J. of Exp. Psychol.*, 1916, 1, 247-283.
11. FERREE, C. E. & RAND, G. A Substitute for an Artificial Pupil. *Psychol. Rev.*, 1916, 23, 380-382.
12. FERREE, C. E. & RAND, G. Some Experiments on the Eye with Pendant Reflectors of Different Densities. *Trans. of the Illum. Eng. Soc.*, 1916, 11, 1111-1136.
13. FERREE, C. E. & RAND, G. A Résumé of Experiments on the Effect of Different Conditions of Lighting on the Eye. *Ann. of Ophth.*, 1916.
14. FERREE, C. E. & RAND, G. Miscellaneous Experiments on the Efficiency of the Eye Under Different Conditions of Lighting. *Ophthalmology*, 1916, 12.

15. GOLDSCHMIDT, R. H. Die Frage nach dem Wesen des Eigenlichtes, ein Hauptproblem der psychologischen Optik. *Psychol. Stud.*, 1916, 10, 101-155.
16. HOUSTOUN, R. A. A Theory of Colour Vision. *Proc. of the Roy. Soc.*, 1916, A, 92, 424-432.
17. IVES, H. E. & KINGSBURY, E. F. The Theory of the Flicker Photometer.—II, Unsymmetrical Conditions. *Phil. Mag.*, 1916, 31, 290-321.
18. IVES, H. E. An Improved Visual Acuity Test Object. *J. of the Franklin Inst.*, 1916, 182, 539.
19. IVES, H. E. Visual Diffusivity. *J. of the Franklin Inst.*, 1916, 182, 540-541.
20. IVES, H. E. A Polarization Flicker Photometer. *J. of the Franklin Inst.*, 1916, 182, 541-542.
21. IVES, H. E. Measurements of Brightness-Difference Perception and Hue-Difference Perception by Steady and Intermittent Vision. *J. of the Franklin Inst.*, 1916, 182, 542.
22. IVES, H. E. Hue Difference and Flicker Photometer Speed. *J. of the Franklin Inst.*, 1916, 182, 812.
23. JOHNSON, H. M. A Note on Ferree and Rand's Method of Photometry. *Psychol. Rev.*, 1916, 23, 390-396.
24. KIRSCHMANN, A. Ueber die Herstellung monochromatischen Lichtes in grösseren Flächen. *Psychol. Stud.*, 1916, 10, 185-188.
25. KRIES, J. VON. Messende Versuche über die Funktionsstellung im Sehorgan. *Zsch. f. Sinnesphysiol.*, 1916, 49, 297-315.
26. LADD-FRANKLIN, C. On Color Theories and Chromatic Sensations. *Psychol. Rev.*, 1916, 23, 238-249.
27. LANGFELD, H. S. Concerning the Image. *Psychol. Rev.*, 1916, 23, 180-189.
28. NUTTING, P. G. Effects of Brightness and Contrast in Vision. *Trans. of the Illum. Eng. Soc.*, 1916, 11, 939-946.
29. NUTTING, P. G. The Effects of Brightness on Vision. *J. of the Franklin Inst.*, 1916, 182, 531-532.
30. SWINDLE, P. F. Positive After-Images of Long Duration. *Amer. J. of Psychol.*, 1916, 27, 324-334.
31. TITCHENER, E. B. A Note on the Sensory Character of Black. *J. of Phil., Psychol., &c.*, 1916, 13, 113-121.
32. TITCHENER, E. B. A Further Word on Black. *J. of Phil., Psychol., &c.*, 1916, 13, 649-655.
33. TROLAND, L. T. Apparent Brightness; Its Conditions and Properties. *Trans. of the Illum. Eng. Soc.*, 1916, 11, 947-966.
34. TROLAND, L. T. The Laws of Visual Minuthesis: the Threshold Preëxposure Time and the Equilibrium Time for a Projected Negative After-Image. *J. of the Franklin Inst.*, 1916, 181, 579-581.
35. TROLAND, L. T. Notes on Flicker Photometry: Flicker-Photometer Frequency as a Function of the Color of the Standard, and of the Measured Light. *J. of the Franklin Inst.*, 1916, 181, 853-855.
36. TROLAND, L. T. The Laws of Visual Minuthesis: the Influence of Intensity on the Equality Time-Function. *J. of the Franklin Inst.*, 1916, 181, 855-856.
37. TROLAND, L. T. The Absence of the Purkinje Phenomenon on the Fovea. *J. of the Franklin Inst.*, 1916, 182, 111-112.
38. TROLAND, L. T. The Heterochromatic Brightness Discrimination Threshold. *J. of the Franklin Inst.*, 1916, 182, 113-115.

39. TROLAND, L. T. Notes on Flicker Photometry: Flicker-Photometer Frequency as a Function of Light Intensity. *J. of the Franklin Inst.*, 1916, 182, 261-263.
40. TROLAND, L. T. The Revival of a Faded Negative After-Image by Brightening the Stimulus Field. *J. of the Franklin Inst.*, 1916, 182, 529-530.
41. WARD, J. A Further Note on the Sensory Character of Black. *Brit. J. of Psychol.*, 1915, 8, 212-221.
42. WEISS, A. P. Purkinje Demonstration. *Psychol. Bull.*, 1916, 13, 442-444.
43. WOODWORTH, C. W. A New Fundamental Equation in Optics. *Science*, 1916, 43, 824-825.

CUTANEOUS AND KINÆSTHETIC SENSES

BY JOHN TRUMBULL METCALF

Smith College

Although the war has greatly reduced the number of articles on cutaneous and kinæsthetic senses, still the literature of the past year in some ways makes up in intensive interest what it lacks in extensity. Most important is the continued interest in the line of work started by Head and his collaborators. The study of the cutaneous sensibility of skin areas supplied by regenerating nerves was continued by Trotter and Davies, and with somewhat different technique by Hacker whose results were reported last year.¹ During the past year Boring (2) has published the results of his experiment on nerve-division. He has added to the results a critical review of Head's theory and a theoretical proposal of his own. Moreover, Carr (3) has made a painstaking examination of Head's results in connection with a critique of the Head theory. Boring's work was undertaken under rigidly controlled psychological conditions, the author himself acting as subject. The investigation therefore had the advantage of having a trained psychologist as observer. Quantitative methods were employed to test the sensitivity of the affected area. There was also a preliminary period of practice. This covered fifteen months before the operation during which time the subject devoted from three to eight hours a week to practice in discrimination of different cutaneous sense qualities, localization, and discrimination of two points. The nerve chosen for section was the anterior branch of the internal cutaneous of the left arm. As a result of the operation a small area of skin on the left forearm became anæsthetic. This area was located near the wrist on the volar surface and slightly to the

¹ See this BULLETIN, 1916, 13, 138.