AN ANTISPASMODIC

That is Safe—Non-narcotic

Benzyl Benzoate Miscible, H. W. & D.

Indicated and Effective

In Spasms of the Smooth Muscles · Colics

Especially In

TRUE ASTHMA—BRONCHIAL SPASM

And for Relief of

DYSMENORRHEA

See "The Journal" A.M.A., August 23rd, 1919

and Southern Medical Journal, July, 1919

Circulars Upon Request

HYNSON, WESTCOTT & DUNNING

BALTIMORE

The Natural Coagulant of Blood

Thromboplastin Solution (Armour) is a specific hemostatic and is made from the brain substance of Kosher killed cattle. This brain tissue of cattle killed according to Mosaic law is uninjured and by the Armour process this "principle" which causes coagulation is extracted and supplied to the medical profession in standardized and sterilized form.

Thromboplastin Solution (Armour), is useful in the treatment of hemorrhage, especially that from oozing surface, scar tissue and the nose and throat.

25 c. c. vials, in dated packages.

Pituitary Liquid (Armour) is the most trustworthy solution of the Posterior Pituitary Substance. It is free from preservatives and is standardized physiologically by the Roth Method. ½ c. c. and 1 c. c. ampoules.

Thyroids (Armour) runs uniformly 0.2 per cent organic iodine in Thyroid combination. Thyroid Tablets (Armour) ¼, ½, 1 and 2 grain. When Thyroids is indicated specify Armour's.

We offer all the endocrine gland preparations in powder and tablets. All drying of the glands is done in vacuum ovens at a low temperature. This insures uninjured therapeutic value.

Circulars on request

ARMOUR & COMPANY
CHICAGO

SOUTHERN MEDICAL JOURNAL

JOURNAL OF THE SOUTHERN MEDICAL ASSOCIATION

PUBLISHED MONTHLY AT BIRMINGHAM, ALABAMA

Volume XIII

FEBRUARY 1920

Number 2

MEDICINE

INTERNAL DISEASES, PEDIATRICS, NEUROLOGY, DIAGNOSTIC METHODS, ETC.

FOCAL INFECTION*

By THOMAS D. COLEMAN, A.M., M.D.,
Professor of Medicine, Medical College of
Georgia,
Augusta, Ga.

The title of this paper like "De Profundis" is selected not from any notion on my part that I can compass all its problems, but because many of them invite elucidation and discussion by the laboratory worker and the clinician. Infection may be defined as the invasion of the living body by pathogenic micro-organisms, which by their life processes produce perversion of function. Focal infection as generally understood and accepted, means the invasion, and in consequence, perversion of function of some organ or organs, more or less removed from the focus of infection. The phraseology is catchy, because it invites one to reason from cause to effect, but is not altogether satisfying, since etymologically, the word focus means literally a point, or sharply circumscribed area, whereas the focus in some cases of infection may extend over wide areas, e.g., the bronchial tree, the intestinal canal, genito-urinary tract, etc.

Less objectionable would seem to be the term autoinfection, meaning thereby an infection of the living body by pathogenic organisms within itself; and yet in narrow limits, similar objections might be urged. Focal infection is not a new conception. In the first century it found expression in these words: "If thine eye offend thee pluck it out and cast it from

thee." The principle has long been recognized, but in relatively recent years we are indebted to medical workers for calling attention to the fact of focal infection and its role in the production of disease. It is not the purpose of this paper to mention the tissues that are liable to focal infection, for this would involve a description of the entire human anatomy, nor to describe the micro-organisms concerned in such processes, for it would entail a dissertation on bacteriology; but I would feel myself derelict to my medical ideals if I did not call attention to certain extremes, not to say errors, that inevitably follow the exposition of facts presented in a new light. Discoveries of a serviceable nature to mankind when freshly made, or reproduced and presented plausibly, are apt to lead to fads, and fads lead to extremes of thought and action.

It is as natural for medical men to follow fads as it is for the human race to follow any "will o' the whisp" that promises relief from the ills that flesh is heir to. To speak not at random, fads often lead the honest man unconsciously into error. They are the food and drink of the shallow-pated and the charlatan. They are utilized by the wise. All fads have a medium of merit to account for their existence. Men of science strive to winnow out the wheat from the chaff. The evolution of an idea like the evolution of a species, is apt at times to produce sports and monstrosities, so one need not expect that medicine can escape these tendencies. The main thing is not to ignore but to profit by them.

A notion of focal infection is as old as the human race. The role which bacteria play in the tragedy was unappreciated

*Read in Section on Medicine, Southern Medical Association, Thirteenth Annual Meeting, Asheville, N. C., Nov. 10-13, 1919.

until the days of Pasteur and Lister. A retrospection at this time and in this connection, will, I am sure, be well worth the moments' consideration. It is within comparatively recent years that the ovary attracted the needlessly assiduous and sometimes rude attention of the surgeon. It was regarded as being the source of a host of woman's ills, complaints or idiosyncracies. That it is the cause of many of their ills cannot be disputed. Diseases real and imaginary were attributed to the always fecund, but sometimes innocent ovary. It became a matter of distinction to have had one's ovaries removed, and discussion of it even with one of the opposite sex was not taboo. In this, if I may be pardoned the diversion, the bar sinister of the sex is established; one could not even in imagination picture a bunch of men glorying over their own enucleation; but such is the nature of women. In the last public utterance which I was privileged to hear from the late Abraham Jacobi, he said to a body of my students: "I was recently in the office of a surgeon and he showed me with great pride a jar filled with ovaries that he had removed. Instead of praising him as he had expected, I told him I knew a better place for ovaries than glass jars." This sums up what it took many years to learn. It is not now regarded as the mark of a good surgeon to cut out an ovary, but rather to cut out only that which is diseased and leave the rest of the organ for the normal uses that nature intended. Another organ to go under the ban in the fight against focal infection was the appendix vermiformis, and so violent was this onslaught that it was said jestingly at one time that people in the larger centers were having stamped on their under clothing "mine has been removed," so that if they were found unconscious they might not be operated on a second or third time. "An inch and a half incision and a week and a half in bed" has passed into a proverb; it was enunciated first, I believe, by one of the great geniuses of modern surgery, to whom we are all debtors, and yet one can but be impressed by the relative infrequency of appendix operations now compared with a decade ago. Except in fulminating cases Oxner's dictum paraphrased; of watchful waiting, I believe, is now adhered to, although some surgeons still stand for immediate removal.

Then followed the fad of removing tonsils and adenoids. Many of the generation of men had had tonsillitis, and lived to tell the tale, but it devolved upon the focal fad followers, if I may so style them, to propagate the dictum that all enlarged tonsils must be immediately removed root and branch; and with this idea very radical operations are being performed even unto this day. It was held that the tonsils were not only useless, but were the atria of entrance into the body for germs, that many, if not most, of the bacteria that lurked in the air found entrance into the body through the tonsils; and that those which they could not admit immediately into the blood stream they held in captivity until some convenient time when they could do so. As a result large numbers of full grown tonsils, many adolescent ones, some even infantile ones, were removed. Some patients were benefited; disappointment was the lot of others. In the hands of conservative men it is, I believe, the practice now, to remove only such tonsils as are manifestly diseased, or which hold in their structure infectious organisms.

The present fad extends several years back, and expresses itself in war on the teeth—shall I say to the teeth? And so teeth are coming out of the mouths of people like corn out of a hopper, and dentists are everywhere apparently unequal to the demands upon them. I fear the fad is making a generation of toothless young people, and one horror of old age is anticipated. Artificial teeth are not the equal of those furnished by nature, and I am persuaded that many of the latter are being unwisely sacrificed, which could be preserved by appropriate treatment *in situ*. It is not the work of a good dentist to pull teeth, but to save them and I believe that conservation will yet assert itself; the pendulum will swing back and only teeth will be removed which cannot be saved or which are causing demonstrated systemic infection.

Even the colon with its expanse of surface has not escaped attention, and colons have been removed in whole or in part, with doubtful benefit, and sometimes grievous harm to their possessors, in the hope of relieving focal infection—intestinal stasis, etc. As previously stated, these extremes of practice should not lead to the denial of the theory of focal infection, for

in the light of present knowledge it would make one ridiculous; nor should one too readily condemn wholesale those who in their enthusiasm advise and employ operative procedures of such nature. One should be impressed, however, with the serious necessity of careful study of all the premises before deciding that a given organ is the source of systemic disturbance and should be removed. Such organs as the ovaries, the appendix, the tonsils and the teeth are not placed in the body for purely cosmetic reasons.

According to Blum, in an article on "Focal Infection in Childhood":

"The mere presence of pathogenic micro-organisms within a circumscribed area does not *per se*, constitute an infective focus. At most this condition may be regarded as a potential focus of infection. Bacteria, for example diphtheria bacilli, may be harbored indefinitely in a given locality and be perfectly innocuous. They may be non-virulent or walled off, or the patient may be immune to their specific action or conditions necessary for their metastatic propagation may be lacking.

"Failure to recognize this fact has resulted in much misdirected treatment, especially enucleation of the tonsils and extracting of the teeth. For the creation of infections from existing foci two additional factors are indispensable—the bacteria must invade the circulation (hemic or lymphatic) and the part to be infected must be ready for their reception.

"Eradication of the primary focus of infection is certainly to be desired but, as has been stated, it is not always feasible. In toxemia due to a local abscess, putrefaction in the intestines, etc., cure of the focus may alone be sufficient treatment. But to assume that in cases in which a distinct metastatic bacterial process has been established, as in endocarditis or arthritis, eradication of the primary focus will cure the secondary lesion is fallacious. Such processes, once established, are selfsupporting, the most that can be expected from removal of the primary focus under such conditions is that the possibility of further deleterious increments from the original source would be excluded. The question whether treatment should be aimed at the cure of the infection or removal of the infected structure must be decided on its merits in each individual case."

How often has it occurred within the experience of medical practice that the ovaries when removed, as they were e.g., in Battey's operation, did not improve the nervous phenomena from which the patient suffered; that the extirpated appendix showed no evidence of infection, gross or microscopic; that removal of the tonsils and the teeth had no effect whatever on the arthritis, myositis or pancarditis from

which the patient may have been suffering and that in the language of the scriptures, "Verily the last end of that man was worse than the first."

How often does one not find that patients with badly diseased tonsils, alveolar abscesses, and sinus disease, present no evidence whatever of general or systemic infection.

It goes without saying that where an organ is diseased to an extent that it is a menace to the body, if it does not entail too great danger to other and more important tissues, it ought to be eradicated. "Ubi pus ibi evacua" may not be classical Latin, but it is an axiom, sound and true, but it also should be borne in mind, that often even pathogenic organisms remain strictly localized, and produce no systemic effect; and that, in the process of removing these foci, openings into the blood stream or lymphatics may be made which lead to general infection, the very thing which one wishes to shun. Numbers of such cases have been reported. Unstinted praise should be given to the pioneers in this field who have so enriched our knowledge of focal infection; to such workers as Winge, Rosenow, Billings, Pointon, Paine, and a host of others for giving a rational basis for systemic infection resulting from focal lesions, our debt is great; but the value of their work and the allurements of their theories may, when pushed too far, lead to excesses in treatment. The fact that a patient has a circumscribed abscess at the root of one tooth does not keep him from having a cholecystitis, or open pulmonary abscess that may be accounting for his systemic symptoms; under such circumstances all the teeth might be extracted and the patient be not a whit relieved. And while it is true that many acute and chronic infections which may be general, may and often do, start from infections in the mouth, nose, ear, sinuses, and genito-urinary tract, it is also true that the alimentary canal is a source of infection with its twenty feet or more of gut; much more frequently than is commonly appreciated.

In making a diagnosis of focal infection, and by this I mean systemic infection from some focus or foci, it is needful to employ every means of diagnosis—clinical, biochemical, bacteriological and radio-graphic. Even when all these are exhausted, one may fail.

It is not only necessary to determine that the systemic infection is due to an infective focus and secondly to locate that focus; but often a focus which seems potentially capable of spreading systemic infection is innocent or innocuous, and some focus far removed and unsuspected is producing the trouble.

Bodily or systemic infection is effected through the medium of the blood and lymphatics, micro-organisms attack with frequency endothelial membranes which seem to possess small resistance to invading bacteria. This is shown by the frequency of involvement of the endocardium, pericardium, synovial covering of joints, peritoneum, pleura, meninges, and the mucous membranes of the mouth, and accessory passages, the alimentary canal, and the genito-urinary tract. Sometimes it may be by extension, simply as one sees in erysipelas, and yet toxins are set free which produce systemic disturbance. It is of moment, therefore, if one is to combat effectually the process that not only the primary but also the secondary foci shall be discovered.

When a focus of infection is established, it may remain innocuous to the body in general, e.g., in an encysted tuberculous nodule or a furuncle in the subcutaneous tissues. A few organisms may at times escape the barriers, but they are destroyed by the phagocytes or the inimical plasma. When they escape in larger numbers or their virulence is intensified, or the defensive processes of the tissues is impaired, general infection ensues.

A case from a distant city referred to me by a colleague last January will illustrate, as well as many, the problems involved, both in the manner of the spread and the difficulties involved in the treatment of focal infection.

Mrs. C., a robust woman of middle age, had noticed a number of months previously a glandular swelling in her left axilla. After long persuasion she consented to have it removed. It proved to be a tuberculous gland. Following this a tuberculous process was discovered at the apex of the right lung. This in turn was followed by pain in the region of the right kidney. The urine was examined and found to contain tubercle bacilli; the right ureter was catheterized and tubercle bacilli found. The involved kidney was removed. A cystoscopic examination showed tuberculous ulceration of the bladder. Then the right hip became involved. Her weight and general condition remained good, and she returned

to her physician in apparently as good condition as she came with the hip condition improved, the lung condition stationary, the bladder condition apparently unchanged, and still complaining of pain in her back. I saw nothing to be gained by putting her through the ordeal of catheterizing the left ureter, since if the left kidney was involved treatment other than general and symptomatic could be instituted. She could not possibly part with the remaining kidney.

The lessons to be learned from this case were, (1) that it was reasonably certain that the gland in the axilla was not the primary focus. More than likely it was a bronchial gland.

(2) That the removal of a single focus of infection does not always eradicate the primary focus.

(3) That there may exist in the body foci of infection that may not be removed, in this case in the lung, hip-joint, bladder and possibly the remaining kidney.

In the treatment of cases of focal infection it is essential that (1) the focus or foci should be removed where the end justifies the means, and one should keep in mind the possibility not only of not doing good, but also of doing harm "Non Nocere."

(2) Every agency both in building up the bodily defenses and ameliorating symptoms as they arise should be employed.

To all those who lean to radicalism, I would suggest a motto, "Festinate lente."

CONCLUSIONS

(1) The fact of focal infection is not disputed.

(2) Focal infection is caused by pyogenic organisms that are distributed through the body by means of the blood and lymph streams.

(3) Foci of infection when removable should be removed, but removal should be a treatment and not a diagnostic procedure.

DISCUSSION

Dr. Douglas VanderHoof, Richmond, Va.—Medicine of the future is unquestionably preventive medicine, and I feel that the study of our patients should invariably include a search for possible foci of infection which we have reason to believe may subsequently become a pronounced menace to their health.

Dr. Coleman's paper draws attention to the fact that ultra-radicalism is to be condemned, and in that I am sure we all agree; at the same time every one of us is daily confronted in our consulting rooms by horrible examples of neglected chronic infections. The chief advantage of a regular systematic oral examination as has

been instituted by certain industrial companies and certain insurance organizations, is to be found in the fact that such an examination not infrequently discloses apparently simple foci of infection which, if not removed, may ultimately prove a great menace to the health.

Why is it, however, that certain individuals with infected tonsils, or infected teeth, suffer from arthritis, whereas other individuals with the same amount of focal infection are apparently immune? That question is so frequently asked me by my students. We do not know. Rosenow explains it on the basis of a chemical tropism; an affinity, if you wish to call it, of the joint tissues of certain individuals to the toxins or the organisms originating in the focus of infection. In other words, Dr. Coleman and I each might have equally infected tonsils; he may be in apparent perfect health, while I may be horribly crippled with arthritis. The tissues of my joints possess apparently some attraction or some affinity for the toxins in the tonsils, whereas Dr. Coleman, being more fortunate, has joint tissues which do not have this tropism or chemical attraction.

In regard to the failures, and there are many, that result from the removal of chronic foci of infection, we must admit that we are not yet wise enough, when confronted with a constitutional condition associated with certain focal infections, to be able to establish a direct relationship between the two. In our ignorance, it seems to me, we should proceed carefully but definitely to remove obvious foci of infection in an attempt to restore the patient to health. Our failures, it seems to me, are due to one of two things: our diagnosis is at fault or else we have operated too late. In the latter instance metastatic foci of infection have occurred and the patient is not relieved by the operation to remove the primary focus. We know when a woman finds a small lump in the breast and consults a surgeon and has the breast removed, she has a certain chance of getting absolutely well. There is almost an equally big chance that she will subsequently die of carcinomatosis. We believe, and we have every reason to believe, if that small lump in the breast is removed early enough we remove the original focus of malignant disease and the patient never has any recurrence. The analogy applies, I believe, in our failures in constitutional conditions dependent upon foci of infection. These foci are not removed until the infection has spread and has involved other tissues, possibly the joints, possibly the endocardium, and so on. Dr. Coleman touched upon that possibility in his paper. It is extremely important in the study of a patient to be always on the lookout for chronic focal infections which are possible menaces to the future health of that individual. If they are found it is my opinion that they should be removed.

The question of finding these foci of infection emphasizes the fact that we are very dependent upon our colleagues. My dentist, my throat man, my surgeon, my x-ray man, men who are associated with me in clinical studies, must be competent in their judgment and, above all, hon-

est in their opinions, and it is only by being associated with a group of men in whose work we have absolute faith that we can hope to succeed.

Dr. Bruce James, Danville, Va.—I would like to say that while this matter of focal infection is certainly very important, we are going too fast with it. About a year ago a woman came to my office with eczema all over her face, which was very intractable. She had consulted two specialists who looked for focal infection and after x-raying her mouth decided that her teeth had to come out. She was almost paralyzed at this advice, for she had beautiful teeth. She came to me in despair. Suspecting intestinal toxemia, I referred her to a man whom I thought would find out what was the matter with her. He examined her very carefully and found the trouble was in the intestines, and she escaped having her teeth taken out.

Since I have been here I recall the fact that a young woman was sent to this town (Asheville) for a continuous fever. It was thought she had tuberculosis, though we could not find any lesion in the lung. She was weak, had this continuous fever and loss of appetite. The physicians here thought she had a tubercular infection but could not find it, and finally they decided to have six of her teeth extracted. This was done, but following the extraction of these beautiful teeth she was in the same condition as before.

We are getting too faddish about these things, especially the tooth proposition. I do not think taking out the tonsils does any harm or even taking out the appendix or gall-bladder, but when we advocate and take out the teeth it is a different proposition, and we ought to be very conservative about the teeth.

Dr. W. L. Dunn, Asheville, N. C.—I am sure that Dr. Coleman, like the rest of us, has seen remarkable things happen after the removal of foci of infection. He evidently has seen the same thing as the rest of us, a veritable mania on the subject of removal of all sorts of foci that might be infective, and, therefore, I think his paper is very timely.

Those who have had numbers of cases referred to them do not realize to what a tremendous extent the teeth are being sacrificed today by men everywhere and without very careful consideration. Dr. VanderHoff touched a very important point when he said we are dependent upon those who are associated with us and must be certain of their ability as well as their honesty. This is not a reflection upon men in special fields, but the men who are familiar with our work in our own particular field seem to lose their sense of proportion; they lose their perspective.

As to the removal of the teeth, we know patients are having duplicate films made to carry around and keep them as special souvenirs showing what teeth were taken out. It is not at all infrequent to have a patient tell us that she has false teeth and in her pocket a set of films.

I have made it a point to study these cases carefully, and I am perfectly certain that a large number of teeth have been sacrificed when there was absolutely no excuse for it. We have not reached that degree of perfection in our

x-ray diagnosis where we can say with definiteness that a tooth has slightly erupted, or that the periodontal membrane around the tooth is inflamed and the tooth should therefore be taken out. There are so many chances for error in overexposure or underexposure of the teeth. We must be certain of our interpretation.

I quite agree with Dr. VanderHoof's position that we are in the day of preventive medicine and should look out for the things that may happen to our patients as well as the things that have already happened. We must not swing in the other direction.

I am reminded of a patient who consulted a physician who referred the patient to a hospital and she had six beautiful teeth pulled out.

Another patient who had for more than two years tuberculosis, or which was said to be tuberculosis, without any marked physical evidences so far as the lung was concerned, but because of some minimal x-ray findings and the fact that she ran a temperature and had some other symptoms which ordinarily go with tuberculosis, she was thought to have tuberculosis. Examination showed that this patient had had her teeth operated upon by a dentist, and through some slip he let a piece of his broach slide into the antrum, and that had been resting there peacefully for years. On opening the antrum and removing the piece of dental instrument, and likewise treating the antrum, the patient was cured of what was thought to be an intractable tuberculosis.

*Dr. Coleman (closing).—*The main object of my paper was to try to preach a little conservatism because I felt we were going too far in the opposite direction.

I have followed many cases to the operating table, not operating myself, and I would not like to say how many healthy ovaries I have seen removed, or how many perfectly normal appendices I have seen excised, how many teeth I have seen extracted without the relief of systemic infection, etc. The burden of my paper, and I perhaps repeated it enough to be a little redundant, was that where a focus is found that is actually causing systemic disturbance it should be removed, but in the removal of foci be sure you are doing it as an operative procedure and not as a diagnostic one.

SUBACUTE COMBINED DEGENERATION OF THE SPINAL CORD*

BY WILLIAM G. SOMERVILLE, A.B., M.D.,
Memphis, Tenn.

It is not the intention in this essay to refer to Friedreich's ataxia, nor to those forms of combined sclerosis which are sometimes caused by luetic infections. The title of the essay is one used by English authors for a symptom complex consisting

of anemia and a combined degeneration of the posterior and lateral columns of the cord, running a fairly acute or subacute course. According to the pathology of the disease, it would seem that combined degeneration is a more suitable term than combined sclerosis, as used and first described by Putnam and Dana, of this country.

It occurs as a rule during the fifth and sixth decade of life. Secondary anemia is present at some stage of the disease and the nervous symptoms may precede the anemia or vice versa.

The pathology consists of lesions in the posterior and lateral columns of the cord and frequently throughout the brain, as shown by Lurie.* These lesions are foci of degeneration, apparently arising from the blood vessels.

The secondary anemia, which is present at one time or another during the course of the disease, may or may not be pernicious; this is not a cause of the degenerative changes found in the central nervous system, but both are caused by some toxic agent. When fully developed the paraplegia may assume one of two forms, according to whether the lesions predominate in the posterior or lateral columns of the cord. The one closely resembles tabes, with its ataxia, hypotonus, and loss of knee jerks; the other is an ataxic spastic type with increased knee jerks and Babinski's sign; though this latter type usually, sooner or later, changes to the flaccid type, as the foci of degeneration in the posterior columns extend downward in the lumbar segments of the cord. A very important and striking point in diagnosis is that even in the tabetic form we find Babinski's sign.

The early nervous symptoms are mainly subjective, and by no means assume such clear clinical types as just mentioned, but begin as "stocking glove" tingling and numbness; also a tight feeling a sensation as if the feet or legs were tightly bandaged. These subjective symptoms are persistent and characteristic; but when one looks for objective signs of sensory disturbance, they are very slight in the early stage; there being usually no diminution of touch and very slight loss of temperature and pain sensation. These paræsthesias and beginning sensory changes make one think of

*Read in Section on Medicine, Southern Medical Association, Thirteenth Annual Meeting, Asheville, N. C., Nov. 10-13, 1919.

multiple neuritis, and may lead to error in diagnosis.

The lower extremities are affected earlier and to a greater extent than the upper extremities; the paræsthesias gradually extending up the limbs on to the trunk, frequently with the sensation as if there were a tight band around the body. This may occur before there are any marked objective signs of sensory loss. One of the earliest sensations lost is the vibratory, the next is joint and muscle; then follow pain and temperature; while tactile sense is frequently the last to go; so that we may even find at some stage a dissociated anesthesia. Batten and Collier call attention to a "heavy fixed pain in the hepatic regions," of which complaint is frequently made. Throughout the course of the disease there occur at intervals, gastro-intestinal attacks, vomiting and diarrhœa with irregular temperature, and coinciding with these attacks there is an exacerbation in the nervous symptoms. Then there develops the *pæraplegia*, either the spastic or flaccid type; though the former usually changes to the latter, but with both types there is nearly always present Babinski's sign.

Mental symptoms, varying in intensity, may develop later in the disease indifference, dullness, mental depression or excitement, or a toxic exhaustive psychosis. These mental symptoms with the brain pathology have been fully discussed by Lurie.*

The case of Mrs. H. P. C., recently seen by me, may be regarded as a fairly typical one. She is 55 years of age, and her trouble began insidiously fifteen months ago, in the spring of 1918, with numbness and tingling in the feet and legs which has gradually extended up above the umbilicus. Recently there have been numbness and tingling of the hands to the wrists. For several months past there have been weakness of the lower extremities and a rather marked ataxia. There is some difficulty in voiding urine, but no retention nor incontinence. She complains of a sensation in the lower extremities as if they were tightly bound with bandages. She has been in bed, at intervals, since June, 1918, with periods of improvement, and there have been gastro-intestinal attacks with some rise of temperature at intervals of every few weeks. Her color is pale and sallow; the heart and lungs are negative, she is poorly nourished; hemoglobin is 63 %, R. B. C. 2,300,000; there was slight variation in size, but no nucleated red blood cells were found.

The neurological examination showed pupils

normal; no diplopia nor nystagmus; optic discs normal; no cranial nerve paralyses. There were marked tingling and numbness of hands and of lower extremities as high as waist line; limbs felt as if bound by bandages. She had a sensation as if belts were around the body at three levels, viz: the bladder, umbilicus, and ensiform cartilage. This latter had been noticed only during the previous week, following a gastro intestinal attack.

The sensation of touch was slightly if at all diminished; pain and temperature were diminished, but not lost; sense of position was present, but vibratory sense was completely lost as high as the wrists and throughout the lower extremities to and including the crest of ilium. There was muscular weakness of the lower extremities, more noticeable in the hip muscles. There was no atrophy; but both wrist, knee and ankle jerks were absent. Babinski, Chaddock, Oppenheim and Gordon's signs were present, both right and left. In every stool examined there was found the *Cercomonas intestinalis*.

Jelks, of Memphis, finds this same *parameba* in all cases of pellagra. We wish to draw attention to the similarity between the nervous symptoms of subacute combined degeneration of the spinal cord and pellagra. Both diseases have in common secondary anemia, gastro-intestinal disturbances, degeneration of the posterior and lateral columns of the cord; pigmentation of nerve cells and frequent occurrence of mental symptoms. One of the chief points of difference is, the presence in pellagra of the characteristic skin lesions, which, it is agreed, are due to the trophic disturbances, from involvement of the spinal ganglia—that is the extension of the foci of degeneration into the posterior nerve roots and their ganglia. The secondary anemia is common to both, but more intense in the subacute combined degeneration, and more likely to assume a pernicious type. The *cercomonas intestinalis hominis* has been found in the intestinal discharges of both—in every case of pellagra, according to Jelks, and in the case of subacute combined degeneration reported here.

It is very evident that one case proves nothing, but this statement is made as a preliminary to further investigation, and to call attention to the very great importance of examination of the intestinal discharges of these patients, as well as of a careful comparison of the clinical signs and pathological findings in these two types of disease. For, after all, we may find they are one and the same, and that all their symptoms are produced by a common cause, an intestinal parasite.

*Lurie, L. A.: Pernicious Anemia with Mental Symptoms. Archives of Neurology and Psychiatry, Vol. II, p. 67.