It seems certain that the early years of life are those in which tuberculous infection takes place in the great majority of cases, and primary infection of adults is much less frequent than formerly believed. Clinical histories cannot be relied on to determine the period of infection. It seems equally certain that infection of adults does occur, and that no age is exempt.

CONCLUSIONS

1. The evidence at hand indicates that in the majority of cases the respiratory tract is the route of infection in tuberculosis.
2. The alimentary tract is a frequent portal of entry for the tubercle bacillus.
3. The tubercle bacillus is able to pass through the intact mucous membrane of the alimentary tract without producing a lesion at the point of entrance. This takes place most readily during the digestion of fats.
4. The bacilli pass with the chyle through the lacteals and thoracic duct into the blood, which conveys them to the lungs, where they are retained largely by the filtering action of the tissues.
5. Infection through the alimentary tract is especially frequent in children.
6. Infancy and childhood are preeminently the periods of life when the individual is susceptible to tuberculous infection, and the majority of cases of infection occur during these early years.
7. Any campaign against tuberculosis which leaves out of consideration the protection of children against infection will fail of success.
8. Tuberculous infection in adult life occurs, but not so frequently or readily as generally believed.
9. Tuberculous infection may occur at any age.

TRANSFUSION OF BLOOD WITH SPECIAL REFERENCE TO THE USE OF ANTICOAGULANTS*

HENRY S. SATTERLEE, M.D.

AND

RANSOM S. HOOKER, M.D.

NEW YORK

We previously reported¹ a method of transfusing blood by means of specially constructed pipets and canulas designed for use with either a lining of hardened paraffin or a preliminary internal coating of an anticoagulant solution such as leech extract or herudin². During the past eighteen months we have experimented with various anticoagulants other than paraffin and herudin, with a view to determining the best anticoagulative agents for purposes of transfusion. We desire here to report the results of these experimental transfusions and to summarize the theoretical and practical advantages of the anticoagulants which we have studied; and in order to make more evident at the outset the practical bearing of this study, we propose, first, to discuss briefly the present status of blood transfusion as a rational therapeutic procedure, and to call attention to certain aspects of this problem which are not usually taken into consideration. It is not our

intention by this discussion to disparage the valuable and often brilliant results which have been and are being constantly achieved by blood transfusion, nor to argue for any special restriction of its field of usefulness, but we do wish to point out the need for a more exact knowledge of the biologic processes which are involved in transfusion of blood (or which may be involved under special conditions) in order that the certainty of its usefulness may be increased and that the liability to any untoward results may be minimized beyond what is realized at the present time.

In the constantly enlarging field in which transfusion of the blood is being employed, the questions which arise concerning the indications and contra-indications for this operation, the amount of new blood to be supplied, and the advisability of removing some of the patient's own blood at the time of transfusion, are all matters closely concerned with the probable benefit or futility of the operation. For instance, in the emergency of acute hemorrhage occurring in an otherwise healthy subject, or following the artificial removal of acutely poisoned blood, as in the treatment of gas poisoning, we believe that the problem is a comparatively simple one and that the established tests for hemolysis and agglutination between prospective donor and recipient are probably adequate, if indeed the urgency of the conditions may not make it imperative to proceed without taking time for even these tests. But the problem is far more complex when it is a question of the benefit to be derived from the transfusion of a so-called "normal" blood into the circulation of a patient suffering from one or more of the various pathologic conditions for which transfusion has been recommended, and we believe that this question could be more satisfactorily answered if laboratory methods were available for the detection of other possible grounds of incompatibility than those which are covered by the tests which are now in use.

Among the recent contributions to the literature of transfusion we can find only one reference³ to accidents which appear definitely to suggest inadequacy of the present tests for blood compatibility, although there are numerous citations of unexplained chills and febrile reactions.

Percy⁴, in a series of fifty-four transfusions, met with toxic reactions in three cases in which the hemolysis and agglutination tests were perfectly negative, and two of these cases resulted fatally.

Ottenberg and Kaliski⁵ estimated that there are toxic reactions not referable to hemolysis or agglutination, in 10 per cent. of all cases of transfusion, which estimate appears to be based on their own series of 128 cases.

Ottenberg and Libman,⁶ in their series of 212 transfusions on 189 cases collected from hospital and private sources and performed by a number of different surgeons, have contributed a valuable classified report. These authors, in summarizing the untoward results due to transfusion, assign the most unfavorable reactions of their series, including five fatalities, to "incompatible blood" (meaning, apparently, hemolysis or agglutination only) or to hypertransfusion. But they also state that a "more or less febrile reaction" occurred in about 10 per cent. of all cases and that skin eruptions, usually urticarial, occasionally petechial, appeared in another 10 per cent. In referring to the hemolytic and agglutinative reactions as being an insufficient guide to the value of a given donor's blood for the purpose of stimulating a remission of the disease in pernicious anemia, Ottenberg and Libman

* From the Laboratory of Clinical Pathology, Department of Medicine, College of Physicians and Surgeons, Columbia University, New York.

1. Satterlee and Hooker: Surg., Gynec. and Obst., 1914, xix, 235.

2. Satterlee, H. S., and Hooker, R. S.: The Use of Herudin in the Transfusion of Blood, THE JOURNAL A. M. A., June 6, 1914, p. 1781.

3. Percy: Surg., Gynec. and Obst., 1915, xxi, 360.

4. Ottenberg, Reuben, and Kaliski, David J.: Accidents in Transfusion, THE JOURNAL A. M. A., Dec. 13, 1913, p. 2138.

5. Ottenberg and Libman: Am. Jour. Med. Sc., 1915, xl, 36.

make the interesting observation that "there are probably other not yet recognized differences between the bloods of donors."

Lindeman⁶ reports 4 chills and 14 febrile reactions in 46 transfusions with related donors, and 22 chills and 32 febrile reactions in 62 transfusions with unrelated donors.

Lewisohn⁷ reports 5 febrile reactions and 3 chills in 22 transfusions.

Two cases of toxic reactions in which the hemolysis and agglutination tests were favorable have been reported to us by Dr. Robert A. Cooke of New York in a series of twelve transfusions.

Without going further into statistics it will be enough to say that those who have had experience with the transfusion of blood in varying pathologic conditions and by various methods have observed that certain toxic symptoms varying all the way from a slight chill and rise of temperature to marked anaphylactoid phenomena and even death have occasionally resulted as an immediate sequel to transfusion and where the usual serologic tests for hemolysis and agglutination have been entirely favorable.

An extended discussion of the available hypotheses by which to explain such phenomena would occupy too much space here, but we believe that certain grounds of explanation are more strongly suggestive than others and are therefore worthy of special attention. Recent researches on the blood, and especially on the action of serum ferments, have assumed a general significance as affecting many questions of immunity, and are most particularly pertinent to the problem which here concerns us. We will therefore consider three hypotheses, any or all of which may be applicable.

1. *It is possible that the trypsin-antitrypsin balance in the circulating blood of the recipient may be so disturbed by the commingling with the donor's blood as to result in the immediate formation of serotoxin from cleavage of serum protein.*

For instance, Jobling and Petersen have shown⁸ that homologous serum may be toxic for animals under conditions which remove the antiferment and permit the serum protease to act, that⁹ serotoxin (anaphylotoxin) formation is coincident with the adsorption of the serum antiferment and¹⁰ that an increase of antiferment diminishes susceptibility to trypsin shock. These investigators have further demonstrated¹¹ that acute anaphylactic shock is accompanied by the intravenous mobilization of a large amount of nonspecific protease, a decrease in antiferment, an increase in noncoagulable nitrogen of the serum and an increase in amino-acids, which they attribute to the cleavage of serum proteins through the peptone (toxic) stage by a nonspecific protease. They conclude that the specific elements of this reaction consist in the rapid mobilization of the protease and in colloidal serum changes which bring about a change in antiferment titer, and that these changes are probably due mainly to an adsorption or lessening of the dispersion of the unsaturated lipoids on which the antiferment property depends.

2. *It is possible that the action of protective colloids in the body cells of the recipient may be so disturbed that these cells are thereby exposed to a reaction of antigen and antibody present in the circulation of the recipient but harmless to the protected cell.*

In support of such an explanation, the work of Zinsser and Young¹² on the possible importance of colloidal protection in

certain phases of the precipitin reaction, has a distinct bearing.

If further investigation and better knowledge should prove the validity of these assumptions, it is most probable that laboratory tests can be evolved which will show in advance the liability to harmful results from such causes. Meanwhile it would be well to recognize that delicate physicochemical balances may be disturbed and toxic biologic reactions may take place as a result of transfusing blood under certain conditions, and that the usual tests for hemolysis and agglutination, although of primary importance, do not afford the complete assurance of freedom from unfavorable reaction which is commonly believed.

The increasing attention which is being given in the current medical literature to simplified methods of transfusion has tended to popularize the idea that almost any technic of operations may be employed which will serve to transfer blood under aseptic precautions and in a fluid state from the circulation of a donor to that of the recipient. As opposed to this trend of opinion, we believe that the removal of so complex a tissue as the circulating blood and its exposure to foreign physical influences during transfer to another human system should be regarded as at least as delicate an operation as the transplantation of any other living tissue, and should be given the same degree of consideration for favorable biologic conditions as any other surgical transplant. These conditions are not satisfied by providing only for the transfer of the blood in a germ-free and fluid state; they should also preclude the possibility of other changes tending to lessen its functional value as a transplant, and most particularly should the conditions preclude the possibility of harmful disturbances in the recipient's circulation induced by such changes. This brings us to another possible causation of toxic reaction which is not concerned with incompatibility between two bloods and which therefore is not to be anticipated by preliminary tests. This is:

3. *The possibility of a toxic disturbance in the circulation of the recipient by the introduction of blood which, though perfectly fluid, may nevertheless be undergoing incipient coagulative changes due to the physical influences to which it is subjected in process of transfer.*

Doerr¹³ observes that substances which influence coagulation also induce anaphylactic manifestations, and he states that coagulative changes may produce such toxic reactions as evidenced by work carried out in his laboratory by Moldovan. Although Doerr specifically states that these effects may result from the injection of undefibrinated homologous blood, the only protocols which we can find of such experiments are those reported by Moldovan¹⁴ demonstrating the fatal effects of the intravenous injection of recently defibrinated homologous blood. In explaining the anaphylactoid phenomena so produced, Doerr says that there is a reaction between an antigen and antibody, this being his conception of the incipient changes of coagulation according to the theory of Nolf, which results in an adsorptive action on the colloids of the blood, and that this latter disturbance—an essentially physical process—is responsible for the anaphylactic symptoms.

The mechanism of toxic action as suggested by this theory is similar to the two preceding hypotheses which we have introduced, in that all are concerned with adsorptive phenomena affecting the colloidal substances

6. Lindeman, Edward: Blood Transfusion, THE JOURNAL A. M. A., March 28, 1914, p. 993.

7. Lewisohn: Surg., Gynec. and Obst., 1915, xxi, 37.

8. Jobling and Petersen: Jour. Exper. Med., 1914, xix, 480.

9. Jobling and Petersen: Jour. Exper. Med., 1914, xx, 38.

10. Jobling and Petersen: Jour. Exper. Med., 1914, xx, 468.

11. Jobling and Petersen: Jour. Exper. Med., 1915, xxii, 401.

12. Zinsser and Young: Jour. Exper. Med., 1913, xvii, 396.

13. Doerr: Wien. klin. Wchnschr., 1912, xxv, 339.

14. Moldovan: Deutsch. med. Wchnschr., 1910, xxxvi, 2422.

of the blood or of the endothelial cells, and resulting in exposure to chemical change or physical shock. But the special significance of this explanation of toxic action is in connection with the physical influences to which the blood is exposed by the method of transfusion. Defibrination, either by agitation with glass beads or by "whipping," is, of course, an extreme degree of physical violence, but the argument of toxic effect from incipient coagulative factors applies, *pari passu*, with the lesser forms of physical violence which are incident in varying degree to any method of transfusion which is not a direct anastomosis. For instance, Minot¹⁵ has conducted some experiments which demonstrate that blood allowed to remain in brief contact with the glass barrel of a syringe before being mixed with oxalate solution almost always clots when chloroform is added, whereas blood allowed to run from an artery through a paraffined cannula directly into oxalate solution does not clot on the addition of chloroform. Minot's explanation of this is that chloroform renders inactive the antithrombin, thus allowing any free thrombin present to clot the fibrinogen; so that in an oxalated blood the presence of free thrombin indicates that this thrombin was formed in the blood before the blood met with the oxalate influence. The effect of the glass syringe as causing this incipient thrombin formation is, of course, evidenced by his experiments.

The adhesive forces of capillary action, so-called, or, more definitely, the effect of surface tension forces, which act in and about any zone of contact with moistenable foreign surfaces, must be to set free thromboplastic substances in the blood by abrasion and disintegration of platelets. Other more obvious physical violence, such as forcing through small artificial channels, pressure from piston syringes, etc., would naturally increase this effect.

Anticoagulants.—The rational and most desirable function of an anticoagulant is therefore, in our conception of the premises, to restrain *incipient* coagulative changes at the zones of foreign contact, where such changes are most liable to occur. The use of an anticoagulant to render the blood incoagulable in mass, as with the calcium-converting agents, merely as an expedient for handling the blood in a fluid state, does not seem to us justifiable, not only because this involves mixing with the injected blood a much larger quantity of the anticoagulant than is necessary, but also because, by calcium inactivation, only the end-result of the clotting process is prevented. What assurance is there, for instance, that blood rendered incoagulable by citrate, but which has been subjected to physical contact with glass and metal surfaces, does not contain free thrombin, as evidenced by Minot's investigation, or a considerable amount of thromboplastin derived from contact with tissue juices and from the disintegration of platelets? In this connection it should be said that the preserving influence on the platelet cell membrane which is ascribed to sodium citrate can have no effect on the platelets which have already been disintegrated by passage through foreign channels before meeting the citrate influence.

Paraffin.—The natural advantage of the paraffin method for the transfusion of blood is the effectiveness of hardened paraffin as an anticoagulant without admixture of any foreign substance to the transfused blood. Objection to paraffin has nevertheless been made on the ground that special care is necessary to

insure a perfect coating of the transfusion pipets or cylinders. It cannot be denied that special care in the matter of coating is essential to success, but the process is not a difficult one, and the mere trouble of coating in advance is offset by the convenience of having the pipets always ready for immediate use without further preparation.

Herudin.—The advantages of leech extract as an anticoagulant for purposes of transfusion may be briefly summarized: The derivation of herudin, the product of specialized glandular activity of the leech, indicates its biologic adaptation to the single purpose which it is known to serve in nature, namely, to prevent coagulation and thereby to facilitate digestion of blood in the stomach of the leech. From a physiologic standpoint it should be, and until recently has been regarded as, the ideal anticoagulant for animal experimentation. It acts not only as a preventive of platelet disintegration,¹⁶ but it appears also to exert a neutralizing or binding action on the positive factors of coagulation (thromboplastin, prothrombin)¹⁷ when these have been rendered active as a result of platelet disintegration.¹⁸ Furthermore, it does not, as far as we know, disturb the protective lipoids of the tissue cells, as has lately been suggested¹⁹ to be the action of the calcium precipitants and other calcium-converting agents.

As against the advantages which have just been enumerated, a very serious practical objection to leech extract has arisen in the uncertainty of procuring a reliable preparation of this substance. Quite recently Turner, Marshall and Lamson²⁰ discovered that a certain lot of the commercial preparation of "Hirudin" manufactured by Sachse & Co. of Leipzig, purchased by them in the early part of 1915 for use in their experiments with plasmapheresis, contained some very poisonous substance. Marshall,²¹ in a special investigation of this subject, reviews the findings of other workers with herudin and reports experiments showing the action of aqueous extract of leech heads, as prepared by Abel, Rowntree and Turner²² in comparison with similar experiments showing the action of the commercial preparation which contained the toxic principle previously mentioned. Marshall found that 20 to 25 mg. of this toxic preparation per kilo of body weight proved fatal for dogs, and ascribes its toxicity not to anything obtained from leech heads, but to some accidental product of putrefaction resembling sepsin. He confirms the findings of Abel, Rowntree and Turner,²² that the aqueous leech extract, as obtained by their method, is nontoxic in quantities which render the circulating blood incoagulable. While referring to toxic reactions from the use of herudin it is important to call attention to the reactions noted by some observers who have employed much larger quantities of this substance for transfusion than we have advised. Lewisohn,⁷ in attempting to employ herudin for his method of transfusion, experimented with the

16. Bürker: Arch. f. d. ges. Physiol., 1904, cii, 92; Zentralbl. f. Physiol., 1907, xxi, 653. Deetjen: Ztschr. f. physiol. Chem. (Hoppe-Seyler's), 1909, lxi, 1.

17. Mellanby: Jour. Physiol., 1909, xxxviii, 491.

18. Morawitz: Deutsch. Arch. f. klin. Med., 1904, lxxix, 215. Bayne-Jones: Am. Jour. Physiol., 1912, xxx, 74.

19. Overton: Arch. f. d. ges. Physiol., 1904, cv, 176. Boehm: Arch. f. exper. Path. u. Pharmacol., 1914, lxxv, 230. Clark: Jour. Physiol., 1913, xlvii, 66. Clowes: Proc. Soc. Exper. Biol. and Med., 1913, xi, 1. Clowes: Ibid., p. 4. Clowes and West: Ibid., p. 6. Salant and Hecht: Am. Jour. Physiol., 1915, xxxvi, 126.

20. Turner, Marshall and Lamson: Jour. Pharm. and Exper. Therap., 1915, vii, 145.

21. Marshall: Jour. Pharm. and Exper. Therap., 1915, vii, 157.

22. Abel, Rowntree and Turner: Jour. Pharm. and Exper. Therap., 1914, v, 257.

15. Minot: Am. Jour. Physiol., 1915, xxxix, 137.

intravenous injection of large quantities of this drug and reported that the dose of 100 mg. of herudin caused death in one dog, no ill effect in a second dog, and most alarming symptoms in a postoperative patient, so that she was in a precarious condition for over thirty-six hours. Another physician has informed us that he employed this same dose (0.1 gm.) in transfusing two patients by means of Kimpton's cylinders. In one case there was no reaction whatever, and in the other case a very severe one. Whether or not any of these reactions were due to the use of a toxic preparation of herudin cannot be ascertained, but the dosage noted above is far in excess of what we have recommended or deemed advisable. Our observations with herudin have been confined to the commercial preparation manufactured by Sachse & Co., comprising five lots of 10 mg. tubes

by chills and febrile reactions; and as these symptoms were associated only with the latter half of our herudin series, we are inclined to believe that they indicated some fault in the herudin which was used. Although we have never had any such reactions in our experimental transfusions with herudin while following precisely similar technic, this may be explained by the fact that we were working in the laboratory with a different lot of herudin; or it may be that dogs are less susceptible than human beings to the contaminating toxic substance which was discovered in the commercial "Hirudin" by Marshall, just as this author found that rabbits were much less susceptible to its effects than dogs. Other disadvantages of leech extract are that its activity is destroyed by heat sterilization, that it must therefore be produced under rigidly aseptic

TABLE 1.—PRECIPITATION REACTIONS WITH CALCIUM CHLORID

Substances in Aqueous Sol.	Reaction to Litmus (L) and Phenolphth. (P)	Addition of 1% CaCl ₂ Sol. in Small Amt.	Addition of 1% CaCl ₂ Sol. in Excess	CaCl ₂ Excess + Dil. HCl to Acid Reaction	Excess of Aq. Sol. of Substance to Already Formed Ppt.
Sodium citrate.....	Faintly alk. L.; Neg. P.	No ppt.	No ppt.	No ppt.	
Magnesium sulphate.....	Neg. L.; Neg. P.	No ppt.	No ppt.	No ppt.	
Sodium tartrate.....	Faintly acid L.; Neg. P.	No ppt.	Ppt. forms slowly on standing	Ppt. remains.....	Ppt. remains
Sodium metaphosphate.....	Faintly acid L.; Neg. P.	No ppt.	Ppt.	Ppt. remains.....	Ppt. dissolves
Dipotassium phosphate.....	Neg. L.; Neg. P.	Ppt.	Ppt.	Ppt. remains.....	Ppt. remains
Sodium fluorid.....	Alk. L.; Alk. P.	Ppt.	Ppt.	Ppt. remains.....	Ppt. remains
Sodium oxalate.....	Alk. L.; Alk. P.	Ppt.	Ppt.	Ppt. remains.....	Ppt. remains

TABLE 2.—EXPERIMENTAL TRANSFUSIONS WITH SODIUM CITRATE

	Amount of Solution Used, c.c.	Amount of Blood, c.c.	Amount of Anticoag. to 100 c.c. Blood, mg.	Clotting Time in First Tube, Min.	Clotting Time in Second Tube, Min.	Clotting Time of Residual Blood in Pipet, Min.
Sodium citrate: 10% sol. (1.40 m.). (Transfusion 1, Dog 1)	3	215	140	38	45	No sign of coagulation at 67½
Sodium citrate 10%, NaCl 0.8%, equal parts. (Transfusion 2, Dog 1)	1.8	215	42	10½	11½	No sign of coagulation at 50½
Sodium citrate 10%, NaCl 0.8%, equal parts. (Transfusion 7, Dog 2)	1	215	23	15½	10½	9½
Sodium citrate: 6.46% sol. (0.9 m.). (Transfusion 9, Dog 3)	1	215	30	36½	27½	16½
Herudin 10 mg. in NaCl 0.9% sol. (4.5 c.c.). (Transfusion 10,* Dog 3)	1.5	215	1.5	11½	16½	16½
Sodium citrate 10% sol. (1.40 m.). (Transfusion 11, Dog 3)	1	215	46	11½	19	32½
Sodium citrate 10% sol. (1.40 m.). (Transfusion 15, Dog 4)	1	215	46	16½	17½	31½ †
Sodium citrate 10% sol. (1.40 m.). (Transfusion 16, Dog 4)	1	215	46	23½	23½	90
Paraffin only, full coating. (Transfusion 17,* Dog 4)	0	215	0	12½	12½	Trace of fibrin at 39, still fluid at 24 hours ‡
Sodium citrate 10% sol. (1.40 m.). (Transfusion 27, Dog 6)	1	265	38	18½	18½	No sign of coagulation at 24 hours
Sodium citrate 10% sol. (1.40 m.). (Transfusion 28, Dog 6)	1	265	38	12	23½	No sign of coagulation at 90
				21	21½	
				27½	25½	
				15½	15½	15½
				17½	17½	18
				7½	no clot at 60	No clot at 60
				11½		
				9½	no clot at 54	45%; 52 (incomplete)
				12½		

* Transfusion 10 and transfusion 17 with herudin and paraffin, respectively, are included in the table for purposes of comparison. Where two sets of figures are given in columns five, six and seven, the first refers to first appearance of fibrin, the second to time of complete coagulation; where but one figure is given, the time of complete coagulation is indicated unless otherwise stated.

† Stained smear of this blood made at forty-three minutes showed normal staining and morphology.

‡ Stained smear of this blood made at forty-two minutes showed normal staining and morphology.

of this substance purchased prior to Jan. 1, 1915, and one lot purchased subsequently. We have never used, whether experimentally or in clinical practice, more than 7 mg. of herudin for transfusing 500 c.c. of blood, and in no one of numerous experimental transfusions on dogs have we had any ill effects from herudin in this proportion of admixture, although in some such cases we have removed and reinjected considerably more than the estimated total volume of the dog's blood. We wish, however, to record here the following interesting facts in this connection, which are derived from our series of clinical transfusions. We have made 14 clinical transfusions with herudin and 13 with paraffin. Among the 13 paraffin cases there were no chills or febrile reactions, nor have we had any such manifestations following our citrate transfusions. On the other hand, 6 of our 14 herudin transfusions were followed

by chills and febrile reactions; and as these symptoms were associated only with the latter half of our herudin series, we are inclined to believe that they indicated some fault in the herudin which was used.

Inorganic Salts.—Many investigators have reported the effects on blood from a purely physiologic standpoint of various anticoagulant substances of inorganic origin. The work of Bürker and Deetjen,¹⁸ Gessard,²³ Zucker²⁴ and Clowes,²⁵ is especially worthy of note in this respect. The observations here reported were made in the course of an experimental study of the anticoagulative effects of certain calcium-converting agents, to determine the minimal proportions of admixture with blood which would be needed for transfusion

23. Gessard: Compt. rend. Acad. d. sc., 1911, cliii, 1241; 1913, clxvii, 799.

24. Zucker: Proc. Soc. Exper. Biol. and Med., 1914, xi, 60.

25. Clowes: Proc. Soc. Exper. Biol. and Med., 1913, xi, 1; *ibid.*, p. 4. Clowes and West: *Ibid.*, p. 6.

with our apparatus and the relative safety and utility of these agents for this purpose. The following substances were selected for examination: sodium citrate, magnesium sulphate, sodium tartrate, sodium metaphosphate, dipotassium phosphate, sodium fluorid and sodium oxalate.

By reference to Table 1 it will be seen that, with aqueous solutions of sodium citrate and magnesium sulphate, no precipitate is formed on the addition of calcium chlorid, whether in acid or alkaline solution or in any proportion of admixture. With sodium tartrate and sodium metaphosphate a permanent precipitate is formed only when there is a considerable excess of calcium chlorid. In the case of sodium metaphosphate the precipitate caused by excess of calcium chlorid disappears with the further addition of sodium metaphosphate solution.

In determining the anticoagulative effect of these substances the plan of study was as follows:

A dog was anesthetized with ether and the jugular and femoral veins were exposed and served as donating and

was laid aside and this portion as well as the two portions in the paraffin-lined tubes were inspected at short intervals for evidence of coagulation, and the elapsed time from the entry of the tip of the pipet in the donating vein to the first appearance of fibrin formation was noted, as well as the time when firm clotting occurred.

Inasmuch as dipotassium phosphate, sodium fluorid and sodium oxalate form precipitates with calcium chlorid (Table 1) they were left out of practical consideration at the outset and very little was done to determine their relative anticoagulative effects. Preliminary trials indicated that a solution of sodium tartrate equimolecular with 0.9 per cent. sodium chlorid (m 1.54) had only a feeble anticoagulative influence. A hypertonic solution (m 4.35) was therefore employed, but even with this concentration it was necessary to use from twice to ten times the amount necessary to coat the pipet in order to produce any appreciable anticoagulative effect. Magnesium sulphate in m 1.54 solution showed a greater anticoagulative action than sodium tartrate, but neither showed sufficient

TABLE 3.—EXPERIMENTAL TRANSFUSIONS WITH SODIUM METAPHOSPHATE

	Amount of Solution Used, c.c.	Amount of Blood, c.c.	Amount of Anticoag. in 100 c.c. Blood, mg.	Clotting Time in First Tube, Min.	Clotting Time in Second Tube, Min.	Clotting Time of Residual Blood in Pipet, Min.
Sodium metaphosphate 1.57% (Transfusion 18, Dog 4)	1	215	7.3	11½; incomplete at 24 hours	No sign of coag. at 24 hours	No sign of coagulation at 24 hours
Sodium metaphosphate 0.8% (Transfusion 21, Dog 5)	1	215	3.7	9½ 10½	10½ 15	5½ 9½
Sodium metaphosphate 1.2% (Transfusion 25, Dog 6)	1	265	4.5	8½ 12½	22½ 24½	No sign of coagulation at 70
Sodium metaphosphate 1.2% (Transfusion 26, Dog 6)	1	265	4.5	12 16½	19½ 28½	No sign of coagulation at 64½
Sodium metaphosphate 1.46% (Transfusion 30, Dog 6, nine days after last transfusion)	1	265	5.5	13 17½	No sign of coag. at 2 hours	No sign of coagulation at 2 hours
Paraffin only. (Transfusion 31, Dog 7)*.....	0	265	0	17 19	16½ 18½	16½ 18
Sodium metaphosphate 1.46% (Transfusion 32, Dog 7)	1	265	5.5	13½ 16½	18½ 24½	13½ 9
Sodium metaphosphate 1.5% (Transfusion 33, Dog 7)†	1	265	4.0	12½ 13½	8½ 9½	11
Sodium metaphosphate 1.46% (Transfusion 34, Dog 7)	1	265	5.5	8½ 12½	8½ 11½	† Faint trace of fibrin at 11. No further sign of coag. at 24 h.r.
Sodium metaphosphate 1.46% (Transfusion 35, Dog 7)	1	265	5.5	7½ 11½	18½ 19½	13½ 15½
Paraffin only. (Transfusion 36, Dog 7)‡....	0	265	0	18 22	16½ 17½	14½ 18

* Coagulation time of Dog 7, at commencement of operation (4:10 p. m.) was 6 min. 15 sec. The coagulation times here noted were determined by allowing blood from the jugular vein to flow through a gold cannula (15 mm. long and 2.3 mm. diam.) directly into a clean glass test tube 15 mm. in diameter. Ten c.c. of blood was collected for each determination. The fluidity of this blood was examined every ten or fifteen seconds, and the time noted when the blood was sufficiently coagulated to permit inversion of the tube.

† Coagulation time of Dog 7, (4:42 p. m.) was 9 min. and 30 sec.

‡ Stained smear of this blood made at forty-five minutes showed normal staining and morphology.

§ Transfusions 31 and 36 with paraffin only were interpolated for purposes of comparison.

¶ Coagulation time of Dog 7 at end of operation (5:12 p. m.) was 7 min. 10 sec.

recipient vessels for autotransfusion. The transfusions were in all cases carried out by means of our special pipets and cannulas. During process of transfer, the pressure exerted on the blood averaged 40 mm. of mercury, and was controlled by a simple automatic contrivance so as never to exceed 80 mm. of mercury. The pipets were calibrated, of 300 c.c. capacity, and, just previous to use, were coated internally with an aqueous solution of the salt under examination. In each instance from 215 to 265 c.c. of blood were withdrawn from the jugular vein, which required about two minutes. As soon as the pipet was withdrawn from the cannula in the donating vein, about 5 c.c. of blood were immediately expelled from the pipet into a paraffin-lined test tube of 2 cm. diameter. The bulk of the blood in the pipet (from 190 to 240 c.c.) was then returned to the animal's circulation through a recipient cannula in the femoral vein, which required another two minutes, and, when this had been accomplished, another 5 c.c. portion of blood from the pipet was expelled into a second paraffin-lined test tube immediately on withdrawal of the tip of the pipet from the recipient cannula. The blood finally retained in the pipet, amounting to from 15 to 20 c.c.,

effect to warrant further investigation, inasmuch as both of these substances have been shown to have a toxic action when injected intravenously²⁶ which would render them undesirable for use in any quantity. It may be said, however, that in the amounts employed by us, neither of these agents appeared to exert any immediate effect on the experimental animal. The tabulated results of experimental transfusions with sodium tartrate and magnesium sulphate are omitted for the sake of brevity.

Having eliminated as undesirable five of the seven substances selected for study, there remains for consideration the action of sodium metaphosphate and sodium citrate. A synopsis of experimental transfusions with these agents is given in Tables 2 and 3.

Sodium Citrate.—The intravenous use of sodium citrate, in any quantity, as a therapeutic procedure, is open to ques-

26. Meltzer and Auer: *Am. Jour. Physiol.*, 1905-1906, xv, 387. Joseph and Meltzer: *Jour. Pharm. and Exper. Therap.*, 1909-1910, i, 24. Salant and Smith: *Am. Jour. Physiol.*, 1914, xxv, 239.

tion on theoretic grounds from the experimental findings of several investigators, notably Gros,²⁷ Clark,²⁸ Sakai,²⁹ Salant and Hecht.³⁰ There is, however, the recent assurance of those who advocate sodium citrate as an anticoagulant for transfusion that this substance can be injected with transfused blood in considerable quantity. The amounts used by them in clinical practice range from 1.0 gm. (Weil) to 0.2 gm. (Lewisohn) per 100 c.c. of transfused blood. These authors say that they have seen no undesirable effect of such dosage, although Weil has noted a change in the coagulation of the blood. In commenting on the change in coagulation time, which is induced by citrate transfusion, Lewisohn⁷ says that 300 c.c. of a dog's blood removed by his method of transfusion mixed with citrate (0.2 per cent.) and reinjected changed the coagulation time of the dog's blood from five minutes before operation to ten seconds when tested three minutes and again six minutes after operation.

Rous and Turner,³¹ in reporting their findings with respect to preservative fluids for red blood cells, refer to the transfusion of exsanguinated guinea-pigs and dogs with citrated blood by Weil,³² and state that, according to their observations, human cells and sheep cells rapidly disintegrate in citrated plasma, whereas in the preservative solutions which they have devised these cells survive for considerable periods.

Perfusion experiments indicate that sodium citrate has a potential toxic effect on the tissues of the circulatory apparatus which may be offset by the presence of available calcium, and that this toxic effect probably is prevented to some extent by the protective action of calcium ions in the tissues of the recipient. The mechanism of this protective action of the tissue calcium against the citrate ion is still a matter of speculation, but a most interesting and plausible theory of action has lately been formulated by Clowes³³ in a series of papers previously referred to. The extent to which this calcium-ion conversion occurs in the circulating blood and in other tissues would seem to depend on the amount of citrate which is introduced. Gros²⁷ and Salant and Hecht³⁰ found that sodium citrate has a toxic action on the isolated perfused frog heart, the latter authors demonstrating a well-marked effect in a concentration of thousandth-normal in Ringer's solution, and that it exceeded oxalate and tartrate in toxicity. Some as yet unpublished experiments in Professor Salant's laboratory in Washington³³ have shown that the intravenous injection of 100 mg. of sodium citrate per kilo in rabbits has caused serious symptoms and death.

Whatever the mode of action of citrated blood in transfusion, it is evident that when the citrate-ion concentration is sufficient to induce permanent incoagulability in vitro, the physicochemical effect in the circulation of the recipient must be considerable, and that this effect probably varies in direct ratio with the amount of citrated blood which is introduced. For purposes of transfusion, minimum amounts of this agent would therefore seem advantageous, and this point we believe worthy of special emphasis. We have previously found that the pipet-cannula method affords a technic which permits approximately one-eighth of the quantity of herudin which is necessary to produce an equal anticoagulative effect when using the simple syringe method; and a similar advantage is shown, by the present experiments, to hold true with sodium citrate. In coating the pipets with this anticoagulant, 20 to 30 c.c. of a 10 per cent. solution of sodium citrate is poured into the pipet just previous to use. This fluid is thoroughly flowed over the interior of the pipet by rotating it in the horizontal position and all excess is then allowed to drain out through the tip, leaving

approximately 1 c.c. of the solution which adheres as a thin film to the glass wall of the cylinder. We have found (Table 2) that 100 mg. of sodium citrate (1 c.c. of solution), when so used as a liquid coating in one of our pipets, is amply sufficient for the purpose of transfusing 250 c.c. of blood when used with our cannulas, and that the only fully citrated blood so treated is that small residual portion (about 12 c.c.) which is retained in the pipets. That this residual portion of blood is usually subjected to a greater anticoagulant influence than the blood which is discharged from the pipet, may be seen by referring to the tables, and appears to be due to contact with a relatively larger proportion of the anticoagulant solution, which tends to remain adherent to the walls of the cylinder. Without making any deduction on account of this greater proportion of citrate solution, which presumably adheres to the pipet and is not transfused, but remains in the residual portion of blood, it is evident that not more than 38 mg. of sodium citrate per 100 c.c. of blood is necessary for transfusion by this method. This proportion of citrate is from one twenty-fifth to one fifth of the amounts which are recommended by those who have used citrate as an anticoagulant with other methods of transfusion, and we believe that this proportion can produce very little disturbance of the circulating blood of the recipient, inasmuch as there was no noticeable effect on the coagulation time of the blood of the animals used in our experiments.

During the past six months we have adopted this use of sodium citrate as an anticoagulant with our pipet-cannula apparatus in clinical practice and are well satisfied with its practical operation.

Sodium Metaphosphate: (NaPO_3)_x. The sodium metaphosphate used in these experiments was of C. Kahlbaum's manufacture, and consisted of broken bits of vitreous fused plates, having a conchoidal fracture and a solubility of 5.5 parts in 100 parts of distilled water at 15 C. A phosphorus determination, made for us in the laboratory of Messrs. Eimer and Amend, indicated that it contained two atoms of phosphorus in its molecule, corresponding to the formula (NaPO_3)₂ or $\text{Na}_2\text{P}_2\text{O}_6$.

As to the toxic effect on animals of this salt, our preliminary experiments by intravenous injection of 1.6 per cent. solution in rabbits show that the minimum lethal dose for rabbits is between 80 and 100 mg. per kilogram of body weight.

The results obtained in experimental transfusions with this agent are shown in Table 3 and indicate, by comparison with Table 2, that it has a more powerful anticoagulative effect than sodium citrate in proportion to its weight. The susceptibility to recalcification of blood treated with sodium metaphosphate as compared with blood treated with sodium citrate is a matter which we have, as yet, only touched on in our experimental work. We cannot therefore make any statement on the subject, as we have insufficient data for definite conclusions. It would appear, however, to be worthy of further investigation as tending to show what, if any, derangement of the coagulative mechanism is produced by these agents as regards subsequent reactivation by ionized calcium.

SUMMARY

1. A better knowledge of the physicochemical changes which may occur in the blood of the recipient as a result of transfusion in varying pathologic conditions is needed in order to define more clearly the rational indications for this operation as a therapeutic measure.

2. Certain hypotheses are suggested to explain the occurrence of febrile and toxic reactions following

27. Gros: Arch. f. Exper. Path. u. Pharmacol., 1913, lxxi, 395.
28. Clark: Jour. Physiol., 1913, xlii, 66.
29. Sakai: Ztschr. f. Biol., 1914, lxiv, 1.
30. Salant and Hecht: Am. Jour. Physiol., 1915, xxxvi, 126.
31. Rous and Turner: Proc. Soc. Exper. Biol. and Med., 1915, xii, 124.
32. Weil, Richard: Sodium Citrate in the Transfusion of Blood, THE JOURNAL A. M. A., Jan. 30, 1915, p. 425.
33. Personal communication.

transfusion which cannot be prognosticated by the existing laboratory tests for blood compatibility.

3. Importance is attached to the belief that many of the febrile and toxic reactions, not to be explained by hemolysis or agglutination, are due to the transfusion of blood which is undergoing incipient coagulative changes, or which contains potential coagulative factors such as thromboplastin and thrombin.

4. In judging of the safety of any method of transfusion, the extent and mode of action of the anticoagulant which is used, the toxicity of the substance, the amount of it which is necessary, and the mechanical technic in transferring the blood are the most important factors to be considered.

5. The employment of an antithrombin such as leech extract or the commercial "Hirudin," while appearing to be ideal as an anticoagulant from the physiologic standpoint, is subject to uncertainty of action and possible danger from contamination with toxic substances. Minimum doses of tested and standardized preparations and optimum anticoagulative technic should be used if such agents are employed.

6. The use of paraffin is theoretically the method of choice in providing an anticoagulant agent for purposes of transfusion and has been found practicable, but not without certain precautions and proper apparatus.

7. Of the inorganic calcium-converting agents which we have investigated experimentally, the citrate of sodium and the metaphosphate of sodium are the only substances which exhibit desirable characteristics. Both of these salts, however, have toxic effects which demand consideration of the amounts which can be safely employed for purposes of transfusion. Our experience with sodium metaphosphate is limited to experimental transfusion on animals.

8. Sodium citrate in the proportion of 120 mg. to 300 c.c. of blood, when used with the pipet-cannula apparatus and in the manner described, serves as an effective anticoagulant for transfusion. This proportion (0.04 per cent.) is from one fifth to one twenty-fifth of the amount of citrate necessarily employed with other methods. Because of the considerations which have been discussed, we believe it to be the best method of transfusion for general use.

Street Dust and Disease.—Following the announcement by the public health committee of the New York Academy of Medicine that a system of street cleaning which does not eliminate gross street dust and which employed dry sweeping methods cannot be satisfactory, a study was made by Prof. Thomas W. Hastings of Cornell University Medical School concerning the relation of street dust to disease. On examination of the available data he summarized his findings about as follows: Numerous bacteriologic examinations made in New York as well as elsewhere show that street dust contains a variety of living pathogenic organisms such as tubercle bacilli and various types of streptococci which are recognized as causative agents of many respiratory and other diseases. Although few authentic cases of disease can be traced to bacteria in street dust, yet the fact that appreciable numbers are found in street dust and inhaled or ingested, establishes a presumption that street dust may be the cause of disease. Studies of the incidence of respiratory diseases shows that persons free from exposure to city dust are less liable to suffer from such diseases. Dust has long been recognized as an injurious mechanical irritant and as a cause of lowered bodily resistance. Dry sweeping should be entirely abandoned, and street flushing should be introduced in street cleaning processes. All efforts should be made to render streets as free from dust as possible.

REACTIONS FOLLOWING BLOOD TRANSFUSION BY THE SYRINGE CANNULA SYSTEM*

EDWARD LINDEMAN, M.D.

NEW YORK

Since the simplification of technic,¹ blood transfusion has acquired considerable popularity. The syringe cannula system has greatly simplified the procedure over the old arteriovenous method. It now occupies a prominent and permanent place in the field of therapeutics. Today a hundred transfusions are performed where one was done but a few years ago. Though the technic has been thus simplified, there is a tendency on the part of some to slide over details of the work and, unequal to the task of even simple technic, they may resort to anticoagulants or inferior methods by which it may be possible to put blood into the patient even subcutaneously. The attempt to simplify the method further has not been attended without the sacrifice of important points of merit. The pitfalls and errors in the further simplification are the very things I have purposely avoided by the method originally outlined. The operative procedure is a life saving measure and should be regarded as something more than an injection of saline fluid or salvarsan administration, and any tendency to curtail detail of execution to meet the convenience or inexperience of the operator at the expense of the patient should be discouraged.

The reactions following transfusion will deter some from employing the measure with a patient in a critical state, in the fear of doing harm by the procedure itself, and the possible benefits to be derived may be overbalanced by the harm done. In order for any therapeutic measure to gain wide favor, it must be free of any danger incident to the procedure itself, or at least a very high percental relationship must obtain between risk and gain. It is my object in this communication to take up the problem of reactions to transfusion, to analyze their cause and to show the way to prevention.

In the first eighteen transfusions which I performed by the syringe cannula system, preliminary blood tests were made in only two instances. In this group of eighteen cases there were no chills and no fever reaction above 2 degrees. The quantity transfused was small—never more than 700 c.c. The donors used were immediate relatives of the patients, and the diseases treated were not primary blood diseases. (Primary blood diseases and infection offer most difficulty in finding compatible donors.) The conspicuous absence of hemolysis in the absence of blood tests in sixteen of these cases must be attributed to good fortune. Later in the course of the work, though blood tests were made in nearly all the cases, chills and fever were encountered more frequently.

In the first 150 transfusions which were performed by the syringe cannula system,² the incidents of chill followed by fever were approximately 33 per cent. The quantities transfused were usually 1,000 c.c. or more in adults. The work was done in many different

* Because of lack of space, this article is abbreviated in THE JOURNAL by omission of Table 1. The complete article appears in the author's reprints. A copy of the latter will be sent by the author on receipt of a stamped addressed envelope.

1. Lindeman, Edward: Simple Syringe Transfusion with Special Cannulas, *Am. Jour. Dis. Child.*, July, 1913, p. 28.

2. Lindeman, Edward: Blood Transfusion, *THE JOURNAL A. M. A.*, March 28, 1914, p. 993.