epidermis at the apex of the lesion was considerably thickened; although not nearly so much as in similar tumors occurring on the hands and feet. In the sections stained for elastic fibers there was a striking difference between the normal integument used as a control and the angiokeratomatous tissue. In the former, the elastic filaments were long, wavy, and sharply defined, and stained evenly and well. Surrounding the vessels there was a close, almost solid network of fibers, the appearance presented being suggestive of a small tube, snugly encircled by numerous turns of fine wire and passing at right angles through a mass of similar material, the strands of which closely encircled and intermingled with those forming the outer covering of the cylinder. In the affected skin, elastic tissue was to be found in all of the regions in which it usually occurs, but the fibers were short, fragmentary, and irregularly arranged, and apparently furnished very little support to the vessel walls. The coats of the capillaries appeared to be normal, but the surrounding network of "braces and guy-ropes," which was so prominent a feature in the control sections, was almost entirely lacking. This held true not only in the immediate vicinity of the tumors, but in the microscopically normal areas as well. In those sections which contained lacunæ, the elastic tissue immediately surrounding the vessel was apparently normal in amount, although comparatively thinned because of the increased diameter of the

Fig. 5.—Section from Case 2, stained with Weigert's solution, showing deficient amount of elastic tissue. (Spencer Obj. : $\frac{1}{4}$; no ocular.)

capillary; but the short, slender, poorly stained strands in the immediately adjacent areas appeared even less numerous and weaker than those in the more remote parts of the interlesional regions.

The findings in this case are of very great interest as pointing to a probable explanation of the cause of angiokeratoma. While it is the consensus of opinion among those who have carefully studied the disease that the lesions are the result of capillary dilatation, followed by thickening and other hypertrophic changes in the epidermis, the fact that there must be present some antecedent pathologic condition in either the vessel wall or in the structures surrounding it has never been emphasized, although it was suggested by Fordyce⁷ as early as 1896.

The majority of cases of angiokeratoma of the extremities have been seen in young persons, and practically all of the patients were subject to chilblains.

While some of the contributory factors to chilblain are well known, a careful and exhaustive histopathologic study of the affected tissue has never, so far as I can find, been made. We see very few instances of the dis-

order in the middle west, and I have not yet had an opportunity to secure material from a case. It is probable that alterations are to be found in the elastic elements in this disease also, but whether the changes precede, and, to some extent, superinduce, the localized congestion, or whether they develop as sequelæ, following repeated attacks of the malady, remains to be demonstrated. The matter is certainly worthy of thorough investigation. In those examples of angiokeratoma in which regions other than the extremities are involved, the case records enable us to wholly dismiss chilblain circulation as a possible contributory factor in the etiology. In the majority of cases, however, there is a history of venous obstruction, in the form of a varicocele, tumor of the epididymis, or postoperative venous occlusion, preceding the development of the angiokeratoma, and the evidence is almost conclusive that these little vascular tumors are simply the result of increased blood-pressure in superficial venous capillaries, the walls of which are insufficiently supported. The accompanying changes in the epidermis may be interpreted as Nature's attempt to prevent further dilatation, and possible rupture, of the involved vessels. Inasmuch as there is a fairly uniform insufficiency of elastic fibers throughout the integument of the scrotum, it is probable that the deficiency is congenital in origin.

The result of a careful consideration of the clinical histories of those cases of the disease in which the eruption was more generalized is also very suggestive of the possibility of a similar lack of elastic tissue in the affected areas.

610 Commerce Building.

INTRA-OCULAR NEOPLASMS

WITH A REPORT OF FIVE CASES OF NEOPLASM AND ONE OF SIMULATED NEOPLASM

F. PARK LEWIS, M.D.

BUFFALO, N. Y.

Malignant intra-ocular growths in children are rare, but in the eye-ball as in almost no other part of the body they are encapsulated. If they are discovered early enough, therefore, and enucleation is at once performed, life may thereby be saved. If, on the other hand, their presence is not recognized until the integrity of sclera or cornea has been destroyed and the orbital or extra-orbital tissues has become involved, the invasion of the brain and face with the separation of the cranial bones results in such a horrible method of death that even if the number of cases of retinal glioma is not great, it is worth while to emphasize the characteristic diagnostic features in order that no child may be denied the right of help—while help is yet possible.

The possibilities are that if all cases were reported, their aggregate number might not be so small. Wintersteiner considers that they constitute 0.04 per cent. of all eye diseases. Of 204,612 cases presenting at the New York Eye and Ear Infirmary in ten years, 0.017 were glioma retinae. It is not at all unlikely that in the country districts or in the small towns remote from large eye clinics, cases which are never reported occur and many more children die than the records would lead one to suppose. The etiology is still uncertain. A number of instances are reported of several cases occurring in a single family, and the assumption has been drawn there-

* Read in the Section on Ophthalmology of the American Medical Association, at the Sixty-Second Annual Session, held at Los Angeles, June, 1911.

Fig. 1.—Case 1; long band of glomatous tissue extends from optic nerve through the vitreous to the lens. May not a vestigial hyaloid artery have been the channel of development? No recurrence after nine years.

Fig. 4.—Case 3; gloma originating at margin of the optic disk. Beginning staphyloma. Enucleation. No recurrence after eleven years.

Fig. 2.—Case 2; glioma retinæ; neoplasm absolutely fills posterior chamber of eye. Partial ectasia of sclera with involvement of the optic nerve; enucleation; no recurrence after three years.

Fig. 5.—Case 4; melanosarcoma; enucleation; no recurrence after a year.

Fig. 3.—Case 2; glioma retinæ. Closely massed round cells, large nuclei; no rosettes found.

Fig. 6.—Case 4; melanosarcoma. Coarse masses of pigment interspersed with spindle-shaped cells, originating in the choroid.

fore that heredity exercises a controlling influence. Van Hoffman, Jr.,¹ at the last session of the Heidelberg Ophthalmological Society, described a double case of glioma in a mother and her infant child. The mother had been operated on about twenty-eight years before. In the Royal London Eye Hospital reports for 1905, Owen related a similar case. A woman operated on in 1859 later bore a child which also had a glioma, likewise removed. A full sister of the second woman bore two children who had glioma. One was cured by operation, the other died of the disease. The author reports a third family case. He operated on a boy 2 years old in 1872. This boy married at 21 and his wife bore seven children. The second and third both had gliomata at the ages of two years and five months respectively. The first was operated on but suffered recurrence in orbit and both eyes. An operation was refused in the second case and both children died.

Many cases are reported of several children in one family having this disease, but in all of these as in the classical one described in the *Australian Medical Gazette*, in which, out of a family of sixteen children, ten had glioma, it seems far more likely that some congenital

Cohnheim's theory that an adventitious fetal development of the nerve fiber and ganglion cells results in the establishment of "rests." From these the glioma takes its origin. As Treacher Collins has shown, the glioma cells and those of the undifferentiated cells of the three-months fetus bear a close resemblance to each other, and it is more than possible that the developed tumor bears a direct relation to the halted development of the fetal retina, and the glia and ganglion cells may be in Axen-

Fig. 7.—Case 5; extradural glioma of optic nerve; enucleation of orbit. No recurrence afterward, although probable later.

fetal defect was present in each of them which made the development of the disease possible.

This is borne out by Fuchs' observations. A mother brought her 4-year-old child to his clinic with an advanced glioma of the right eye. The ball was much enlarged and the neoplasm was growing into the orbit. The entire orbital contents were removed, but the growth recurred and within a year the child died. Some months after the mother brought the second child, who likewise had a glioma which was removed, but which also returned, resulting in the death of the child. Soon afterward she brought her baby, then only a few months old, with a defect of its left eye. In this case it was not a glioma, but a congenital coloboma of the iris and chorioid from which Fuchs concluded that a congenital morbid disposition must have been present which was the basic cause of the glioma in each of the cases.

Recent pathologic studies have added little of value to our knowledge of the intrinsic nature of these tumors. The consensus of opinion seems to be in accord with

Fig. 8.—Case 5; extradural glioma of optic nerve; spider cells from glia.

feld's opinion the outgrowth of the undifferentiated cells of the tumor.

While much has been written concerning retinal glioma, no new facts have been deduced from our recent studies which to any degree change existing views as to the nature of these malignant tumors. The purpose, then, of this report is not to add to our pathologic knowledge of these rare growths, although the four described are of sufficient interest to merit special reference; but it is to urge that these rare conditions be brought more frequently before the notice of the general profession, because they occur so infrequently that a physician in general practice may allow the possibility of such a growth to pass wholly out of his mind, and because without distinctive ophthalmologic appliances, the recognition is so easy on the part of any intelligent observer,

Fig. 9.—Case 5; extradural glioma of optic nerve.

and because by its early recognition life may be saved in a large proportion of cases.

Every case should, therefore, be reported and if possible a section of the tumor made and the character of the neoplasm determined. We shall in this way not only get fuller statistical records as to the actual number of cases which occur, but we may, by calling attention to its manifest characteristics, render it possible to make a

1. Van Hoffman: Ann. d'ocul., 1910, cxliii, 32.

diagnosis before the growth has passed outside of the eye-ball and while enucleation may yet save the life of the child.

The yellowish reflex coming from the depths of the vitreous lead to the common name given to it by the earlier writers of "amaurotic cat's eye"; and if a light is allowed to shine obliquely into the large pupil, the whitish reflex coming from behind the lens together with the age of the patient is almost absolutely characteristic. When such cases present themselves, skilled advice, if obtainable, should be immediately sought. It is quite different in appearance from the whitish color of a cataract or the surface opacity of a leukoma. It may, indeed, be confounded with plastic deposits in the vitreous from the uveal tract, but these are rare in young children; and when they occur, the usefulness of the eye has probably already been lost.

The first three cases reported below are interesting as being typical of this form of neoplasm. The fourth, that having its origin in the optic nerve, is exceedingly rare. All of these cases occurred, as glioma almost invariably does, in children under 4 years of age. The fifth, that of melanosarcoma, was found in a boy of 13. At present there is no evidence of a recurrence. The sixth is of interest as being the result of an uveitis, possibly prenatal, developing a glaucomatous condition of the eye and simulating in appearance an intra-ocular tumor. An exact diagnosis was impossible until the eye had been removed and sectioned. The illustrations give more clearly than any verbal description the exact conditions which were found in the first five cases.

REPORTS OF CASES

CASE 1.—Glioma retinae; enucleation; no recurrence after nine years.

Patient.—A. F., of Pembroke, Genesee County, N. Y., the daughter of simple, honest, hard-working parents, was brought to my clinic on Nov. 26, 1910. She had nearly reached four years, and although there had been some peculiarity noticed in the right eye since she was 1½ years old, it was only during the past four weeks that a marked difference was observed by the parents.

Examination.—The child manifested considerable uneasiness, pulling her hair on that side and occasionally manifesting evidences of discomfort in the eye. The pupil was dilated and maximum and immovable. The cornea and lens were perfectly clear, while a lobulated whitish yellow mass could be seen apparently filling the posterior chamber. It was traversed by fine vessels, giving in places a light reddish reflection. Tension was apparently slightly increased. There was no response to light, and the eye was evidently perfectly blind. The child was apparently in robust health and seemed to be unusually well developed.

Course of Disease and Operation.—The diagnosis of glioma was made and an enucleation advised. This was refused for the time, however, by the parents, and the child did not appear at the clinic until April 3, 1902. The tumor had then markedly increased, the eye-ball was flushed, injected and painful, tension plus 2, the lens pressed forward toward the cornea but still clear and in the upper portion of the mass, which was discernible when seen by reflected light, evidence of hemorrhage which had partially absorbed. It was no longer safe to postpone operative measures, so the child was taken into the hospital and an enucleation immediately performed. The eye was hardened and on section of it the diagnosis of glioma was confirmed. Figure 1 shows the extent to which the growth had filled the eye-ball. On April 15, 1911, the child was again brought to my clinic, when the orbit was found to be in perfectly normal condition and there was no evidence of any return of the growth, so that it may be considered to have been permanently cured by the excision.

CASE 2.—Glioma retinae; enucleation; no recurrence.

Patient.—C. S., the little daughter of well-to-do parents from another town in Genesee County, was brought to my office by her grandfather, a well-known physician of that place, on Feb. 15, 1908. The little girl was 3 years old. The previous summer she had fallen and struck her nose and from that time a glassy look appeared in the left eye, which seemed to diverge slightly. The day before bringing the child to me, the parents had covered the right eye and had discovered to their alarm that the child was unable to see from the left eye. The little girl had gray eyes, both pupils being normal and both reacting to light readily. There was no history of cancer and the family are quite sure that the child had been able to see out of the left eye for some time after she was born. Lately she had been running against objects and seemed unable to guide herself. For the past week she had been unusually irritable.

Examination.—The eye-ball showed no apparent redness externally; the cornea and lens were both clear, but in the vitreous, flocculent masses that seemed of a creamy whiteness could easily be seen by oblique light without the aid of the ophthalmoscope.

Course of Disease and Operation.—The diagnosis of glioma was made and assurance given that within a short time the eye must be removed. It was, however, two months before final permission was obtained. Then tension had rapidly become increased, the eye being injected. The child was immediately taken to the hospital and the eye-ball removed, the optic nerve being divided as far back as possible. A section of the eye was made after it was suitably hardened, and Figure 2 shows that the vitreous had become completely invaded by the gliomatous mass. Had enucleation been much longer delayed, the growth would have passed through the ocular structures, involving the orbit, when an exenteration of the orbit would have been required, and even that might not have saved the child's life.

CASE 3.—Glioma retinae; enucleation of eye-ball; no recurrence after eleven years.

Patient.—F. S., of Buffalo, was brought to my clinic at the Charity Eye and Ear Hospital on Oct. 16, 1900, aged 2½, one of eight children. The family was wretchedly poor, often lacking the necessities of life. The youngest of the family, a baby of 6 months, had had spasms from birth. F. had twice had measles; the left eye during the first attack had become swollen with much mucopurulent secretion. He had had a second attack last winter, at which time the eye had become inflamed but apparently had recovered again. The eye began to have a "hollow look," as described by the mother. Three weeks before I saw him, the child had fallen down-stairs, after which the eye-ball began to swell; and during the last three weeks, much of the time the child had held his hand to his head and given evidence of suffering.

Examination.—The cornea and lens were still clear, but a yellowish mass tinged with pink filled the vitreous. The red fundus reflex could still be seen around the margin of the enlarged pupil. The pupil did not react to light and tension was increased.

Operation.—An immediate enucleation was advised and made. The diagnosis of glioma on section was confirmed and the mass was found already occupying the vitreous space.

CASE 4.—Melanosarcoma in boy of 13 years; enucleation; no recurrence. As differentiated from these cases, all of which seem to have originated in the optic nerve, and in none of which was the patient older than 4 years, this case is of interest and value.

Patient.—C. B., of Falconer, Chautauqua County, New York, a schoolboy of 13½ years, was brought to my office by his parents, who gave a record of good heredity. The patient had, however, not been seeing from the right eye for some time.

Examination.—In this eye (the right) he was found to have light perception, recognizing only moving shadows, with entire loss of the temporal field, while the light could be perceived on the nasal side. In the left eye, with a plus .50 D. Sph. plus .50 D. Cyl. axis 180, vision equaled 20/20; external choroidal degeneration with scattered pigment masses was found to be present. A grayish elevated mass was discovered on examination with the ophthalmoscope, occupying nearly a

third of the nasal side and leaving a wide margin which was undisturbed between the growth and the optic disk.

The diagnosis of malignant neoplasm was made and the parents assured that enucleation must necessarily be made. The eye, however, was kept under observation for nearly a year before it was removed, when the growth which is shown in Figures 5 and 6 was shown to be a melanosaecoma.

CASE 5.—Intracanal glioma of the optic nerve; exenteration of the orbit. Glioma of the optic nerve is exceedingly rare. The report of the following case will be of value.

Patient.—Baby S. was brought to my office by Dr. William M. Mehl, of Buffalo, for consultation because of a marked exophthalmos. She is the little daughter of parents in moderate circumstances living in Allegheny County. One year previously, the mother noticed that the child frequently rubbed the right eye and shortly afterward it was seen to protrude. This protrusion had continued, slowly increasing to the time of examination.

Examination.—There was no evidence of pain and the child was well nourished. All the movements of the eye were still retained and notwithstanding the excessive proptosis, the child was still able to close the lids completely over the ball. It would seem that the lids had grown with the ball so as to accommodate themselves to the excessive protrusion. The eye could not be returned to the orbit with pressure, thereby excluding the possibility of an orbital vascular growth as the cause of the protrusion. There was no pulsation of the eye, nor was it apparently irritable or reddened. The pupil responded to light slightly. The vessels in the fundus were dark and tortuous, and a view of the disk, as far as it could be obtained in a nervous, unsteady child, showed a woolly and uncertain outline. The possibility of a gliomatous involvement was not considered because of the great rarity of tumors of this kind and it was thought that possibly a postauricular abscess or cyst might be causing the protrusion. The child, therefore, was placed under chloroform and an opening was made by the side of the eye-ball and the finger inserted, when it was found that a large roundish growth, apparently coming from the optic nerve, occupied the larger portion of the orbit. Incision was made into this growth, when it was found filled with a translucent honey-like yellowish mass, the tumor being separated by membranous septa. It was evident, therefore, that the growth was gliomatous.

Operation and Course of Disease.—The entire mass was removed on Aug. 13, 1910, by Dr. Mehl in my presence. Figure 8 indicates the character of the substance which was found to constitute the tumor. So extensive was the involvement of the optic nerve that the entire exenteration of all of the tissue in the orbit was made. It seemed inevitable that the growth must recur and an unfavorable prognosis was made. In March, 1911, however, the child was again seen and the tissues of the orbit had filled in apparently normally. The lid was not greatly sunken, and there was no evidence whatever of malignancy in the new tissue. It is quite possible, however, that ultimately the growth may recur.

CASE 6.—Iridochorioiditis with occlusion of the pupil simulating an intra-ocular neoplasm.

Patient.—E. H., of Fillmore, New York, was referred for consultation by Dr. S. Mitchell of Hornell, in January, 1910. From the October previous the child had been suffering with the right eye. One night a sharp pain developed in the eye and the next morning the eye-ball was very red and swollen. The patient was not seen by Dr. Mitchell for some months; then the eye was found deeply injected, the pupil contracted and filled with a plastic gray mass, and lens and iris pushed forward until it was in contact with the cornea. The eye-ball was intensely hard and evidently exceedingly painful. Dr. Mitchell had advised enucleation. The eye was evidently entirely blind.

Examination.—The conditions present when the boy was brought to my office were quite the same as those which Dr. Mitchell had found with the exception that they were intensified. The grayish mass in the pupil had recently become more marked. Glaucoma following a general uveitis occurring in a child three years old and involving only one eye, all led to the suspicion that a glioma was present and was rapidly developing.

Operation.—The eye-ball was removed by Dr. Mitchell and a section made, when it was found that there was no tumor in the eye whatever. The glaucoma was the result of the obstruction of the filtration angle and of the intense iridocyclitis, which in all probability had existed for a long time before Dr. Mitchell was consulted.

The case is important as one simulating an intra-ocular tumor and showing the great difficulty in diagnosis prior to enucleation. Happily in either instance, the eye had to be removed and diagnosis was one of secondary, although scientific, interest.

454 Franklin Street.

[THE DISCUSSION ON THIS PAPER WILL BE FOUND ON A LATER PAGE OF THIS ISSUE UNDER THE HEADING OF "SECTION DISCUSSIONS."]

THE IMMEDIATE REMOVAL OF TRAUMATIC CATARACT *

JOHN A. DONOVAN, M.D.
BUTTE, MONT.

A paper relating to the treatment of eye injuries, to be properly understood, should be confined to specific cases.

The impossibility of the universal application of any single line of treatment is well known to all surgeons, but especially so to those most frequently called on to treat eye injuries. In these cases it is often the most experienced who find it hardest to decide on the line of treatment.

Sweet, in his report¹ of 702 cases, says: "Unlike other operative procedures on eyes in which there is uniformity of technic, each foreign body injury must be studied as a separate and distinct condition."

Bowman² is quoted as saying: "I have never laid down a method of extraction of cataract universally applicable. I always operated as seemed best at the time for the individual treated."

Another difficulty in obtaining uniform results is the difference not only in classes of people, but classes of injuries. Sweet's report, which probably fairly represents the proportion of cases in the large cities, gives 81 per cent. magnet cases, while my cases are in almost inverse proportion. He gives 5 per cent. lens injuries. Here, among the penetrating wound cases the proportion of lens injuries is several times this percentage.

When the patient is seen immediately or soon after the injury, after he has recovered from the shock and before the local reaction has made careful work almost impossible, the lens should be extracted. This may be done under a local anesthetic, though a general anesthetic is preferable in many cases. If local reaction is already marked, the ophthalmologist should wait about four to six days when it subsides, then remove the lens before the secondary inflammatory symptoms have become severe.

A general anesthetic is necessary at this time. Though the breaking up of the lens and washing it out has been a recognized procedure for some time, a regular cataract extraction, combined or simple, has in many cases been more successful. If there is a large central corneal wound or if the lens is already badly mutilated, cutting it up with a cystotome and washing it out is the operation of choice. Unusual care must be exercised in doing this, as the zonula, if not already ruptured, may

* Read in the Section on Ophthalmology of the American Medical Association, at the Sixty-Second Annual Session, held at Los Angeles, June, 1911.

1. Sweet: Third Series of Cases of Injuries from Foreign Bodies, Tr. Am. Ophth. Soc., 1900.

2. Wood's Ophthalmic Operations, p. 1155.