the smaller nodules had been absorbed. Through my going home on leave this case was unfortunately lost sight of.

Conclusions.

Of the 4 tubercular cases treated 3 showed undoubted improvement, and in the fourth the signs were distinctly encouraging at the time the treatment was unavoidably discontinued. In all of these cases the injections were given locally into the nodules as well as intravenously, and it might be thought at first that the ulceration described was merely the result of irritating fluid in the nodule, and that there was only a local effect produced. The course of the cases described is, however, opposed to this view, as in none of them were the local injections given on the face, and yet the nodules on the face underwent great improvement, and to a large extent disappeared entirely. One of the mixed cases, too, which received no local injections into nodules improved greatly as regards the nodular portion of the disease.

From my experience, then, in these cases I am of opinion that it would be quite worth while to give this remedy a more extended trial and in a larger series of cases, using both intravenous injections and also local injections into the nodules, care being taken that the area of ulceration is controlled to a reasonable size, so as not to produce too severe a drain on the patient's health.

Post-mortem reports on the bodies of lepers appear to indicate that, at any rate for some years, in the vast majority of cases the disease does not produce any internal lesions; the bacilli are seldom found in the blood or urine and only very occasionally in the fæces. The disease is largely, then, one whose manifestations are confined to the skin, and the cure of which attempted by nature is by means of ulceration and sloughing of the leprosy deposits therein. Any drug, therefore, which will bring about this natural process in the nodules, causing extrusion of the deposits and with them of millions of bacilli, and leaving behind soundly healed, apparently healthy skin, is, I think, worthy of consideration. These remarks apply entirely and only to tubercular cases. In anæsthetic cases, where permanent changes have evidently taken place in nerve sheaths and fibres, the chances of benefit are obviously *nil*, and in any early case before such changes have occurred I have had no opportunity of trying this method of treatment.

Of the remedies that are being tried in various places at the present time the most diverse results are reported by different observers in the use of the same remedy. To take only four of these remedies, leproline, nastin, salvarsan, and chaulmoogra oil, and reading the extracts from various papers given in the *Tropical Diseases Bulletin* during the last two years, on them the authors express the most divergent conclusions. Taking leproline, Rutherford found no improvement in 32 cases, while H. Gougerot writes that "it has a distinct curative action," and "the best treatment we know of." Taking nastin, Schumacher and Max Rudolph had cases which improved under its use, while E. P. Minett considers that it has "very little beneficial effect." With regard to salvarsan, Gaucher and Boinet speak of it favourably, the course of treatment being combined with bee stings. Paldock had 9 cases which improved temporarily followed by a relapse, and Veillon and Serra consider it has no benefit. As to chaulmoogra oil, Kirby Smith found that there was no improvement from it, while G. Victor Heiser

obtained an apparent cure from the treatment when combined with oil of camphor and resorcin.

There are, of course, many reasons which may account for these very divergent results, but I suggest that one of the chief may be that the disease is attacked by different observers whilst in different stages. The initial lesions of the disease are unknown, and the knowledge of its subsequent stages very imperfect, and it may be that a substance has the power of destroying the bacilli and arresting the disease in certain stages and is quite useless at other periods. Therefore, until our knowledge of the pathology of leprosy is more accurate, given any case which does not improve after fair trial under one kind of treatment, I think that other forms which have been found of benefit in other hands should be tried, and careful notes made of the stage in which the disease is. This would, I think, lead to greater uniformity of results.

In conclusion, I should like to draw attention to the history of fish-eating in all the preceding cases. Certain kinds of fish locally known as "taragassa" and "kulumi"—the latter not unlike the cat-fish in appearance—are, according to popular report, the cause of leprosy if eaten frequently. The native method of dealing with fish is to open them, clean them roughly, and then dry them for about 12 hours in smoke; they are then sold in the market. It is probable, however, that the drying is very imperfect, and rotting fish would, I think, be a better description of the product as sold than dried fish; but the native view is that it is not every kind of fish that can convey leprosy, but only certain kinds, especially the two mentioned above.

Leprosy is extremely prevalent in Sokoto province, and has been for many years, and the natives have of course evolved this fish-eating theory without any knowledge of European views. I mention it accordingly as affording an interesting and independent piece of information in support of the theory of the late Sir Jonathan Hutchinson regarding the cause of leprosy.

Sokoto, Northern Nigeria.

SOME CASES OF TRAUMATIC HYSTERIA.

By J. W. GEARY GRANT, F.R.C.S. ENG.

For some years past whilst examining workmen's compensation and other cases for insurance companies I have happened to run across a considerable number of cases of definite traumatic hysteria (as distinguished from the ubiquitous traumatic neurasthenia); cases which I believe Janet calls "hysterical accidents"—that is to say, local manifestations of the hysterical state, determined by some fortuitous event, focusing the mind on the particular part concerned. Whilst of the cases I have examined these have furnished quite a considerable proportion, they can hardly be very common, for on inquiring among one or two medical men, who exclusively examine for large collieries, they appear to have had few or none. The following cases selected from many similar ones illustrate different types of this "hysterical accident." Since I put them together I have had the advantage of reading the Lumleian lectures by Dr. J. A. Ormerod,¹ and I think that these cases which I am fully aware are very short and

¹ THE LANCET, April 25th (p. 1163), and May 2nd (p. 1233) and 9th (p. 1299), 1914.

incomplete, may be of interest in connexion with his lucid exposition of the modern theories of hysteria.

The cases of anæsthesia have been far the most common, but I have only given one, for they all have nearly the same features; usually the stocking, glove, hind-quarter, or fore-quarter type. In most of them I have remarked what Dr. Ormerod terms Janet's sign, answering "No" when the anæsthetic part is touched. I have also found that in hemi-anæsthesia the vibrations of a tuning-fork are not felt when on the supposed anæsthetic side of the sternum and are at once appreciated when brought half an inch or so to the other side. Similarly with hysterical deafness, when the tuning-fork is passed to and fro across the middle line of the forehead it is heard or not heard according as it is or is not on the supposed deaf side.

As Dr. Ormerod points out, the anæsthetic areas are ideational and represent the anatomical ideas of the man in the street, who very fortunately for the examiner has no knowledge as yet of Head's segmental areas: but in some cases after repeated examinations with needles and suggestions as to where they feel and where they do not, the subjects make more or less good shots at peripheral nerve distributions, such as those of the median and ulnar.

In connexion with anæsthesia the absence of the plantar reflex in many cases is remarkable, and I think that attention is called to this point in Professor T. R. Glynn's Bradshaw lecture,² and the inference is drawn that anæsthesia cannot be purely ideational in hysteria. The normal plantar reflex is, however, a cortical reflex and its absence might therefore be explained by some block in the higher centres. Further, limbs the subject of hysterical anæsthesia, contractures, or paralysis show, as might be expected, some interference with their nutrition; they are cold, they do not bleed readily when pricked, and the electrical response of the muscles, whilst normal qualitatively, is not so quantitatively; by which I mean that a stronger faradic current is often required to obtain contraction than is the case on the sound side. A frequent association I have found is an absent plantar reflex with a considerably increased knee-jerk, the corresponding reflexes on the sound side being normal.

A further point to which I should like to call attention is the question whether some cases do not start from a small patch of organic anæsthesia due to injury of a peripheral nerve. In Mr. James Sherren's "Injuries and Diseases of Nerves" he cites one or two such cases where an extensive hysterical anæsthesia depended on a lesion of a small cutaneous nerve, in one case the anterior tibial, this being both verified and cured by the operation. In several of my cases this has occurred to me as possibly having been the starting-point, but I have never had the opportunity of verifying it by operation. It does seem feasible that a patch of anæsthesia in a hysterical subject may by suggestion determine anæsthesia of a limb. In one case, a brachial monoplegia, the history of the accident suggested injury to descending cutaneous branches of the cervical plexus; in another wasting of the vasti after a blow on the thigh, followed at a much later date by anæsthesia of the whole limb, suggested damage to a motor branch of the anterior crural, and possibly to cutaneous branches also; and in the first case of the following there may

have been, as in Mr. Sherren's, damage to the anterior tibial nerve.

In connexion with contractures I have never had the opportunity of examining them when the patient was asleep or under an anæsthetic, and I was very interested to find that in the cases mentioned by Dr. Ormerod the contracture persisted during sleep.

My greatest difficulty in these cases has been to determine whether they were genuine hysterical subjects or malingerers. I frankly confess, in spite of the statements of some authorities that this is a simple matter, that I usually find it quite beyond my powers to give my reasons why I regard one case as that of a malingerer and the other as that of a hysterical subject. Possibly a very elaborate examination, with determination of the visual fields and search for all the known stigmata of the condition, might corroborate, but in the cases where a claim under the Workmen's Compensation Acts or at Common Law is pending so many other factors enter in that the difficulty is extreme. Hysteria in many cases is associated with a large element of moral perversity, and as the result of a not inconsiderable experience I feel convinced that some at least have started by deliberate feigning and exaggeration and have at last reached a stage where they are quite unable to control the condition, whether a paralysis or contracture, much as a habitual liar in time is said to believe his own tales.

Further, it seems to me that to go so far as to say that "hysteria is as real as small-pox or cancer"³ is somewhat misleading. Any mental state is, of course, a reality *per se*, but taking these cases attributed to accident it is obvious that the hysterical condition is, in reality, precedent to the accident which merely determines the local manifestation of the underlying condition. Most of the following cases also arose as the result of very minor injuries, where there was little evidence of severe nervous shock; the accident itself acted by auto-suggestion, the auto-suggestions being strongly reinforced by injudicious treatment and the pernicious suggestion of stupid, ignorant, or designing relatives, the object in many instances being to magnify a condition in order to enhance its financial value.

Very different are the cases which Page and Thorburn relate, where there have been all the elements of profound psychic shock, railway accidents, explosions, and such occurrences. Into their pathology, so clearly explained by Thorburn, I need not enter, but in them it is evident that the accident truly is the *causa causans* of the resulting condition. By its profound effects on the nervous system it really is the cause of the resulting neurasthenia or hysteria, whichever may eventuate; but looking at the matter from what I conceive to be a commonsense point of view, it appears to me that to mulct a defendant in heavy damages for an imaginary condition in a plaintiff is not justice. Because a man after an abrasion of the hand allows himself to imagine that the injured member is useless, it is nothing less than an absurdity, I hold, to suggest that the person who has been held to have caused that simple injury should be also responsible for the mental state into which the injured person allows himself to fall. Moreover, the end result for the man, whatever sum he may obtain, is disastrous, for not with impunity do these people permit themselves to drift into such states.

² THE LANCET, Nov. 8th, 1913, p. 1303.

³ Purves Stewart: Lectures on Nervous Disease.

The consulting medical officer to a certain company, after reading my report on one case, annotated it as follows: "This man requires not massage or electrical treatment but a horse-whip," and I am not sure that this somewhat drastic, if impracticable, treatment might not have been more effective in causing speedy recovery than the £150 he received in settlement after two years of loafing. Without, however, venturing to suggest this as an appropriate stimulus to supply the lacking will-power, I only desire in citing these cases to draw attention to their medico-legal relations and to what I look on as two distinct classes in regard to causation, firstly, those due to severe mental shock; and secondly, those due to trivial causes, injuries occurring in persons with a strong neuropathic tendency and often in their genesis depending on wilful exaggeration of symptoms, or deliberate feigning.

Anæsthesia of lower limb.—The patient was a collier aged 34. In May, 1906, a stone fell on the dorsum of his right foot, which was bruised. He was laid up for a few weeks, and early in the case some loss of sensation was found on the dorsum (character and extent of this not specified). Later he was treated in an out-patient department with high frequency, &c., but he got worse and was only able to walk with the help of a stick. When I examined him 18 months later he presented loss of all forms of sensation, epicritic and protopathic, to a line drawn circularly round the knee. Calf an inch less in circumference than the left. Knee- and ankle-jerks brisk. No ankle clonus. No plantar reflex. Muscles of leg reacted to faradism, but required a much stronger current than left to produce the same result. Complained of "agonising pain" in foot and leg, and desired to have it amputated, but in spite of this he had a most serene and self-satisfied expression. Asked to stand without support his right leg gave way, and he gently and carefully subsided to the ground. This case resulted in prolonged litigation, and was the subject of two appeals. When examined two years later anæsthesia had spread to fold of groin and nates. Disposed of for a large lump sum.

Hysterical contracture.—The patient was a domestic servant. There was a small septic sinus on the back of the left hand between the fourth and fifth metacarpals, the result of operation for removal of a needle, and unconnected with bone, tendon or nerve. I laid it open and scraped it and it healed. Some weeks later she found she could not flex the fingers. Any attempt at passive flexion produced spasmodic contraction of extensors, but taking each finger separately and applying gentle pressure they could be passively flexed to full extent into palm, the extensors relaxing. No sensory loss. An arbitration followed with a decision in her favour. She passed from one medical man to another and to different hospitals. At one I believe she had an anæsthetic and the hand was fixed with the fingers bent, but about a year later her condition was the same, and as far as one could judge she never bent her fingers; she was then lost sight of.

Paralysis without sensory loss.—The patient, a collier's boy, in October, 1905, was kicked in the hollow behind the left great trochanter by a horse. Examined in September, 1910. Has done no work since. Walks with stick, and without it leg gives way and he gently subsides to ground. Lying down limb when passively raised drops as soon as support removed. No sensory loss. No ankle clonus. Right knee-jerk normal; left exaggerated. Right great toe response flexor; left absent. All muscles of limb react perfectly to weak current. A prolonged attempt at treatment with electricity and massage at hospital did no good, and he was disposed of for a large sum.

Hysterical monoplegia with anæsthesia.—The patient, a chemical plumber aged 41, met with an accident in February, 1911. When bending a leaden pipe, 12 feet long, over his right shoulder he felt a jar in his shoulder and pain running down the upper outer side of the arm. He worked for a week. He was examined by various medical men and reported to have ruptured the brachial plexus. When examined in the following November the arm was kept helplessly by the side, but with help could be raised above the level of the shoulder. All the muscles of the limb,

including the trapezius, rhomboids, and spinati, reacted readily to a faradic current of the same strength as required by those of the left arm. Light touch, prick, minor degrees of temperature and vibration sense loss over whole limb, limited by posterior border of scapula behind and midline of sternum in front, fore-quarter type. Pressure sense retained. Report: hysterical paralysis. After a course of electrical treatment, which resulted in no improvement, compensation was stopped, and the patient then obtained an appointment abroad, receiving a small sum in settlement.

A second case was that of a male, aged 40, who in February scraped the back of his right hand on a bin causing a simple abrasion, which healed in three weeks. He then lost all power in the hand and found weakness going up the arm and affecting the right leg. When examined at the end of April he was a robust, well-developed man, with his right hand hanging limply by the side. He made no attempt at grasping. The hand was colder than the other. Sensation of light touch and prick lost to a line drawn circularly round the elbow. The muscles gave a perfectly normal response to faradism. Right knee-jerk exaggerated, no ankle clonus, plantar reflex flexor. His claim was settled at once for a small sum, and he was not seen again.

Hysteria with spasmodic movements.—The patient on July 6th, 1910, when standing on top of a lift put his foot through a glass roof. No physical injury whatever. His health was said to have gradually broken down since then, but he did not cease work until February, 1911. He was examined in May. As he entered the room he screamed and started the most extraordinary writhing movements of the hands, arms, and face. These movements, which resembled those of posthemiplegic athetosis, consisted of slow flexion of wrist, pronation of forearm, flexion of forearm, fingers in accoucheur's position, and hands gradually passed behind the back and then to position of rest. The shoulders were raised and head bent over until they touched, and the mouth was drawn over to the side. Both sides were affected; speech was jerky and staccato. His aspect indicated terror of examination. Knee-jerks appeared grossly exaggerated, but this proved to be a purely voluntary act, as shown by the long reaction period and the fact that tapping the patella or tibia caused the jerk. On diverting his attention found to be normal. No sensory loss. Pupils normal. No apparent lack of intelligence. He remembered dates perfectly and discussed his case intelligently with me. He was sent by his medical attendant to University College Hospital under Dr. J. S. Risien Russell, who got, I am informed, his house physician to write to the solicitors of the insurance company pointing out the boy's condition, that he was improving, but that this was delayed by impending litigation, and suggesting a compromise on the basis of a small sum and a promise of work when certified cured. On my advice this was agreed to, and for a trifling sum and this guarantee, which was faithfully carried out, the case was settled, and the boy is now well and back at work.

Hysteria with astasia abasia and fits.—The patient, aged 31, a labourer underground, in January, 1903, was caught in the right calf by the hitching plate of a runaway tram. "Ruptured some sinews." He was in bed for two months and on crutches for two years. He was in hospital for a month and was treated with faradism. There was loss of power in the right leg, right arm, and right side of the face. Latterly he had developed fits with aphasia, coming on at night. He fell out of bed and rolled underneath. His face was drawn to the right. He lost speech for from four to nine days, during which period he had to express his wants on a slate. He could not see or hear so well on the right side. When seen on April 18th he was a powerfully built man of fine muscular development. The circumference of the lower limbs was approximately equal. On walking across the room he reeled to the left, but stood with eyes shut perfectly steadily. When he sat or stood on the sound leg he could move the affected one in any direction, and held it straight out in front of his body, a condition known as astasia abasia. Superficial reflexes were brisk and Babinski sign was not present; a spurious clonus on right side and greatly exaggerated jerk on tapping bone or tendon obviously like the clonus voluntary. It was found normal on distracting attention. On testing muscles with faradism the left was normal; right, cried out with a weak current;

electrode disconnected without his knowledge with battery working, cried out loudly with the pain. His medical attendant assured me he was absolutely genuine, and thought it was a case of cerebellar disease, though how a blow on the calf could cause this he did not try to explain.

If the account the last patient and his friends gave is true it is a curious one, as he would appear to have had fits of pure motor aphasia of subcortical type. In these he was unable to speak a word for days and expressed his wants in writing, being able to understand written and spoken speech and to write. Anatomically this would be due to a block of the outgoing fibres whilst mental speech processes and memory of words is unimpaired, and is a variety which is known to occur in hysteria and an illustration of which is the scriptural one of Zacharias, the father of John the Baptist.⁴ It is noteworthy that the paralysis was on the correct side, the right, and the patient's symptoms therefore fitted a hypothetical organic lesion. The view I took of him was that he was a hysterical malingerer.

Cardiff.

MENTAL SYMPTOMS ASSOCIATED WITH EXOPHTHALMIC GOITRE.

By W. LEGGETT, B.A., M.D. DUB.,

SENIOR ASSISTANT PHYSICIAN, ROYAL ASYLUM, MONTROSE.

THE following case appears to be of interest on account of the well-marked mental symptoms.

The patient is an unmarried female, aged 34 years. Having been certified as insane, she came under my care for treatment. She was described as suffering from hallucinations of hearing and as having the delusional idea that her relations and neighbours were conspiring to poison her food and give her chloroform. It is her first attack, and the mental symptoms are said to have existed for two weeks before certification. Her father was insane and committed suicide. Most of the physical signs usually seen in cases of this kind are present. The thyroid gland is moderately and symmetrically enlarged, a thrill is easily felt by the hand, and a murmur heard by the stethoscope over the gland. No dyspnoea or cough is noticeable. The pulsation of the carotid arteries is most remarkable, as is also the apex beat of the heart, for as a result of the latter a wave of pulsation may be seen all over the chest and abdominal walls. The pulse-rate is 142 per minute and tends to become more frequent while the patient is under observation. There is a well-marked fine tremor of the muscles of the face and limbs, and also a staring expression of the eyes. The eyeballs are prominent, but this sign is not well advanced at present. Neither Stellwag's nor von Graefe's sign has developed. The pupil seems to be unaffected, but on looking downwards both the upper eyelids enter into a state of spasmodic twitchings. The patient is thin, anæmic, and becomes very breathless on slight exertion. The superficial reflexes are increased.

The chief mental features noticed are irritability, restlessness, and a tendency to resent any interference by those about her. She appears to be very preoccupied and often sits or lies in bed with her eyes closed apparently listening to imaginary voices. She rarely makes an attempt to initiate a conversation, but when spoken to she smiles or answers in monosyllables and sometimes incoherently. She does not ask questions or try to

continue a conversation. Her manner is rather dignified and commanding, yet she is nervous and tremulous, and becomes more so under observation. Her method of speaking is rather abrupt and somewhat halting. She always says she feels well. She mistakes the nurses and doctor for people she has known at her home and gives them names accordingly. Periodically she jumps impulsively out of bed and takes up a threatening or abusive and sometimes violent attitude towards those who happen to be near to her. On one occasion she struck a fellow patient for throwing dirt on her, whereas the patient was merely dusting the floor. Certain noises which occur about the room she mistakes for guns which are fired to shoot her. She apparently does not realise where she is at present, nor has she any idea as to the length of time she has been with us. Her general condition varies considerably from time to time; sometimes she is comparatively composed and almost polite, but usually she is excitable, exalted, and rather noisy and difficult to deal with.

The most prominent mental features seem to be irritability, impulsiveness, suspiciousness, disorientation, incoherence of speech, exaltation, and a tendency to take up a threatening and abusive attitude. Drugs or other treatment have had no appreciable effect on the condition.

Montrose.

A CASE OF FIBROSIS OF LUNG TREATED WITH BACILLUS FRIEDLÄNDER VACCINE.

By ALEXANDER SANDISON, M.B., B.C. CANTAB.,
B.Sc. LOND.,

TUBERCULOSIS OFFICER TO THE COUNTY BOROUGH OF CROYDON.

THE patient, a married man aged 33, first came under my notice on Nov. 8th, 1913. He gave a history of having suffered from left-sided pleurisy in 1907 and severe bronchitis in 1910. Except for these illnesses he had enjoyed good general health and was in regular employment as a clerk. In January, 1913, he had a severe cold, with night sweats, and noticed streaks of blood in his sputum. A doctor who attended him diagnosed "phthisis and heart trouble" and ordered a change. Tubercle bacilli were not found in the sputum. He gave up work in consequence and went down to Devonshire, where he gained weight and improved in general health, returning to Croydon in October, shortly before I saw him first, but was nevertheless unable to work on account of dyspnoea and palpitation. My notes on examining him on Nov. 8th were as follows:—

Family history: Negative, except for a sister who died from acute phthisis about 1900, aged 29. Previous diseases: None beyond those described above. Symptoms: Cough and sputum moderate, but troublesome; sputum not foul and never copious. No night sweats nor hæmoptysis. Digestion good and larynx normal. Chief trouble excessive dyspnoea, combined with cardiac palpitation on any exertion. Weight 10 st. 5 lb. Temperature morning 97° F., evening 98°; pulse 96.

Physical signs:—Left lung: Percussion note dull all over front and back. Movement diminished and chest flattened in front. Loud broncho-vesicular breathing with increased vocal resonance. Numerous coarse râles and rhonchi heard all over. Right lung: Impaired note, broncho-vesicular breathing, and coarse râles over the upper two-thirds of the upper lobe in front and behind. Heart: Apex beat displaced upwards and outwards towards the left axilla, about an inch outside the nipple line. Occasional systolic bruit in the

⁴ Purves Stewart: loc. cit.