The Journal of the American Medical Association

Published Under the Auspices of the Board of Trustees

VOL. LXVII, No. 6

CHICAGO, ILLINOIS

AUGUST 5, 1916

ADENOMYOMA OF THE RECTOVAGINAL SEPTUM*

THOMAS S. CULLEN, M.D.
BALTIMORE

In 1914, I referred¹ to two cases of adenomyoma of the rectovaginal septum recently reported by Cuthbert Lockyer,² and mentioned observations made by Macnaughton-Jones, Clifford White and Herbert Spencer. I also briefly mentioned two cases that Dr. D. S. D. Jessup of New York had observed and which he was about to publish.

Since then Jessup³ has reported his cases in full. I have reported another case.⁴ Stevens⁵ reports five cases of adenomyoma that he observed in the course of eighteen months. He also refers to a case which he had accurately described in 1910.

Pierre Nadal,⁶ of Bordeaux, reported a case of adenomyoma of the posterior vaginal wall in 1911. Other cases may have been recorded, but a careful survey of the literature in the *Index Medicus* for the last years failed to yield me more data on this subject.

I shall briefly refer to two other cases that have come under my personal observation and shall then try to sketch the course of adenomyoma of the rectovaginal septum as evidenced by the cases thus far reported. I shall also endeavor to indicate the appropriate methods of treating such tumors.

CASE 4.—Adenomyoma of the rectovaginal septum extending out to the right pelvic wall and constricting some of the pelvic nerves; hysterectomy; recovery.

Dec. 28, 1914, Dr. John Calvin Huckins of Ashland, N. H., wrote me as follows: "In THE JOURNAL OF THE AMERICAN MEDICAL ASSOCIATION under date of March 14, 1914, I saw a report of a few cases of adenomyoma of the rectovaginal septum. On May 10, 1913, I had a patient with a small tumor in the rectovaginal septum. The mucous membrane of the right vaginal fornix had a bluish color and the mucosa was puckered. High up in the right vaginal fornix a small hard mass could be felt. Upon rectal examination, there was some bulging of the anterior rectal wall, and the mass could be made out more distinctly by a finger in the rectum. What the mass was I do not know. I had two surgeons in

consultation, neither of whom knew what the growth was, or where it had originated. Dr. A. H. Harriman and I did an exploratory operation through the posterior vaginal wall and removed two thirds of the tumor, but we were of the opinion that it was malignant and did not complete the operation. When I read your article ten months later, I then knew what this tumor must have been. I took pains to keep track of this woman and for over a year I could not detect any lump upon digital examination. Only a few days ago, on rectal examination, was I able to find a small lump which seemed to be nearly as large as the one found at the time of our exploratory operation (about the size of a small hen's egg)."

Under date of Dec. 31, 1914, Dr. Huckins says: "This case seems to me to be identical with Lockyer's Case 2, only not quite as marked. I am of the opinion that the growth is connected with the uterus and broad ligament."

The patient, Mrs. O. P. S., aged 43, was admitted to the Johns Hopkins Hospital, Jan. 18, 1915. She had had no children. Her chief complaint was pelvic pain. The present illness had begun about three years before. At that time she had been for a long ride in a buggy over a rough country road. On her return home she had been taken with severe pain in the lower part of the pelvis and a feeling of pressure in the rectum. These symptoms had persisted for a few days and then disappeared. The following month, just after the menstrual flow had ceased, she had again experienced this peculiar grinding pain in the lower part of the pelvis. It had lasted for four or five days and then suddenly had disappeared. This pain had continued each month after the cessation of the menstrual flow. The discomfort would last from four to five days, and then until the following month she would be perfectly well. She said that the character of these pains was unusual. They were difficult to describe. They were more like a dull aching or grinding which, although not severe, was very annoying and when present made her very nervous. The pain was most severe at night and seemed more pronounced when she was on her feet. It was usually localized in one spot, but at times would radiate down the right leg.

The menstrual history was unimportant except for the fact that on two occasions during the past summer the flow had been profuse and prolonged to from eight to ten days. The bowels had been somewhat constipated; there had been no pain on defecation, and no history of bloody stools at any time.

On pelvic examination under anesthesia I found the uterus normal in size, slightly retroverted. No thickening could be made out on either side.

In the right vaginal vault and almost glued down to the lateral pelvic wall was a hard indurated mass, apparently about 3 cm. long and 2 cm. broad (Fig. 1). It felt distinctly nodular, and reminded one somewhat of an adenomyoma. I made a median incision and did a complete hysterectomy, but saved both tubes and ovaries. On the left side I kept close to the cervix. On the right side we dissected out the ureter and, after removal of the uterus, separated the rectum from the vagina for a distance of several centimeters, and then cleared the right broad ligament. The mass in question was about 3 cm. long and densely adherent to the side of the rectum, to the posterior vaginal wall, and also to the lateral

* From the Gynecological Department of the Johns Hopkins Hospital and of the Johns Hopkins University.

* Read before the Section on Obstetrics, Gynecology and Abdominal Surgery, at the Sixty-Seventh Annual Session of the American Medical Association, Detroit, June, 1916.

1. Cullen, Thomas S.: Adenomyoma of the Rectovaginal Septum, THE JOURNAL A. M. A., March 14, 1914, p. 835.

2. Lockyer, Cuthbert: Proc. Roy. Soc. Med., 1913, vi, No. 4.

3. Jessup, D. S. D.: Adenomyoma of the Rectovaginal Septum, THE JOURNAL A. M. A., Aug. 1, 1914, p. 385.

4. Cullen, T. S.: Surg., Gynec. and Obst., March, 1915, p. 263.

5. Stevens, T. G.: Adenomyoma of the Rectovaginal Septum, Proc. Roy. Soc. Med., 1916, ix, Obst. and Gynec. Section, p. 1.

6. Nadal, Pierre: Bull. de l'Assn. franç. pour l'étude du cancer, 1911, iv, 338.

wall of the pelvis. It really had to be dug out. When loosened, it was still found attached to the rectum. It was removed. The pelvic peritoneum was brought together; a cigaret drain was laid in the pelvis and carried out through the vagina. The abdomen was closed. The patient made a perfectly satisfactory recovery.

Gyn.-Path. No. 20828: Sections from the nodule show that it is composed of bundles of nonstriated muscle, of fibrous tissue, and of a moderate amount of adipose tissue. Scattered throughout the tumor is uterine mucosa. Sometimes glands can be seen lying in direct contact with the muscle. At other points two or three glands are embedded in a characteristic stroma (Figure 2). Here and there is a slight tendency toward the formation of a miniature uterine cavity. The glands are lined with one layer of cylindric epithelium. The stroma is identical with that usually found surrounding the glands of the endometrium. Here and there the stroma is rarefied. At other points it shows a moderate degree of hemorrhage. Scattered here and there throughout the tumor are brownish or yellowish-brown masses of pigment, the remains of old hemorrhage. The outlying portions of the tumor are particularly instructive. The adipose tissue of the broad ligament is being gradually replaced first of all by small round cells, the fat cells looking like vacuoles in a mass of small round cells. In other places the tissue is in part or in its entirety replaced by connective tissue and here and there, as indicated in Figure 3, muscle and connective tissues are enveloping the nerves, thus in a measure accounting for the pain that the patient has had on the right side and which had extended down the right leg. The tumor is an adenomyoma of the type usually found in the rectovaginal septum, and in parts cannot be differentiated from an adenomyoma of the uterine wall.

CASE 5.—*Adenomyoma of the rectovaginal septum.*

Miss J., aged 38, was referred to me by Dr. Alfred Whitehead, Jan. 8, 1910. For several months she had been complaining of pain in the lower abdomen, chiefly in the region of the rectum. For three months she had been growing weaker and for the last three weeks had been confined to bed. Her periods were excessive.

On pelvic examination I found, behind the uterus, a somewhat irregular mass, apparently 5 or 6 cm. in diameter. The thickening could be much more easily outlined through the rectum than through the vagina. Her pain at the menstrual period was so severe that I advised her removal to the Johns Hopkins Hospital.

Operation was performed Jan. 10, 1910. On opening the abdomen I found a myoma, 1 cm. in diameter, on the left side of the fundus. This was removed. The mass very low down behind the uterus seemed to be due to an inflammatory thickening between the cervix posteriorly and the rectum (Figure 4). The rectum at this point was drawn up and adherent to the back of the cervix. After the cervix had been separated from the rectum, the area of thickening was found to extend backward and to implicate part of the left uterosacral ligament. Although the surface of the rectum was thickened over an area 2 by 3 cm., the lumen of the bowel was in no way encroached on.

To have removed satisfactorily the area of thickening in the rectum would have necessitated, as a preliminary, a complete hysterectomy. This we were very anxious to avoid, as the patient was engaged, and hoped soon to be married. As the growth in no way suggested cancer or tuberculosis, we laid a drain in the pelvis, bringing it out through the vagina in such a manner that it separated the raw area of the bowel from the raw area on the back of the cervix.

Fig. 1 (Case 4).—Adenomyoma of the rectovaginal septum. The black area posterior to the cervix indicates the location of the growth noted in Case 4. It had encroached on the right vaginal vault and at operation was found to have extended as far as the pelvic wall on the right. To this it was densely adherent.

The patient made a good recovery. I saw her again at her home, Feb. 28, 1910. At the menstrual period following her dismissal from the hospital she had terrific rectal pain. This persisted, and I again examined her under anesthesia and found the supposed inflammatory thickening in the vaginal vault just as marked as before the abdominal operation. I opened up the old drainage tract in the vaginal vault and found that the area of thickening in the bowel had folded over on itself. After straightening out the fold I again drained through the vagina. We thought the kinking of the rectum might possibly have caused the severe pain. Complete removal of the uterus followed by removal of the rectal thickening was again considered but, on account of the contemplated marriage, it was thought wiser to continue palliative measures for a time at least. During my absence, in the summer of 1910, she came under the care of Dr. E. H. Richardson, and to him I am indebted for the further clinical notes on the case. As her symptoms still persisted, in October, 1910, Dr. Richardson dilated and curetted, removing a large amount of endometrium. After this the patient improved for several months, although she was never free from menstrual pains, which were referred to the left hip and to the left iliac fossa. She was practically an invalid

for three weeks out of every month. Nov. 1, 1911, Dr. Richardson did a supravaginal hysterectomy, releasing extensive pelvic adhesions, removing at the same time the left tube and ovary, and resecting the right ovary. Enough of the uterus was left to allow the patient to menstruate. Then the operator encountered dense pelvic adhesions. The cervix was firmly fixed to the rectum, and a mass, several centimeters in diameter, was found at this point. The patient's condition was such that resection of the rectum could not be undertaken. Her convalescence was a stormy one and she was afterward kept on a rest cure for four weeks. She improved markedly so far as her general condition was concerned, but from the first menstrual period after operation she complained of the same pain as before; it was, however, less severe. Finally, she again became an invalid for three weeks out of each month. Her condition remained about the same.

In July, 1912, the patient was suffering so intensely and so constantly that another thorough pelvic examination was made. At this time there was a marked increase in the size of the "inflammatory mass" in the rectovaginal septum. She was admitted to the Johns Hopkins Hospital Aug. 7, 1912. At that time she was poorly nourished. The abdomen was opened, and numerous omental adhesions were found everywhere. After dissecting out the right ovary, the stump of the cervix was reached with a great deal of difficulty. The rectum was now gradually dissected free and the "inflammatory mass" was exposed. This, Dr. Richardson tells me, was situated about 4 cm. from the anus; it implicated the anterior surface of the bowel and extended out laterally. The mass was about the size of a small hen's egg. The portion of the rectum containing the tumor was resected and the bowel drawn down and attached to the anal margin. The patient stood the early part of the operation very well, but near the close her pulse became almost imperceptible. She did not rally and died ten hours later from surgical shock.

Gyn.-Path. No. 18756: Sections from the tumor show that it is composed of nonstriated muscle and fibrous tissue, with islands of uterine mucosa scattered freely throughout it (Figure 5). Some of these islands contain only one or two glands; in other sections there are eight or ten. The glands occasionally lie in direct contact with the muscle, but nearly all of them are embedded in stroma. The gland epithelium

is thicker than usual and corresponds to the type that we speak of as hyperplasia, being two or three times the normal thickness. The stroma surrounding the glands is also exceptionally abundant. A good many of the glands are dilated and some are evidently cystic. Even in the dilated glands the epithelium is much thicker than usual. The picture is that of a typical adenomyoma. The glands in the myoma encroach more on the rectal wall than usual (Figure 6).

Fig. 2 (Case 4).—Adenomyoma of the rectovaginal septum. In this case the growth had invaded the right broad ligament, and extended out to the pelvic wall, so that it was necessary literally to dig the growth away from the side of the pelvis. In the lower part of the section two or three glands are seen. The dark areas surrounding them are the characteristic stroma of the uterine mucosa. In the left portion of the section it will be noted that the adipose tissue is being replaced by fibrous tissue and muscle. The condition was diagnosed clinically by Dr. Huckins, the family physician.

In a few places they have penetrated the muscle and at one point at least they have broken through the muscle and lie in direct contact with the rectal glands (Figure 7). One gland is seen projecting down nearly half way through the rectal mucosa. Such an intimate blending of uterine mucosa with rectal mucous membrane I have never seen before. To have shelled this growth out or to have cut it away without removing a segment of the rectum would have been an utter impossibility.

DESCRIPTION OF ADENOMYOMAS OF THE RECTO-VAGINAL SEPTUM

Myomas occurring in the septum between the vagina and the rectum have long been known, and some of them have reached such a size that they encroach markedly both on the vagina and on the rectum. In the male, so-called rectal myomas have been noted. In one case the tumor was as large as a coconut and at first was thought to be an inoperable malignant growth. The short but comprehensive article on the subject by F. Descoedres⁷ is well worth a thorough study.

From the cases of adenomyoma of the rectovaginal septum that I have been able to collect, the following tentative classification may be made:

1. Small adenomyomas lying relatively free in the rectovaginal septum (Cullen's Case, 1, Stevens' Case 1, Stevens' Case 5, Nadal's case).

2. Adenomyomas adherent to the posterior surface of the cervix and at the same time to the anterior surface of the rectum (Lockyer's Case 2, Cullen's Case 3, Stevens' Cases 2, 3, 4 and 6, Jessup's Cases 1 and 2).

3. Adenomyomas gluing the cervix and rectum together and spreading out into one or both broad ligaments (Cullen's Case 4, Cullen's and Richardson's Case 5).

4. Adenomyomas involving the posterior surface of the cervix, the rectum and broad ligaments, and forming a dense pelvic mass that cannot be liberated (Cullen's Case 2).

Of course, one group merges imperceptibly into another, and a case which today belongs to Group 1 may in a few years belong to Group 2 or to Group 3.

On vaginal examination one can detect a definite thickening. This may be situated in the posterior wall to the right or left of the cervix, but is usually directly behind it. It may vary from 1 to 3 or 4 cm. in breadth. The thickening is occasionally nodular. As a rule, however, it is diffuse, hard, and reminds one of dense inflammatory tissue or of a small adherent myoma. In some cases the nodule moves with the cervix, as if the two made up a single piece. In other cases the nodule also seems firmly fixed to the rectum.

In a few cases the vaginal mucosa over the nodule has been puckered and is of a bluish tinge. This was noted in Lockyer's Case 2, and in my Case 4. In Stevens' Case 2 the overlying vaginal mucosa was also puckered. The puckering is due to the tumor becoming adherent to the vaginal mucosa.

In those cases in which the growth has invaded all parts of the pelvic floor, one finds induration throughout the vaginal vault. There may also be a slightly cystic feel on bimanual examination, if some of the glands or a miniature uterine cavity in the tumor have filled up with blood. This was observed in my Case 2.

On rectal examination, these tumors, if situated directly behind the cervix, can often be more clearly felt than through the vagina. If the growth be adherent to the rectum, the bowel over this area is splinted and hard, and does not yield. The rectal mucosa, however, even over the growth, is, as a rule, perfectly normal, although the tumor may project slightly into the lumen of the bowel. In my Case 3 the growth had partly encircled the bowel and had produced a considerable degree of narrowing of the lumen. In

Fig. 3 (Case 4).—Muscular and fibrous tissue in an adenomyoma of the broad ligament encircling and compressing nerves. This section is from the right broad ligament. It shows the diffuse myomatous and fibrous tissue surrounding nerves. There were definite symptoms of nerve pressure.

Jessup's Case 2 the patient had had hemorrhoids for thirteen years, and in my Case 1, in which a discrete myoma, 4 cm. in diameter, was encroaching on the lumen of the bowel, the rectal mucosa at this point showed several small polypi projecting from it. These Dr. Samuel T. Earle removed prior to my doing the hysterectomy.

7. Descoedres, F.: Des myomes du rectum, *Rev. de gynéc. et de chir. abd.*, 1910, xiv, 5.

HISTOLOGIC APPEARANCES

These growths, whether large or small, discrete or diffuse, consist of nonstriped muscle and fibrous tissue, and have scattered throughout them typical uterine glands. These glands may occur singly and lie in direct contact with the muscle, or may be found in groups. As a rule, they are separated from the muscle or fibrous tissue by the characteristic stroma of the uterine mucosa, and at many points one can see islands of stroma devoid of glands. In some of the glands fresh blood may be found. More frequently, yellowish-brown pigment is noted in clumps scattered throughout the stroma. This pigment is the remnant of old bleeding.

Along the advancing margin of the diffuse growth, the manner in which the growth extends can sometimes be very clearly followed. This was well shown in my Case 4 (Fig. 2). Here the young connective tissue cells and the muscle fibers gradually replace the adipose tissue of the broad ligament, and finally engulf it completely. They encircle nerves and gradually compress them, thus adding to the discomfort of the patient.

The most widespread growth that I know of was found in my Case 2. In this case a hysterectomy had been performed in San Francisco two years before. When I saw the patient, the lower portion of the pelvis was filled with an adenomyomatous growth. The mass in the left side proved to be a miniature uterine cavity filled with chocolate colored fluid—the residue of the retained menstrual flow.

Histologically these growths are identical with adenomyomas of the uterus.

SYMPTOMS

Age Incidence.—The ages were as follows, 25, 30, 32, 35, 37, 37, 38, 41, 41, 41, 42, 43 and 53. The 53 year old patient still menstruated slightly. She told me that members of her family did not reach the menopause until somewhat late, and that her great uncle was born when his mother was 54. All of the cases occurred among women who were still menstruating.

Menstruation.—The most pronounced symptom was profuse menstruation. In some this was painless, in others accompanied by great discomfort. My patient, in Case 2, was almost exsanguinated, although a supravaginal hysterectomy had been done for removal of a myoma two years before. In my Case 3 the patient had insufferable pain for two days before and after each period. In my Case 4, in which we found the adenomyoma gripping small pelvic nerves on the right, the pain at the period was always on the right side; it also extended to the right hip. Partial removal of the growth relieved her of these pains for from six to eight months. In my Case 5 the pain at the period was very severe, and the patient was finally incapacitated for three out of the four weeks.

Rectal pain was also a prominent feature in some cases. In Lockyer's Case 2 there was great pain in the rectum, and defecation was difficult. The patient said

that something prevented the bowel from moving. The discomfort was more severe at night. In my Case 3, the patient had had for the last two months six bowel movements daily. The discomfort was so severe that she insisted on having something done. It was in this case that we found much narrowing of the lumen of the bowel. In my Case 4 the woman was slightly constipated. Before her first operation defecation had been invariably accompanied by pain. In my Case 5 there was pain in the lower abdomen, chiefly in the rectum.

When the growth is small, is not adherent to the rectum and has not spread out into the broad ligament so that it has commenced to exert pressure, we shall expect few if any symptoms. In those cases, however, in which the adenomyoma has become adherent to the rectum, there is a tendency to pain in the rectum, with or without painful defecation.

When the growth has invaded the broad ligament and encroached on the nerves, we may expect discomfort referable to this region, due in the first place to the pressure on the nerves, and in the second place to the swelling of the tumor at the menstrual period. Profuse periods seem to be the rule.

Fig. 4 (Case 5).—Adenomyoma of the rectovaginal septum. The black area between the cervix and rectum indicates the location and approximate size of the supposed inflammatory thickening noted at operation in Case 5. A reference to the description of the specimen shows that the mass was a typical adenomyoma. In the fundus of the uterus was a small myoma.

CAUSE OF ADENOMYOMA OF THE RECTOVAGINAL SEPTUM

Our knowledge of the distribution of adenomyoma is becoming more stable each year. Adenomyomas of the uterus, formerly thought to be rare, are relatively common.⁸ The source of origin of their glands is also clear, namely, from the uterine mucosa. We know that adenomyomas may be found in the round ligament,⁹ in small umbilical tumors,¹⁰ in the utero-ovarian ligament,¹¹ and recently one has been described as occurring in the broad ligament over the left ureter.¹² My colleague, Dr. William W. Russell,¹³ has found uterine mucosa in the hilum of the ovary.

When discussing cervical adenomyomas, in my work on "Adenomyoma of the Uterus," several years ago, I pictured an island of perfectly normal mucosa of the body

of the uterus situated near the outer surface of the cervix.¹⁴

Subsequent studies may possibly show that this is no great rarity. If so, this growth might readily become the starting point for those adenomyomas found in the rectovaginal septum.

Krönig, Hartz and Breus have reported cases that may be of interest in this connection.

8. Cullen, Thomas S.: Adenomyoma of the Uterus, Philadelphia, W. B. Saunders Company, 1908.

9. Cullen, Thomas S.: Adenomyoma of the Round Ligament, Johns Hopkins Hospital Bull., May and June, 1896.

10. Cullen, Thomas S.: Diseases of the Umbilicus, Philadelphia, W. B. Saunders Company, 1916, p. 373.

11. Cullen, Thomas S.: Adenomyoma of the Uterus, Figure 41, p. 142.

12. Stickney, G. L.: A Case of Diffuse Adenomyoma of the Uterus with Discrete Adenomyoma over the Left Ureter, Johns Hopkins Hospital Bull., 1915, xxvi, 304.

13. Russell, William W.: Aberrant Portions of the Müllerian Duct Found in an Ovary, Johns Hopkins Hosp. Bull., 1899, x, 8.

14. Cullen, Thomas S.: Adenomyoma of the Uterus, Figure 49, p. 169.

Krönig¹⁵ reports a very interesting instance of a cystic adenomyoma springing from the posterior wall of the uterus and extending backward beneath the peritoneum of Douglas' pouch. It consisted of one large thin-walled cyst containing a liter of brownish-red fluid, and of a more solid portion consisting of about thirty small spaces so arranged that they

Fig. 5 (Case 5).—Adenomyoma of the rectovaginal septum. The matrix of the tumor consists of nonstriated muscle and of fibrous tissue. The round colony of glands on the right is composed of typical uterine glands. They are surrounded by the characteristic stroma of the mucosa. The large glands in the center and also those on the left have a thicker epithelium than usual. They look exactly like those noted in hyperplasia of the endometrium. They also are partially embedded in stroma similar to that of the endometrium. (Photomicrograph by Herman Schapiro.)

resembled a honeycomb. The walls of the large cyst, especially of that portion lying free in the abdominal cavity, were very thin, in places measuring scarcely more than 1 mm. The cyst walls were composed of fibrous tissue and of a varying amount of muscle. The inner surface was in places lined with cylindric ciliated epithelium. The more solid portion of the tumor was a typical cystic adenomyoma, which, as Krönig says, in the form and arrangement of glands and in the cystic spaces, corresponded in practically all points with the adenomyomas of von Recklinghausen. There were definite groups of glands surrounded by the characteristic stroma. Krönig thinks that the tumor originated in the uterine wall and was later pushed out into the connective tissue of Douglas' pouch.

Hartz¹⁶ observed a similar case in Sängers clinic.

In Breus' ¹⁷ Case 2 the patient was 51 years of age. There was a tumor, the size of a child's head, springing from the posterior surface of the uterus and covered by the peritoneum of Douglas' sac and the left broad ligament. The tumor, on its upper and posterior surface, was hard. On section it was seen to be composed of myomatous tissue, but the central portion contained several cavities. These varied from a pea to an apple in size, and were in part separated from one another by thick partitions. Several of them, however, communicated with one another. The cysts had smooth inner surfaces and were filled with a thick chocolate-brown fluid. The largest cyst com-

municated directly with the uterine cavity by a funnel-shaped opening just above the internal os. The cysts were lined with cylindric ciliated epithelium, and, where the large cyst communicated with the uterine cavity, the surface epithelium of the uterine mucosa was directly continuous with that of the cyst. Breus considered the tumor as a subperitoneal and intraligamentary cystic myoma of the uterus.

The glands in the adenomyomas of the rectovaginal septum look like, and act exactly like, those of the mucosa of the body of the uterus, and they undoubtedly arise from uterine glands or from remnants of Müller's duct.

DIFFERENTIAL DIAGNOSIS

As pointed out by Jessup, in both of his cases, in which the patients were operated on by Dr. William S. Bainbridge, other surgeons had diagnosed the growths as inoperable cancer. That was permissible in the past, but will not be in the future. These growths occur in women, are noted in the childbearing period, are usually situated directly behind the cervix and, as a rule, are firmly adherent to the rectum. The symptoms are excessive and painful periods, accompanied in severe cases by much rectal pain. On rectal examination the growth is found literally plastered to the anterior surface of the bowel. The rectal wall is fixed at that point, but its mucosa is intact, and is normal, or merely shows a few hemorrhoids or possibly a polyp. I know of nothing in pelvic or rectal surgery that should be confounded with this condition in future.

TREATMENT

Some might argue that simple removal of the appendages would cause atrophy of the uterine mucosa contained in the adenomyomas of the rectovaginal septum. My Case 2 is a sufficient answer. Although

Fig. 6 (Case 5).—Adenomyoma of the rectovaginal septum. The rectal mucosa is normal. Some of the glands of the adenomyoma are small and round; others are large. The large glands have a thickened epithelium and correspond to the characteristic picture noted in hyperplasia of the endometrium. Nearly all of the glands are surrounded by the characteristic stroma of the uterine mucosa. At the right limit of the section one of the uterine glands is lying in direct contact with the rectal mucosa. (Photomicrograph by Herman Schapiro.)

a supravaginal hysterectomy had been performed two years before for a myomatous uterus, the pelvic condition had grown steadily worse.

1. Where small discrete nodules exist in the posterior vaginal vault, these may be readily removed

15. Krönig, B.: Ein retroperitoneal gelegenes voluminöses Polycystom entstanden aus Resten des Wolff'schen Körpers, Beitr. z. Geburtsh. u. Gynäk., 1901, iv, 61.

16. Hartz, A.: Neuere Arbeiten über die mesonephrischen Geschwülste, Monatsschr. f. Geburtsh. u. Gynäk., xiii.

17. Breus, Carl: Ueber wahre epithelführende Cystenbildung in Uterusmyomen, Leipzig, 1894.

through a vaginal incision, as was so successfully done by Stevens.

2. Where the growth occupies the posterior surface of the cervix and extends laterally, after the ureters have been dissected out carefully, a complete abdominal hysterectomy should be performed.

3. If the growth be firmly adherent to the rectum, a wedge of the rectum should be removed, together with the uterus. It has been found best, after freeing the uterus on all sides, to open up the vagina anteriorly and laterally. The uterus and the rectum can then be lifted farther out of the pelvis, thus facilitating the removal of the necessary wedge of the anterior rectal wall. The uterus really acts as a handle, and the necessary rectal tissue and the uterus are removed as one piece.

4. Where the lumen of the bowel is greatly narrowed, a complete segment of the rectum should be removed with the uterus, and an anastomosis should be made.

5. In desperate cases, where everything in the pelvis is glued together, as in my Case 2, an ideal operation is out of the question. The patient will not stand a long operation, and, if she could, a satisfactory result could not be obtained. In such a case it would be better to cut across the sigmoid, invert the lower end, close it, and bring the upper end out through the abdominal wall of the left iliac fossa, making a permanent colostomy. When the patient has to some extent regained her strength, the uterus, the lower portion of the rectum and the broad ligament tissue can be shelled out as one piece.

These growths, while histologically not malignant, remind one of glue. Unless they are completely removed, further trouble is liable to occur.

The fact that Stevens has observed five cases of adenomyoma in eighteen months, and that I have had five cases in my own practice, leads me to believe that adenomyoma of the rectovaginal septum is no great rarity and that the literature of the near future will contain records of many such cases.

20 East Eager Street.

ABSTRACT OF DISCUSSION

DR. CAREY CULBERTSON, Chicago: At first thought one might be led to surmise that adenomyoma of the rectovaginal septum is a pathologic entity so rare as to be of little value. But when we realize that Cuthbert Lochyer's first case was reported as recently as 1913 and that already we have these five cases of Dr. Cullen's, with the six additional ones of Stevens', it can only mean that in all clinics of any considerable size these cases have been overlooked. In addition to his own cases, the first of which, by the way, Stevens claims

he reported as far back as 1909, cases have been described in England by Spencer, Lietch, Gough and Stewart and Bland-Sutton. A. Mayer last November reported five cases of a peculiar tumor-forming inflammatory process in the tissues between the rectum and vagina which he places in the same category as R. Meyer's "posterior paravaginitis and parametritis with proliferation of epithelium," described in 1908. Aman recently described a case which he calls "retrocervical fibro-adenomatous serositis" and refers to five others, accepting Meyer's pathologic explanation and conclusions. An interesting point in the pathology of growths is that, with the exception of two of Stevens' cases, all were associated with inflammatory processes, the uterus usually being retroverted and densely adherent. This suggests at once the similar condition in the tubes, called by Robinowitz "adenomyosalpingitis," but which has also been regarded as a true neoplasm. It is not so difficult to understand how inflammatory pressure might produce such a squeezing out of the mucosa into the wall of the tube; but in the case of the rectovaginal septum or postcervical space, the distance from the uterine cavity is too great to admit of the presence here of typical uterine mucosa as the result of inflammatory pressure. Lack of time prevented Dr. Cullen from expressing his opinion as to the origin of these growths—whether he regards

them as due to inflammatory pressure on the uterus, which seems to be the German idea, or as true neoplasms originating from embryonic rests persisting from the fusion of the müllerian ducts.

DR. THOMAS S. CULLEN, Baltimore: I have had in all five cases of adenomyoma of the rectovaginal septum. When the first case came under my care I was ignorant of the fact that such a condition as adenomyoma of the rectovaginal septum ever existed, and it was only within the last five weeks, when looking over old specimens, that we found out the exact nature of the case. In our cases of adenomyoma of the rectovaginal septum there has been no indication whatever of inflammation, that is, of the pouring out of small round cells or polymorphonuclear leukocytes. There is, however, an inherent tendency for the mucosa and for the nonstriated muscle and connective tissue to spread out and invade surrounding structures. The glands in

Fig. 7 (Case 5).—Glands of adenomyoma of the rectovaginal septum penetrating the rectal mucosa. The rectal mucosa looks normal. In the underlying muscle are a few uterine glands (a) lying in direct contact with the muscle. At b is a uterine gland extending down into the rectal mucosa. The continuity of this gland could be followed in other sections. This is the only instance known to us in which the glands of an adenomyoma have actually invaded the rectal mucosa. (Photomicrograph by Herman Schapiro.)

these cases undoubtedly originate from the uterine mucosa or from remains of Müller's duct. I feel confident that within the next few months more cases of adenomyoma of the rectovaginal septum will be reported.

Tapeworm in Portugal.—The *Medicina Contemporanea* states that the eggs of the *Hymenolepis nana*, the smallest tapeworm known, were found in thirty-three of 503 children at Lisbon whose stools were examined. From one to several hundred ova were found, and in nineteen cases they were associated with other helminths' ova. This is a proportion of 6.5 per cent.; Grassi found this parasite in the stools of 10 per cent. of the children in southern Italy that were examined, and it seems to be prevalent in Belgian miners and in Italian immigrants in Brazil. In the Lisbon cases, there did not seem to be any special symptoms from the presence of the parasite.