- Bordley, James, Jr.: Optic Nerve Disturbances in Diseases of the Posterior Nasal Sinuses. *Jour. A. M. A.*, Vol. lxxv, p. 809, Sept. 18, 1920.
- Knapp, Arnold: *Medical Ophthalmology* (An International System of Ophthalmic Practice, Pyle), 1918, p. 382.
- Vail, D. T.: Monocular Retrobulbar Optic Neuritis from Hyperplasia of the Ethmoid Bone. *Trans. Am. Acad. Oph. and Oto-Laryng.*, Oct. 29, 30, 1917, Aug. 5, 6, 1918, pp. 431-443.
- Loeb, H. W.: Two cases of Blindness Relieved by Ethmoid Excision. *Trans. Am. Laryng., Rhinol., and Otol. Soc.*, Vol. xxvi, p. 346, 1920.
- Loeb and Wiener: The Borderland of Oto-Laryngology and Ophthalmology. *Trans. Am. Laryng., Rhinol., and Otol. Soc.*, Vol. xxvi, p. 136, 1920.
- Skillern, R. H.: The Ethmoidal Problem. *The Laryngoscope*, Vol. xxx, p. 687, November, 1920.
- Present-Day Value of Surgical Procedure of the Sphenoidal Sinus. *Trans. Section Laryngol., Otol., and Rhin., Amer. Med. Assn.*, p. 99, 1916.
- Hajek: Zur Diagnose und intranasalen chirurgischen Behandlung der Eiterungen der Keilbeinhöhle. *Arch. f. Laryng.*, 1904, xiv, 105.
- Roemer, Paul: *Text Book of Ophthalmology*. Translated by M. L. Foster, Vol. iii, p. 712, 1913.
- de Schweinitz George E.: *Diseases of the Eye*, 8th edition, p. 475.

A RARE FRACTURE OF THE LOWER END OF THE HUMERUS.

BY LUTHER G. PAUL, M.D., F.A.C.S., BOSTON.

No mention of the fracture described below has been found, after a moderately thorough search of the literature.

A girl, 14 years of age, was referred to me by Dr. M. F. Cronin of Boston, on April 30, 1919. The day before, she had fallen down several steps while carrying a younger child.

The left elbow was swollen and tender over the external part, and motion was much restricted on account of pain. There was a slight bruise of the skin just above and back of the head of the radius. She had about twenty degrees of flexion, and about the same degree of extension, without pain. The rotation was about one half normal. Ether was given, and after manipulating the part, I was unable to make a diagnosis of the condition. The bony landmarks were in normal relation, and there was no abnormal mobility or crepitus noticed. The arm was put in a comfortable position, which was nearly right angle flexion, and on the next day, an x-ray was taken.

This showed a vertical fracture of the lower end of the humerus, in which the anterior part of the articulating surface had been separated from the rest of the bone, and that this fragment had been forced out of the joint, and was lying on the anterior surface of the lower end of the humerus, fully an inch above its normal position.

Under ether, an unsuccessful attempt was made to reduce the fracture, and on May 9, 1919, she was again etherized, and an incision into the joint was made on the outer side of the elbow. The fragment was found as the x-ray

BEFORE REDUCTION.

AFTER OPERATION.

showed. It was freely movable, and its only point of attachment was by its inner end to a small area of the capsular ligament. The fragment was so loose that it could be almost wholly delivered from the wound, and with retraction of the anterior soft parts, could be plainly seen. It was seen to consist of the capitellum and trochlear surfaces in one piece.

All attempts to replace the fragment were without success, until the following method was tried. The forefinger was placed in the wound over the fragment, and between the fragment and the tendons of the biceps and brachialis anticus muscles, the elbow being flexed to somewhat beyond a right angle. The assistant then extended the forearm, and at the same time exerted a pull to increase the space between the humerus and the upper ends of the radius and ulna. The pressure exerted by the tightened muscles on the finger of the wound was surprisingly powerful, and when to this was added pressure on the outside, the fragment slipped into place.

The arm was put in acute flexion, and the capsule closed with No. 1 chromic catgut, and the skin with silkworm gut. The arm was held in acute flexion with adhesive plaster. There was considerable swelling in twenty-four hours, so that it was necessary to extend the arm a few degrees in order to relieve the circulation.

The wound healed well, the stitches being removed on the ninth day. The adhesive plaster was omitted at the end of three weeks, and at that time, extension was twenty degrees less than a right angle, and flexion was normal. An x-ray at this time showed the fragment in its normal relation. She was given massage and passive motion, and instructed to carry moderate weights, such as a school bag with books in it. The amount of extension gradually increased, and on September 24, 1919, the extension was fifty degrees beyond a right angle, and normal flexion. Three months later, the movements of the joint were normal. The accompanying x-ray photographs show the fracture before and after reduction, and need no detailed description.

RECONSTRUCTION OF EARS

BY HILBERT F. DAY, M.D., F.A.C.S., BOSTON.

It is rare that we are asked to multiply ears, but that was the proposition that was put up to me.

I was asked to replace the lost portion of an ear and by plastic surgery I succeeded in making two perfect ears from one and a half ears.

On the afternoon of January 30, 1920, I had referred to me by Dr. Henry J. Perry, a young man, D. B. C., 22 years old, who stated that twelve days before he had injured one of his ears by glass in a taxicab accident. The upper portion of his ear was cut immediately off in a straight line which sloped downwards and backwards, as is shown in Figure A.

On examination, it was found that the cut edge was nearly healed and various methods of procedure were considered which would close the defect. The possibilities of flaps, either from the head or some other portion of the body, were considered. To my mind, none of these schemes would do more than cover in