LOCAL ANESTHESIA IN CONJUNCTION WITH GENERAL ANESTHESIA IN MASTOID OPERATIONS.*

By T. W. MOORE, M.D.
Huntington, W. Va.

Every otologist has had patients suffer intensely when recovering from the anesthetic after a mastoid operation. Sometimes it is an intolerable aching, at other times a burning or smarting, and again the patient will say "the awful soreness in my neck."

That all that can be done to avoid or lessen the post-operative pain and make our patients comfortable when coming out from the general anesthetic is our imperative duty here as in the other fields of surgery, I think is conceded by all. So if by spending a few additional minutes in preparation our patient, after the general anesthetic passes off, goes into a quiet and restful sleep for from one to three hours, much has been accomplished.

I have purposely omitted any reference to that complex and little understood physiological or biological condition known as shock which is not, as a rule, a marked concomitant of mastoid surgery.

Nor am I by any means convinced that anociassociation as advocated by Crile is theoretically correct or has any practical application as such.

I have only tried this method in five cases, but in all of these the results have been most gratifying. The technique I have employed is as follows:

An hour before the patient goes on the operating table he is given one-sixth of a grain of morphine sulphate and one hundred and fiftieth of a grain of atropine sulphate, hypodermically. After the patient is well under the ether anesthesia, the skin is infiltrated with a solution of which is given by the open drop method, novocaine one-half per cent and epinephrin chloride one-fiftieth per cent. It requires but a small quantity to accomplish this. Then a deep injection, sub-periosteal if possible, is given of urea and quinine hydrochloride 1 per cent; after waiting five

minutes the operation is proceeded with in the usual manner.

This in no way modifies or effects the healing of the wound. I prefer the blood clot dressing and have found healing equally as prompt and satisfactory as when local anesthesia has not been used.

LATERAL SINUS THROMBOSIS.*

By J. H. BARNES, M.D.
Enid, Okla.

Lateral sinus thrombosis is not so common as we may think when we observe the position of the sinus to the middle ear and the mastoid cells. But when it does occur it is imperative that we be able to make a diagnosis early in the life of our patient.

I have nothing new to offer on the subject, but wish to point out some of the points of diagnosis and to report a case that was diagnosed as lateral sinus thrombosis, also as a severe middle ear infection and mastoiditis which I am sure now was neither, but a brain abscess with local meningitis, though there were no tests made to confirm the diagnosis.

ANATOMY.

The lateral sinus is in the lower posterior region of the skull in a depression from the torcular herophili forward to the jugular foramen. It passes close to the mastoid cells just before it terminates at the jugular foramen as the internal jugular vein.

ANATOMY OF LATERAL SINUS.

Its position to the ear and mastoid cells is so variable that we are at a loss just where the sinus is in the mastoid.

On examining twenty skulls in one series the sinus was not in the same position in any two skulls. It passed in one-eighth inch to external auditory canal to an inch or more behind it. It is sometimes close to the cortex or may be an inch or more from the surface.

The mastoid cells are at times surrounding the sinus. All these conditions make it in the road to danger when the middle

*Read in Eye, Ear, Nose and Throat Section, Southern Medical Association, Ninth Annual Meeting, Dallas, Texas, Nov. 8-11, 1915.

*Read in Eye, Ear, Nose and Throat Section, Southern Medical Association, Ninth Annual Meeting, Dallas, Texas, Nov. 8-11, 1915.

ear is suppurating and does not have free drainage.

The lateral sinus is joined by the superior petrosal above and the inferior petrosal below the mastoid.

The cause of lateral sinus disease comes directly from involvement of the mastoid cells, which is secondary to middle ear suppuration. All severe suppurative middle ear disease involves more or less the mastoid cells.

The sinus is more apt to be diseased from an acute exacerbation of an old mastoid disease. There is a history of chronic discharge from the middle ear, or it may be a frequent recurrence of the suppurative ear. It is very seldom that an acute mastoiditis will attack the sinus.

We find that the diseases of childhood are the forerunners of these ear and mastoid diseases such as measles, scarlet fever, diphtheria and influenza and pneumococcus infections.

Any of these infections are very apt to affect the ears if the child has adenoids.

It has been noted that the ears are seldom affected when the child's throat is healthy or the adenoids have been removed early.

The symptoms of lateral sinus diseases are: Discharging ear of a chronic nature with involvement of mastoid, usually an acute attack of an old affection.

The mastoid is tender on pressure and more or less headache. The symptoms that are diagnostic are a severe chill with high fever, which is 103 or 104 while the chill is still on, and then rises to 105 or 106 in a short time, followed by sweat and a lower temperature, even down to normal. Then in a few hours or a day or two we have recurrence of the chill with the same symptoms. The chills are very irregular as to time, but the character of the chill, fever and sweats are the same in each attack. The patient may feel quite well between the attacks.

If the sinus becomes involved low down next to the jugular foramen, the vein below becomes tender and may appear as a hard cord in the path of the jugular vein just under the anterior border of the sterno-cleido-mastoid muscle.

The pathology begins as a phlebitis. There is a deposit thrown out on the inside of the vessel which becomes larger till soon a clot is formed which fills the

sinus and the circulation is cut off. If this clot is infected and the germs are thrown into the circulation we have the symptoms as given above, those of severe septicemia. The clot may be sterile and the phlebitis be so walled off as to protect the circulation. Then we do not have the severe symptoms and the patient may get well without surgical interference.

The complications are brain abscess, meningitis and an extension of the infected thrombosis to the other sinuses. There may be metastatic abscesses and pneumonia.

The treatment is surgical so as to get good drainage. The mastoid cells should be cleaned out thoroughly and the sinus exposed. Before it is opened the jugular should be tied off below so as to protect the general circulation.

Then open the sinus and clean out the clot from above and below until we get free bleeding.

If we do not get free bleeding from below we may be sure the clot extends as far as the inferior petrosal sinus; then we should dissect all the jugular from above our tie-up as far as possible.

After all sources of infection have been removed we pack the wound tightly and bandage. Dress the wound on the third or fifth day, depending on condition of the patient.

If there should be a recurrence of the symptoms then we should open the wound and look for further infection.

If the operation is done early before the system is too much saturated with the infection we may get a hasty recovery.

I will now give you a report of my case that I first diagnosed lateral sinus disease, not from the typical symptoms, but from the sudden rise of temperature. Then, dropping in a few hours to nearly normal.

CASE REPORT.

Boy age 14, father physician, in February, 1915, was taken with what appeared to be acute otitis media. He was sick for a few days before with some disturbance with stomach; he had fever and some aching of limbs as if he had la grippe. He was given a dose of Pluto water and in a few minutes vomited and burst his right ear drum, which was called to the mother's attention by hemorrhage from the external ear.

I was called and cleaned the ear and found the ear drum ruptured. He had had no pain up to this time in the ear. In twenty-four hours pus began to discharge very profusely, which was from the pneumococcus.

The fever continued from 102 to 104, which was thought to be partly from his disturbance in the alimentary canal by his attending physician, who was seeing him every day with me.

He had had for the past four years repeated attacks of discharging ear, which lasted from a week to a month. I saw him a month or two before this attack and the ear was dry and the drum membrane healed, so diagnosed this attack as acute suppurative otitis media. The drainage was good in this attack, but after three or four days had severe headache all over the head. There was no tenderness over mastoid cells and no pain in ear.

His fever and pain continued high for a little more than a week, when it began to subside. The discharge became gradually better and in two weeks was apparently well and started back to school. His ear appeared to have quit discharging. Two weeks afterward he came to my office, saying that his ear was closed up. I cleaned out some foul, thick pus and found the ear drum had not closed and directed him to return for further treatment.

Two days after I was called to the house to see him. His ear was discharging very freely a thick yellow pus, which was pneumococcus variety of the streptococcus form. He had high fever and complaining of his stomach as before. The fever continued to rise from 102 to 105 in the typhoid fever curve. The ear discharged freely with no pain for two days, when he began to complain of headache all over the head. The pain got worse and soon he was unable to move his head without severe pain. There was no tenderness over the mastoid or any one point.

About the fifth day the ear ceased to discharge the ear drum bulged. We called a consultation and it was agreed to incise the ear drum. We opened from above and below into a large opening in the ear drum. It was at this time that we agreed that the mastoid should be opened in twelve hours if the ear did not discharge or the pain, which was very severe, did not subside.

The next morning after we had incised the drum we called the second consultation, as all the symptoms were worse and the ear dry. The temperature up to this time had run a typhoid course till it reached 105; then it suddenly dropped to 101 and in a few hours was 105 again.

The second consultant thought we had a case of mastoiditis and suggested the third consultant, who arrived that night just twenty-four hours after we had freely incised the ear drum. The drum was still bulging and the last consultant thought that his high fever, which had gone up and down three times in twenty-four hours, was due to retained secretions in the middle ear, and suggested a mastoid operation if we did not find pus by reopening the drum membrane.

During the day the physician in charge made a diagnosis of lateral sinus thrombosis, though there was never a chill nor the least signs of one. The temperature was that of a phlebitis where the germs were thrown into the blood, septiceimia. He did not look septic at any time. His face was a good color and he had no sweats up to this time. It was suggested that his blood should be examined, but this was not done, as we felt sure the opening of the mastoid and lateral sinus would relieve him.

We opened the mastoid immediately after the arrival of the third consultant, as we had everything ready for the operation. After the patient was put to sleep we opened the ear drum; there was no pus, so we opened the mastoid. There was only a thin cortex over what had been mastoid cells. We opened into a large cavity which communicated with the mastoid antrum and found the lateral sinus a reasonable distance behind the ear and exposed on posterior wall of the cavity and pus coming up from the lower posterior part of the cavity from around the lateral sinus. The lower exposed part of the sinus was flat, making a depression down below the cavity filled with pus. The sinus was closed for about one-half an inch above the compressed sinus.

We ligated and cut the jugular below and then opened the compression and took out a hard and comparatively healthy clot. The vein bled freely from above and below, so we packed the wound tightly and bandaged.

This was done just one week from the time I was called to see him.

The symptoms continued as before with pain and high fever; sometimes in the morning it was lower than in the afternoon. Then it would run to 105½ very quickly. He began to have sweats and breathe very rapidly, 40 per minute.

The wound was dressed the third day, looked good; no discharge of pus from the wound or ear.

He still got worse; began to show signs of weakness on the fifth day after the operation and the R. ext. rectus became paralyzed and the pu-

pil did not react to light. This was the only sign of paralysis. He died on the eighth day after the operation.

This was evidently a case of chronic mastoid disease with probably an involvement of the sinus and at this time had found its way into the brain in a region which produced but few symptoms other than the severe pain and high fever which went to 107 just before death.

There was nothing that we did that seemed in the least to help him. The sinus was thoroughly cleaned out and we felt sure that he would improve, and if it had been a mastoiditis or lateral sinus thrombosis he would have shown some signs of improvement. So we make a diagnosis by exclusion.

A spinal puncture should have been made for pus and leucocytes. A post mortem is the only means of a definite diagnosis.

MASTOIDITIS WITH LATERAL SINUS THROMBOSIS AND CEREBELLAR ABSCESS—OPERATION, RE- COVERY.*

By JOHN H. FOSTER, A.M., M.D.
Houston, Texas

The report of a single case before a scientific medical body requires in general an apology or an explanation. The one to which your attention is invited today, however, has some very unusual features, sufficiently so, it seems, to justify recording.

The patient, P. R. H., male, aged 25, came under our observation on March 20, 1914. He gave a history of trouble with the left ear for ten days. The membrana tympani had been incised and he had done fairly well until two days previously, when the pain again became severe and some swelling had developed about the ear. Examination showed that he was suffering from furunculosis of the canal of the ear as a complication of the middle ear infection. He was moved to the hospital, and under gas anesthesia the canal was incised and pus evacuated from two or three places. The middle ear discharge was slight.

During the following few days other infected points in the canal developed and were opened; the middle ear discharge ceased, the membrane healed, and at the end of a week he left the hospital apparently well.

On April 2d, five days later, patient appeared at the office complaining of great pain in the left ear again. His temperature was 103° F., and examination revealed the drum red and bulging. It was incised freely and a bloody discharge liberated, and he was sent back to the hospital. His temperature on admission to the hospital was 104° F., and he complained of great pain in the ear. Hot irrigations were ordered with cold sponges to reduce the temperature. Within twelve hours his temperature dropped to 101° F. and he was more comfortable, though never absolutely free from pain, which seemed to him to be deep within the ear itself.

At noon of the fifth, after slight chilly sensation, his temperature rose to 104.2° F. Blood count was made on the fourth day, showing 12,000 whites, 80% polys. On the sixth day patient first complained of pain outside the ear and localized it in the left frontal region, describing it as a severe, boring pain. There was no mastoid tenderness nor sagging of the postero-superior canal wall. On the afternoon of this day a distinct chill was recorded with a rise of temperature to 104.2° F. again. Examination of the discharge from the ear showed the presence of numerous Klebs-Loeffler bacilli. Blood count showed 18,000 whites, 76% polys. X-ray plates showed cloudiness of all the mastoid cells. Ten thousand units of diphtheria antitoxine were administered and patient's mother in Georgia notified of his condition, and of the probable necessity of operation.

The next day he seemed much better. His temperature rose no higher than 101.3° F. and he was decidedly more comfortable. The following morning, however, the temperature rose to 103° F. and it was deemed best to go into the mastoid despite the lack of tenderness. Accordingly, on April 8th, the mastoid operation was performed. All of the cells were found involved, filled with pus and granulation tissue. The wall of the lateral sinus was exposed and found somewhat discolored but not obstructed, and was not incised, but the bone was removed over it for a distance of two inches, including the knee, so that it could be easily opened later if necessary, and the wound was then dressed.

The patient was much depressed when he left the table and had a hard chill in a short time. He rallied, however, and was more comfortable soon.

*Read in Eye, Ear, Nose and Throat Section, Southern Medical Association, Ninth Annual Meeting, Dallas, Texas, Nov. 8-11, 1915.