mic goitre, how could it change well-marked cases of Graves' disease into typical cases of myxedema?

What except the action of the ray could cause the thickening, adhesion and sclerosis of the capsule and trabeculae of the gland which every surgeon claims occurs in all cases of Graves' disease that have had pre-operative Roentgen ray treatments?

Valvular Lesions in the Aged. Graham E. Henson, Jacksonville, Fla. The Journal of the Florida Medical Association, Vol. III., No. 8, February, 1917, p. 225.

Valvular lesions of some sort exist in the great majority of the aged. In a great many instances they cause no subjective symptoms; in fact, are not likely to so long as a compensatory hypertrophy exists. It should be borne in mind that the detection of a most pronounced murmur or murmurs, accompanied by even great hypertrophy of the cardiac muscle and displacement of the apex beat, is not necessarily a factor in the ailment that has caused the aged patient to seek consultation and advice. Having detected a valvular lesion in an aged person, even though at the time it may not be a factor in an existing illness, it should not be ignored; and although at the time it may not call for active therapeutic measures, prophylactic measures for maintaining the existing compensation should be carefully considered.

In considering the treatment of valvular lesions in the aged, it is especially imperative that to a large degree a waiting, watchful policy should be assumed in so far as active therapeutic measures are concerned.

Careful attention should be paid to the examination of the heart in the aged patients. Their span of life may be materially lengthened if the time be taken thoroughly to familiarize one's self with the condition of their cardiac function. The days of compensation may be prolonged by careful prophylactic measures, and when compensation is lost, life may still further be prolonged by the careful and intelligent administration of therapeutic aids. On the other hand, by slipshod diagnostic methods the days of compensation may be shortened and the crisis still further hastened by the administration of contraindicated drugs.

Radium Therapy. Frank E. Simpson, Chicago, Ill. The Radium Quarterly, Mallers Building, Chicago, Vol. I., No. 1, January, 1917, p. 1.

In this article the author strikes a conservative note in his remarks upon the relation of radium to the cure of cancer. Cancer of the skin is usually curable by radium, but cancer of the deeper structures may lead frequently to death by the development of secondary deposits, although the primary growth can often be relieved. Interesting remarks are made especially with reference to burying or implantation of radium needles in malignant growths, a method which promises much in the treatment of cancer of the tongue. The prophylactic use of radium in preventing the recurrence of malignant dis-

ease after operation is commended. Cancer of various organs of the body is briefly mentioned. In carcinoma of the breast encouraging facts have been observed, although only inoperable cancers or inoperable recurrences should be treated. In carcinoma of the uterus radium treatment in inoperable cases accomplishes results not to be obtained by any other method. In carcinoma of the bladder, esophagus, rectum and prostate, radium holds out some hope of relief and even of clinical cure in selected cases. Several cases of sarcoma are reported in which the results from radium treatment have been gratifying. Fibroid disease of the uterus may be relieved symptomatically and in some cases an apparent cure of the disease has been brought about. Splenic leukemia is briefly commented upon and the beneficial results from radium treatment are mentioned.

Value of Spinal Puncture in the Etiologic Diagnosis of Cardio-Vascular Diseases. Leon Bloch, Chicago, Ill. The Journal of the American Medical Association, Vol. LXVIII, No. 9, March 3, 1917, p. 691.

Spinal puncture as a means of determining the etiology of cardio-vascular diseases, where other etiologic factors aside from syphilis could be ruled out, was suggested to the writer by obtaining a positive Wassermann reaction in the spinal fluid of a patient with aortic aneurism in whom, in spite of symptoms suggestive of cerebro-spinal syphilis, the blood Wassermann was negative.

In a series consisting of five cases of aortic aneurism, one of abdominal aneurism, four of aortic regurgitation, nine of myocarditis in various stages of decompensation, two of hypertension and one of angina pectoris, seventeen gave positive Wassermann reactions in the spinal fluid. These comprised four cases of aortic and the one of abdominal aneurism, three of aortic regurgitation, eight of myocarditis, and the one case of angina pectoris. Five cases with negative Wassermann reaction in the spinal fluid yielded such findings as lymphocytosis, and the presence of globulin and albumin with or without high pressure.

Attention is called to the fact that there were no cases of double mitral lesion in this series, a condition which is generally accepted to be of inflammatory origin.

The important evidences in the spinal fluid which indicate syphilitic involvement or origin are a positive Wassermann, high cell count with a preponderance of lymphocytes, a positive Nonne and Noguchi, and a spinal fluid under high pressure. The Lange test was not carried out in this series. Tuberculosis can be easily ruled out in the differential diagnosis by the differences in the clinical course.

While a cure of syphilis of the cardio-vascular system is impossible where the condition is of long standing, temporary relief is sometimes obtainable. Hence, there is a justification for the procedure in those cases where the diagnosis is in doubt.

(Continued on page 560)