and mixed cases are the most difficult to diagnose. The complement-fixation test for gonorrhea in conjunction with the same test for syphilis may point out some of these complicated or mixed cases so that they can be correctly and fully diagnosed and receive the benefit of combined treatments.

DYSENTERIC CASES

Occasionally during the course of or following an attack of dysentery, presumably of bacillary origin, we see a sudden rise of temperature with an effusion into the bursae of the larger joints, accompanied by only slight pain. The patient recovers in a short time with nothing but symptomatic treatment. A knee fluid culture, kindly made by Dr. L. B. Bates in one of these cases several years ago, was sterile.

TUBERCULOUS CASES

The only case of primary tuberculous arthritis in this series was of the usual knee-joint variety, monoarticular. The Wassermann test was negative. The diagnosis was arrived at mainly by exclusion, the presence of continued fever, localized tenderness and exostoses about the epiphysis and head of the bone, and an x-ray plate kindly made by Dr. T. Barhart of this hospital. The x-ray is of the greatest value both in diagnosis and prognosis in all cases of arthritis showing bone destruction. An arrest of the process was obtained in this case by splinting and general rest, in conjunction with the usual dietetic and hygienic treatment of tuberculosis. There was partial ankylosis when the patient was discharged.

SUMMARY

Ninety-one per cent. of the acute arthritides of this series admitted to the male negro medical wards during the past year were caused by venereal disease. Sixty-three per cent. were of syphilitic etiology, the Wassermann test being positive in 93.6 per cent. of these cases. An intensive mixed treatment, carried to the point of maximum tolerance, has enabled these men to return to work temporarily in an average of 18.8 days.

Twenty-eight per cent. were of gonorrheal origin. Vaccine treatment gave very good, sometimes brilliant, results in the great majority of cases. Unfavorable results due to the treatment were not noticed. The average number of days under treatment per man was twenty-four.

All other types of arthritis were in such minority that they are practically negligible.

Acute articular rheumatism was not encountered, and acute or chronic endocarditis was not recognized in any of the total one hundred cases.

In conclusion, I wish to extend my thanks to the following gentlemen for permission to publish: Col. W. C. Gorgas, Medical Corps, U. S. A., chief sanitary officer; Col. Charles P. Mason, Medical Corps, U. S. A., superintendent of Ancon Hospital, and to Dr. W. E. Deeks, chief of the medical clinic, on whose service these cases occurred.

The Milk of Indian Buffaloes.—According to Consul Norton, at Bombay, India, investigations carried on during a period of fifteen months on a number of buffalo cows belonging to a dairy at Poona, show that the Indian buffalo surpasses the best breeds of milk cows in the production of butter fat, 7 to 8 per cent. being common, and 10 per cent. having been found. The milk is white, and the butter usually white. The percentage of proteins in buffalo milk is distinctly greater than in cow's milk. Milk-sugar and minerals correspond closely to those of cow's milk.—*Daily Consular Reports.*

RABIES

A PATHOGNOMONIC SIGN GENERALLY OVERLOOKED

M. B. WESSON, B.S., M.D.

EL PASO, TEXAS

Aside from the pioneer communications on hydrophobia, very little has been written on the subject, and that little is extremely vague and indefinite. Having an opportunity to observe a series of over two hundred rabbits inoculated with rabies (for use in the El Paso Pasteur Institute), as well as a number of animals infected with "street rabies," I was impressed with the uniformity of the development of the symptoms, and particularly of a left hemiplegia starting in the leg (simulating Landry's paralysis) of which I could find only vague references in the literature.

The first symptom of the disease in the rabbits was a general nervousness; the entire body seemed to be trembling and, if the animal was startled by the sudden clapping of hands, it would stagger and apparently have great trouble in maintaining its balance. Soon a limping of the left hind leg was noticed and, if the rabbit was taken from the cage and made to hurry across the floor, that leg gave way entirely. In a short time a weakness in the corresponding front leg developed, then in the neck, as in acute ascending paralysis. Hence when the animal attempted to walk it would move around the cage clockwise, supporting its weak left side against the wall. A few of the rabbits had convulsions, and when that happened they rolled over and over from right to left. If the rabbits are not molested they will remain crouching in a corner of the cage, and then the first apparent weakness is the left torticollis; but if the animal is made to move, the paralysis will be found to be greatest in the left hind leg. The rabbits lie fully extended on the left side as the paralysis progresses, apparently dead, the only sign of life for several hours before the end being an occasional movement of the muscles about the nose and eyes.

In man after the expiration of the incubative period, the patient complains of loss of appetite and headache; he becomes depressed, ill-tempered, apprehensive and excitable; then the breathing becomes difficult and labored, accompanied by spasmodic contractions of pharynx and larynx.

It was at this stage of the disease that I saw J. M., an albino Mexican, who had been bitten about the middle of September by his pet dog (which died the following day in a convulsion). He was sitting up in bed, as he could not breathe in a recumbent position. The stridulous breathing sounded like the bark of a dog, and the frequent shaking of his head emphasized the resemblance and caused the neighbors to make the correct diagnosis. He attempted to convince them that they were wrong by frequently demonstrating how easily he could drink water, but after one or two swallows he always became strangled; salivation was marked. He had a strong grip with his right hand, but that of his left was very weak. On attempting to walk he seemed to have perfect control over his right leg, but his left gave way under him and he had to be supported on that side. As he moved across the room his entire left side was weak, his head almost lying on his left shoulder, which in turn drooped. The following day hallucinations appeared, which caused the patient great anxiety and excitability. A little later paroxysms occurred which resembled maniacal attacks with delirium. They were of short duration, but large doses of morphin, chloral and potassium bromid had no effect in controlling them. Paralysis became general, and death took place suddenly on the fourth day, during a paroxysm, from asphyxia.

On October 5 a neighbor's dog, which had been bitten by that of the Mexican boy, became restless and ill-tempered. He bit the grocery boy (on the hand) and a young hog (on the nose) before he was frightened away by the owner. Three days later it was noticed that something was wrong with the dog's left hind leg. When he wanted to lie down he would turn round and round several times from right to left and then when he allowed his left thigh to touch the floor he would at once bound up with a sharp cry, holding the leg in the air and whining as if in pain. The following day the leg was completely paralyzed, and there was a weakness of the corresponding front leg and neck. The development of the paralysis, preceded by a left hemiplegia, and followed by death, was the same as seen in the rabbits. (The brain was sent to the local Pasteur institute and they reported the presence of Negri bodies.) The groceryman was immediately started on Pasteur treatment, and the bitten hog isolated. On November 12 the hog became restless and began to limp with the left hind leg; it was unable to drink, though it made frequent attempts and would put its head to the level of its eyes in a bucket of water; salivation became marked, and progressive paralysis and death, as seen in the rabbits, followed the next day.

An interesting point here is the fact that a "mad" fox-terrier at almost precisely the same time bit a man and a hog. The man was given Pasteur treatment and is well, while the hog received no treatment and died of rabies.

SUMMARY

- 1. These cases are reported to call attention to the characteristic development of the paralysis of rabies as noted in a series of animals, including man, dog, hog, guinea-pigs and rabbits.
- 2. The initial site of paralysis was the "left hind leg," and this in a short time developed into a left hemiplegia, the paralysis eventually becoming general.
- 3. In an animal that has suddenly become extremely nervous and irritable, the development of a weakness in the left hind leg, without any apparent reason, is sufficient to warrant a tentative diagnosis of rabies.

THE PREPARATION AND EMPLOYMENT, IN
A SERIES OF CASES, OF A POTENT
POLYVALENT ANTISTAPHYLO-
COCCIC SERUM *

B. A. THOMAS, A.M., M.D.
PHILADELPHIA

Impressed by the questionable propriety of the employment of active immunity with bacterins in the acute stage of an infection, whether localized or generalized, as in furuncle or septicemia—conditions in which, it seems logical to infer, the human organism is already sufficiently stimulated by viable bacteria to the production of specific antibodies, if not completely overwhelmed as in the latter instance, rendering the employment of an inactivated specific antigen inadvisable if not positively harmful—I attempted in September, 1909, the preparation of an antiserum which I hoped might prove useful in antagonizing the diverse lesions of that commonest of all pyogenic cocci, the *Micrococcus pyogenes aureus*.

Various attempts have been made on horses and other animals to produce a potent antistaphylococcic serum, but they have almost invariably resulted unsuccessfully, although Doyen and Palchikowsky assert that they have

succeeded partially. Emery states that "it is possible by special methods to produce a serum which has a slight bactericidal effect, but normal human serum or the serum of a patient who has been submitted to a course of antistaphylococcic vaccination is quite powerless in this respect." Schorer asserts that "the value of the serum . . . is inconsiderable, and its injection in the treatment of staphylococcus infections is seldom or never warranted." Such noted authorities as Ehrlich, Bordet and Citron omit even to mention, in their works on immunity, the existence of antistaphylococcic serum. It is therefore with considerable trepidation that I venture to assert the successful preparation of a potent polyvalent antistaphylococcic serum.

The experiments of Neisser and Wechsberg show the production of a typical hemolysin by the *Micrococcus albus* and *M. aureus* in culture growth. Following the subsequent discovery by these two serologists that human and certain other animal serums normally contain antistaphylolysin, although less than is the case with immune serums, Bruck, Michaelis and Schulze have attempted to employ the presence of antistaphylolysin in blood-serum as evidence of the existence of staphylococcic infection.

Furthermore, agglutination tests have shown that the staphylococci in pus may be agglutinated by the serum of the patient, while those from the air, skin, etc., are not so influenced, thus demonstrating, as is the case with many other bacteria, the variability or diversity of the strains of the bacterium in question. I feel convinced that one of the most important factors conducive to the successful production of our serum has been the utilization of staphylococcus cultures from many sources, thereby effecting a polyvalency of unmistakable potency.

PREPARATION OF THE SERUM

Eighteen strains of the *Micrococcus aureus* were isolated in pure culture from as many various sources, as follows:

Furunculosis	8
Carbuncle	4
Subcutaneous abscesses (scalp)	3
Thoracic empyema	1
Axillary abscess	1
Septicemia (osteomyelitic)	1
Total	18

Each of these strains was cultured on agar-agar tube-slants at 37 C. (98.6 F.) for twenty-four hours, and the resultant growths washed off by the addition of 5 c.c. of bouillon to each of the cultures. The combined 90 c.c. were then shaken mechanically for fifteen minutes; 0.25 c.c. of this bacterial suspension was then flooded on the agar-agar surface of each of two Petri dishes, presenting, respectively, an area of 12½ square inches. The growths from the two Petri dish cultures, after an incubation at 37 C. for twenty-four hours, were washed off in 30 c.c. of sterile 0.85 per cent. sodium chlorid solution. This bacterial suspension was standardized so that each cubic centimeter was estimated to contain 32,400,000,000 cocci and then sterilized by submersion in a water-bath at 60 C. (140 F.) for one hour.

From the same polyvalent staphylococcic suspension, agar-agar tube-slant cultures were made preparatory to the determination of the opsonic indices¹—the controls

* From the William Pepper Laboratory of Clinical Medicine, University of Pennsylvania. Read before the Philadelphia Pathological Society, Dec. 12, 1912.

1. In all determinations of the opsonic index of the animal, washed human leukocytes were employed for the purpose of phagocytosis.