Pilcz, Alexander. CONCERNING THE CLINICAL PICTURE OF EPIDEMIC ENCEPHALITIS. [Neurolog. Centralbl., 1920, No. 12.]

Attention has been called by v. Economo to certain atypical cases of encephalitis lethargica. The author here gives five cases to illustrate the variety of forms which the disease may assume. The first case was that of a woman from a neuropathic family who became ill of grippe pneumonia. At the first examination there were no signs of a lethargic encephalitis or of any other form of this disease. The threatening symptoms set in three weeks after the grippe infection and last forty-eight hours, patient becoming somnolent, with ptosis and unilateral abducens paresis, and speech disturbance. Pulse constant at 120. Temperature normal. Other nerve findings normal. This condition was followed by a hypomanic state which receded in a few days. Patient recovered, but four weeks afterward there was recurrence of grippe and death ensued. The author agrees with v. Economo's view that the agent of grippe, which is as yet unknown, sets the virus of encephalitis into activity, or that the virus, which is probably the cause of grippe existing in the same organism as the virus of encephalitis, opens the portals to this latter toxin. Author is unable to say whether the manic psychosis is to be ascribed to the grippe or to the encephalitis. The second case was remarkable from the fact that peracute symptoms of meningeal irritation first made their appearance without any phenomena of grippe, and that only in a third attack, three weeks later did the signs of encephalitis come to light. This lumbar puncture revealed meningeal changes. Hyperkinetic symptoms were only slightly shown in the last days before death. The disturbances of the pupils were rapidly progressive. The third case was that of an epileptic and the author does not attempt to determine whether the psychic symptoms (echolalia, grimacing, monotonous chanting, etc.) were due to the encephalitis or were a prolonged "psychic equivalent." The signs of encephalitis first made their appearance about a month after the beginning of the grippe. The behavior of the pupils in this case was very peculiar. They were miotic, round and reacted only to very strong light and then only slightly and tardily. But if the patient was raised up from his recumbent position, the pupils dilated and were nearly mydriatic; they also reacted promptly but the patient did not awaken from the stupor and when he was replaced in recumbent position, the pupils narrowed again and were in the same position as before. The author believes that the phenomena in sitting position were due to the fact that certain subcortical centers were awakened. The fourth case was remarkable for the fulminating course and for disturbances in the vestibular reaction; the fifth for the recurrent course, and peripheral facialis paralysis. There were also disturbances of the kidneys but it is questionable whether these belonged to the encephalitis picture. [J.]