determined by them. The relative amplitude of waves is incorrectly reproduced, and their relative prominence is often entirely wrong. The curves are frequently complicated by waves which have no existence in the vein or underlying artery but are due to inherent vibrations of the lever. This is particularly true of the Mackenzie tambours and, to a less degree, of those belonging to the Zimmermann apparatus. These artificial waves (labeled? in the records) may come after the *c* wave, as in Figure 5, in which case they may be misinterpreted for *v* waves, or, in case of vigorous heart action, they may give rise to a bifurcated wave which has been variously interpreted but which is probably entirely an artefact; or these oscillations may come after the *v* wave (Fig. 3), in which case they have been interpreted as *h* or *b* waves. While undoubtedly such a late diastolic wave sometimes exists in the case of slow hearts, there can be no question that many waves so interpreted are nothing but artefacts of the lever vibration. If these possible errors are ruled out, however, by an adequate foresight and correlation with other evidences of cardiac activity, as has been so ably done by the leaders in cardiograph work, it is clear that the records of the polygraph reproduce, with a fair degree of accuracy, the onset of the auricular wave and the fall of the *v* wave—two important features of the venous pulse.

It is interesting to inquire how forms of apparatus of such low vibration frequency as is represented in these systems can record the details of the jugular pulse at all. This is due largely to the fact that the veins are transmitting systems of low efficiency and that the details of the intra-auricular waves are considerably distorted or even annulled by the time the oscillations reach the neck veins. To record the waves still present in these veins requires only an apparatus of relatively low efficiency.

Laundry Sterilization.—In New York City some experiments were made to determine the effect of the usual routine of laundry washing in sterilizing clothes contaminated with bacteria. The following tests were made: *B. prodigiosus* and *B. coli* were smeared directly on the garment; placed on squares of gauze and folded into muslin and sewed into the garment; placed on squares made of flannel blanket, folded into muslin and sewed into the garment. These garments were then placed in the center of the net bag containing other clothing and the usual washing process applied. When the garments were returned to the laboratory they were treated as follows: A sterile swab was dipped into sterile broth and rubbed vigorously over the inoculated surface of the garment. The squares of gauze and flannel were cut into small pieces with sterile scissors and inoculated directly into the mediums. Of 42 cultures on nutrient broth 41 were positive, but all of the *prodigiosus* cultures were negative for color. *B. coli* on Conradi was negative; on lactose neutral red fermentation tubes, 17 were positive for color and 3 for gas; 4 were negative for color. Plates made from wash and rinse water (poured agar plates) showed in one instance for 1 c.c., 2,772 colonies, with a smaller number for other specimens. The temperatures of last hot rinsing water were from 50 to 60 C.—*Weekly Bulletin* Department of Health.

THE QUESTION OF SPONTANEOUS CIRRHOSIS OF THE LIVER IN RABBITS AND OTHER LAB- ORATORY ANIMALS*

ARTHUR L. GROVER, M.S., M.D.

Assistant Professor Pathology and Bacteriology, Medical School,
State University of Iowa

IOWA CITY, IOWA

In carrying on a series of experiments during the past three years on the experimental production of cirrhosis of the liver, I have from time to time noted statements by other workers that the ordinary laboratory animals, particularly the rabbit, are apt to show a spontaneous cirrhosis of the liver. In view of the importance of the question to others working along the same lines and along other lines where the spontaneous proliferation of connective tissue in these animals has an important bearing on results of experimental work, it seems as if a review of the literature and the results of my own findings might have some value.

Klopstock,¹ who did considerable work with rabbits in an endeavor to produce experimentally cirrhosis of the liver, stated that rabbits might show a spontaneous affection of the liver, with connective tissue proliferation, extensive enough to be considered a cirrhosis. Siegenbeck van Heukelom² expressed the same opinion.

Laffite³ mentions finding two cases of the disease in rabbits. Von Hanse-
mann⁴ believed that a spontaneous tuberculosis could appear in guinea-pigs and produce cirrhosis of the liver.

Joannovics⁵ and Christian both say that occasionally one finds a cirrhotic condition of the liver in rabbits and guinea-pigs that have not been used for experimental work and were supposed to be healthy animals.

Robertson,⁷ in discussing a previous communication read at the meeting of the American Medical Association, said that at one time, in the Department of Pathology of the University of Minnesota, work was started on experimental alcoholic cirrhosis and that it had to be abandoned, as far as using the rabbit went, because of the numerous sclerotic changes which occurred in this animal, not only in the vessels, but also in the liver, kidneys and heart.

* From the Laboratory of Pathology and Bacteriology, Medical School, State University of Iowa.

1. Klopstock: Zur Lehre von der Lebercirrhose, Berl. klin. Wchnschr. 1910, Nos. 33 and 34.

2. Van Heukelom, Siegenbeck: Die experimentelle cirrhosis hepatitis, Ziegler's Beitr. z. path. Anat. u. z. allg. Path., 1896, xx.

3. Laffite: L'intoxication alcool, experimentale et la cirrhose de Laennec, Thèse de Paris, 1892.

4. Von Hanse-
mann: Dissert. Verhandl. d. Deutsch. path. Gesellsch. 1904, viii.

5. Joannovics, Ueber experimentelle Lebercirrhose, Wien. klin. Wchnschr., 1904, xvii, 25.

6. Christian: Experimental Cardiorenal Disease, Arch. Int. Med., October, 1911, p. 468.

7. Robertson, H. E.: Discussion on Experimental Cirrhosis by Grover, THE JOURNAL A. M. A., 1913, Aug. 16, p. 461.

Fig. 5.—Two consecutive tracings of venous pulses first taken with optical segment capsules (upper) and then with tambour of Mackenzie ink polygraph (lower).

More lately Beitzke⁸ has reported two cases of spontaneous cirrhosis of the liver in rabbits. According to the illustrations accompanying the article, a marked degree of cirrhosis was obtained. The animals were of the series that Stillings previously used in his tumor transplantation experiments.

On the other hand, Saltykow⁹ and Lissauer¹⁰ say that they have both conducted a great many necropsies on rabbits but have never seen spontaneous cirrhosis. Such workers as Fahr,¹¹ Fischler,¹² Kretz,¹³ Ogata,¹⁴ Opie,¹⁵ and many others who have used rabbits very extensively in experimental work, make no mention of finding such conditions. Klopstock¹ and D'Amato¹³ report that an infection in the liver of dogs with *Coccidia* may give a cirrhosis. Saltykow⁹ agrees with this.

During the last four years I have examined the livers from about three hundred rabbits. In every case careful microscopic examination has been made. In no instance has there been any evidence of what might be classed as a spontaneous cirrhosis. A cirrhotic reaction has been experimentally produced in ten cases by the use of alcohol, in one case by tying the common bile duct, in another case by the production of a pyemia, and finally in another case by chloroform intoxication. In this series are about twenty-five rabbits that died as a result of spontaneous infection with pyogenic bacteria. All showed extreme abscesses at necropsy. Three of these showed a well-marked proliferation of connective tissue in the interlobular spaces. In another case there was a marked infection of the liver with *Coccidia ovi-forme*. The accompanying photomicrograph shows two large cysts filled with coccidia and lying between them an area of liver tissue almost entirely obliterated by a dense area of fibrous connective tissue.

Two large liver cysts filled with coccidia, with an area of liver tissue between almost obliterated by dense fibrous connective tissue.

This series of rabbits is made up of animals from a great many sources. Many of them have been raised

in the laboratory and one would expect the result to be the same as in any other series of a similar size.

None of the authors quoted, except Beitzke,⁸ has given specific data. In his case the previous work done on the animals might easily have been complicated with a mild infection that was overlooked, or the transplanted tumor material may have introduced substances toxic enough to have produced a toxic form of cirrhosis. In our series no rabbit has been found that showed any evidence of cirrhotic changes in the liver except where such changes could be accounted for by experimental work or by infection present at the time. From the light of our experience we believe that the cases previously reported can be explained on the ground that either some infection present in the animal at the time had been overlooked, or else such an infection had been present previously and was healed. If the latter condition is the case, it seems that the percentage of its occurrence is not high enough in the least to affect the value of the rabbit as an animal for use in the laboratory for experimental work on cirrhosis of the liver; for if such a condition should occur in a large percentage of rabbits, it would certainly have appeared in our three hundred. Regarding the other laboratory animals, we have never seen a case of cirrhosis in a guinea-pig in a series of several hundred necropsies. As to the dog, our experience is too limited to be of value.

To sum up, we would consider that, so far, not sufficient evidence has ever been brought forward to invalidate the use of animals in experimental work on liver cirrhosis.

The Transmission of Disease.—

Though this is of such vast import, it has received little investigation in the past. To the middle of last century, inquiries had led little further than the subsistence of the subject—we distinguished, classified and named our various bodily afflictions. We also did more; for we had ascertained empirically the physiologic and therapeutic effect of many drugs—quinin, opium, mercury, iodid of potassium, ipecacuanha, and others; and also that great genius Jenner had discovered a wonderful fact which is at the basis of the theory of immunity, prevention, and cure. But the causes of disease and their mode of transference still remained hidden. At that epoch, however, and afterwards Pasteur, Koch, Lister, Behring and others created the science of bacteriology. But still, though the information thus obtained was of prime importance to the pathologist, the physician, and the surgeon, we still remained much in the dark regarding one important issue, namely, the exact path of transmission of many diseases from man to man. Bacteria are easily saprophytic and may have many paths of transmission, and I fear that we are still very much in doubt as to the most important of these numerous routes.—Ronald Ross, *Lancet*, London.

8. Beitzke, H.: Ueber spontan Cirrhose der Leber in Kaninchen Zentralbl. f. allg. Path. u. path. Anat., 1914, xxv, No. 14.

9. Saltykow: Experimentelle Forschung über die pathologische Anatomie des alkoholismus chronicus, Zentralbl. f. allg. Path. u. path. Anat., 1911, xxii, No. 19.

10. Lissauer: Lebercirrhose bei experimenteller Intoxikation, Virchows Arch. f. path. Anat., 1914, ccxvii, Part 1, p. 56.

11. Fahr: Zur Frage des chronischen Alkoholismus, Verhandl. d. deutsch. path. Gesellsch., 1909, xiii.

12. Fischler: Experimentelle Lebercirrhose, Deutsch. Arch. f. klin. Med., 1908, xciii.

13. Kretz, R.: Ueber Lebercirrhose, Wien. klin. Wchnschr., 1900, No. 12, p. 27; Lebercirrhose, Verhandl. d. Deutsch. path. Gesellsch., 1904, viii.

14. Ogata: Beitr. z. exper. erzeugten Lebercirrhose und zur Pathogenese des Icterus mit spezieller Berücksichtigung der Gallenkapillaren bei der Unterbindung des Ductus choledochus und der Iktrogenngiftung, Beitr. z. path. Anat. u. allg. Path., 1913, xi.

15. Opie: Studies from the Rockefeller Institute for Medical Research, 1910, xi, 367.

16. D'Amato: Ueber experimentelle, vom Magendarmkanal aus hervorgerufene Veränderungen der Leber und über die dabei gefundenen Veränderungen die übrigen Bauchorgane, Virchows Arch. f. Path. Anat., 1907, clxxxvii.