Health, under whose supervision the inspectors are placed. Thus, the State Board of Health of Massachusetts, through the State Inspectors of Health, is in position to acquire information relative to the sanitation of local communities throughout the commonwealth, including (1) matters relating to water supplies and sewage; (2) the prevalence of diseases dangerous to the public health in the home, the schoolhouse, the factory or elsewhere; (3) the sanitation of schoolhouses and industrial establishments; and (4) information concerning the health of young persons in factories, *at their work*, and the influence of such occupation upon the health of these minors.

To collect data concerning the health of all young persons of each sex in the factories in Massachusetts is itself a task of considerable magnitude; to consider, in addition, the possible injurious influences to which minors are exposed in certain occupations or processes while at their work makes even greater demands upon the inspectors' time. These facts, taken into consideration with the fact that the duties imposed upon the State Inspectors of Health are varied, make it obvious that the men are handicapped in their study of the health of young persons in relation to occupational diseases.

But were the health inspectors relieved from certain work, as, for example, that of inspecting slaughterhouses and provisions, a duty clearly not intended to fall within their province when they were appointed state inspectors of health, there still would be a difficulty impossible to overcome without further legislation. I refer to the lack of proper places for the physical examination of young persons whenever, in the opinion of a state inspector of health, such examination is advisable or necessary. This is a serious handicap in connection with the work of the inspectors, not only because the absence of suitable places for the examination of minors means inaccurate and incomplete data concerning their health, but because the absence of accurate and complete data means the impossibility of accurate statistical information to show either the percentage of minors who are in ill health in our factories, or the percentage of minors whose health is more or less affected as the probable result of injurious influences which arise from the different occupations or processes.

To be sure, there is another side to this question, as is shown by the very promising work already accomplished by the inspectors. These officials have obtained a considerable knowledge and can, in the course of time, obtain a full knowledge of the conditions of employment. They point out certain dangers which are avoidable, so that the young persons may no longer be subjected to such avoidable unhygienic influences; they point out dangers which, while not entirely avoidable, make the occupation unsuited to young persons; and, finally, they brand as dangerous those processes and trades which from their nature are intrinsically dangerous to health, hence unfit to be followed by boys and girls. All this can be and is done; but with a law providing better

convenience for examining young persons, in addition to the present opportunity of observing them at their work, the system would be complete except for the limited appropriation for carrying out the statute provisions.

An examination of a young person ought to take into account and to record the general health, the personal or family history if important, observations of the minor while at work, fitness for the particular kind of work done, age, sex, height and weight. Such examinations ought to be made with sufficient frequency, which, in any trade or process, should not be less often than once a year, and in some occupations or processes, if permitted at all, not less often than twice or three times a year. Under existing conditions, however, so thorough a system can only partially be carried out.

When conditions of ill health or of physical unfitness of young persons are discovered, aside from communicable diseases, already considered, two courses of action are open to the health inspectors. They may inform the child's parents, or, if it appears to be to the child's advantage, his employer. In the latter way many young persons have been aided, if not completely relieved, by the willingness of an employer at the suggestion of the health inspector to change their work, and, in some instances, by the interests of employers in paying, in part, for the treatment of children found in their employ afflicted with such a disease as tuberculosis. On the other hand, in the absence of authority to exclude from factories young persons found in ill health or physically unfit, the parents of the children take but little notice of information and advice given to them.

In the school, when such affections as pediculosis, scabies, blepharitis, trachoma, etc., are detected, they are vigorously excluded until remedied. Why not in the factory? In the school, each child's sight and hearing are tested annually. Why are not the sight and hearing of each young person in a factory tested annually and a judgment formed for each child based on the kind of work done and the conditions under which the work is done? Why should not the children in factories receive not only as great but greater attention than those in our schools, when many of the factory children are exposed to influences which are far more inimical to health than the conditions to which schoolchildren are subjected in schoolrooms, and for longer daily periods?

SUBMUCOUS FIBROIDS.

BY PHILEMON E. TRUESDALE, M.D., FALL RIVER, MASS.

THESE tumors are commonly considered and described as benign. The influence of their growth, however, upon the health of the individual is sufficiently serious to warrant a consideration of them in the class of malignant growths. Aside from the chronic irritation which they produce, and the consequent carcinomatous and sarcomatous degeneration of the tumors themselves, they

are often unrecognized causes of profound anemia, with accompanying disorders of the heart, blood vessels, and kidneys.

Their presence is most frequently associated with one or both of the subserous and intramural forms. When so, the diagnosis is easily made. On the other hand, when the submucous fibroid occurs alone, its presence is not often suspected until it has attained considerable size, or until the patient's health has suffered to a very marked degree.

In its origin this tumor is probably intramural, but by continued contraction of the surrounding muscle structure it is forced into the uterine cavity, losing its false capsule of plain muscle fiber and becoming pedunculated or sessile. The former, though often a source of persistent bleeding and occasionally of alarming hemorrhage, is usually forced through the cervical canal by the contracting uterus. Here it is readily discovered by digital or visual examination and easily removed with the curette. The sessile fibroid, however, though occasionally expelled, commonly remains an intra-uterine tumor. It frequently has broad basilar attachments and may continue to develop to enormous size, while the uterus itself hypertrophies markedly in its effort to expel the tumor. This is well illustrated in Fig. VI.

In structure, the submucous fibroid is essentially a plain muscle growth. It contains a comparatively small amount of connective tissue and a relatively rich blood supply. It is, therefore, softer than either the subserous or intramural type, more rapid in its growth, and a source of danger from hemorrhage, degeneration, and necrosis sufficiently great to warrant its comparison with a malignant neoplasm.

As a cause of pain, it is a more pronounced factor than either of the other types, especially at the menstrual period, when the uterus contracts forcibly in its effort to expel its contents. Dysmenorrhea with any excess in loss of blood occurring in a woman of thirty years or over should lead one to observe carefully for the presence and development of a submucous fibroid.

The intramural or subserous growth may attain enormous size without causing symptoms of pain or pressure. These forms produce early an irregularity in the contour of the uterus, and, unless their development meets obstruction from adhesions, they merge from the pelvic cavity without difficulty, and, as long as they remain free, may cause very little disturbance from pressure. This is probably accounted for by the frequent changes of the position of the tumor with changes of posture.

We find the circumstances different in the development of the submucous fibroid. It first encroaches upon the uterine cavity, taking up this space; its growth continues in globular form, and the contour of the superimposed uterus will be found very often to fit the concavity of the pelvis into which it has prolapsed by reason of its own weight. The uterus becomes virtually incarcerated, and with the continued growth of the tumor there follows edema of the legs,

phlebitis, bluish discoloration of the vulva, and vaginal mucosa, obstipation and scanty urine with frequency. These symptoms were present in Case No. 477. In delivering this uterus from the pelvic basin considerable traction was required, not because of the tumor weight or adhesions, but for the reason that the mass had grown to fit snugly into the concavity of the pelvis.

We are reminded by modern writers that fibroid tumors of the uterus should be detected early and should be removed as soon as they are the cause of symptoms local or remote. To discover an intramural or subserous fibroid of appreciable size requires very little diagnostic skill, yet to determine the presence of a small submucous fibroid may be difficult for a gynecologist of ability. The symptomatology is of the utmost importance in arriving at a correct diagnosis. It is no longer necessary to palpate an inflammatory mass around the appendix before making a diagnosis of appendicitis, nor to feel a distended gall bladder before determining a cholecystitis; neither should it be essential with an otherwise perfect clinical picture of uterine myoma to wait until the tumor has become as evident to the patient as to the examiner. The size of these tumors as seen at the operating table is sufficient evidence of our inability to discover them before they have caused years of suffering.

In a typical case of submucous fibroid there is gradually increasing menorrhagia and dysmenorrhea, with consequent anemia and debility. At first the increased loss of blood is confined to the menstrual and immediate postmenstrual period. Treatment is rarely sought for local manifestations; instead, the patient seeks advice for the secondary anemia and the so-called nervous debility. Unless the history is fully taken, the abnormal menstruation may not receive mention. This is especially true if the patient is at the period of life approaching the menopause. Any variation from normal may occur at this time without causing anxiety. Hemorrhages may occur during the menopause without pathological significance, and, although this is exceptional, it is for this reason that excessive loss of blood is not looked upon with alarm, but as a final "clearing" before cessation.

A woman's own interpretation of normal menstruation should not be trusted, especially at this period of her life. A careful history and vaginal examination alone are the means by which one may determine the physiological or pathological significance of uterine hemorrhage. Any loss of blood in excess of the normal habit of the individual demands this investigation, and the examiner will frequently find a fibroid as the cause of a group of symptoms which are quite remote from the pelvis. Although the amount of blood loss usually increases with the development of the tumor, this is no criterion by which one may be guided. In Case No. 434 the submucous tumor was no larger than a sparrow's egg, and the hemorrhage was profuse.

Enlargement of the uterus from subinvolution, metritis, endometritis and pregnancy enter into consideration for differential diagnosis. In subinvolution one finds enlargement of the uterus usually with displacement, and the condition a sequence of labor. In metritis and endometritis the changes are usually inflammatory following puerperal or other infection. The passage of a uterine sound is sometimes an aid in determining an irregularity in the uterine cavity. Before this instrument is used one should eliminate beyond doubt a possible pregnancy.

A small submucous fibroid and an early pregnancy may present a striking similarity in physical signs. The soft cervix, patulous os and symmetrical enlargement of the body of the uterus are common to both. There may also exist a bluish discoloration of the vaginal mucosa and enlargement of the veins in the lower extremities by the presence of a submucous myoma.

The following case is an illustration (No. 477, Fig. 1): M. S., age thirty-two, single, bookkeeper, good family history. Catamenia began at sixteen, regular up to July, 1908, twenty-eight-day type, four days' duration, used ten napkins. Patient was well up to July, 1908, when she noticed edema in the right foot, soon involving the entire leg. She remained in bed for four weeks,

Fig. 1 (a).

U. W., Uterine wall. S. T., Submucous tumor. U. C., Uterine cavity. C. U., Cervix uteri. O., Ovary.

and the swelling gradually subsided. Menstruation came on more freely in amount than usual and continued for three weeks. She has had menorrhagia and metrorrhagia since, using from twenty-five to thirty napkins each month. She consulted me for the phlebitis which had returned in her right leg. She had kept at work in spite of the discomfort from her swollen and tender veins. There was obstinate constipation with vesicle

irritability and a diminished amount of urine, 30 oz. in twenty-four hours. Upon vaginal examination, I found a bluish discoloration about the introitus and vaginal mucosa, a soft cervix, an os patulous, admitting the tip of the index finger, a symmetrical enlargement of the uterus corresponding to about the size of a three-months' pregnancy. The breast examination was negative. From history of continued hemorrhage, as substantiated by the patient's mother, there seemed a reasonable certainty of submucous fibroid, and this proved to be the condition found at operation.

Fig. 1 (b).

E., Fundus. S. T., Submucous tumor. E., Endometrium. U. W., Uterine wall. S. S. T., Subserous tumor.

Upon exposing the uterus to view, its symmetry, color and density were consistent with pregnancy, yet a feature which made a distinction possible was the anatomical relation of the broad ligaments in their attachment at the fundus. In pregnancy the enlargement of the uterus is physiological and the broad ligaments retain their normal anatomical relation with the body of the uterus, whereas in fibroid disease the relations of the broad ligament attachments vary according to the position of the tumor within the uterine cavity.

Another case of a somewhat different type is the following, in which I failed to recognize the submucous tumor when it was small, yet the cause of a debilitating flow (Fig. II, No. 436): F. T., thirty-three years, married ten years. Family history negative. One child born 1901, instrumental labor, normal convalescence. Catamenia began at sixteen, regular twenty-eight-day type. For two years the duration has been six days, using sixteen napkins and in bed one day from dysmenorrhea. In February, 1908, she was seen with Dr. E. F. Curry and cured. The uterus was enlarged, but no irregularity was felt bimanually or with the sound. The condition was thought to be subinvolution with endometritis. No benefit followed this procedure, and in May, 1909, an irregularity at the fundus, due to fibroid, was discovered. Hysterectomy was done May

19, 1909. Pathological examination showed the uterine wall 1.5 cm. in thickness. In the cavity, attached near the fundus by a sessile base, was a roundish tumor 4 cm. in diameter. The endometrium near the fundus where the tumor pressed was necrotic looking and had a foul odor. Histological examination of the tumor showed a plain muscle growth. The endometrium showed necrosis of some glands and hypertrophy of others. This tumor within all probability existed as a cause of her symptoms at the examination which I made in 1908.

FIG. II.

U. W., Uterine wall. R. T., Submucous tumor. U. C., Uterine cavity. C., Cervix.

The nervous symptoms which many women manifest following a period of uterine hemorrhage vary widely from simple irritability to a state of marked prostration. In a careful observation of women of nervous temperament one is impressed by their extreme sensitiveness to the loss of blood.

No. 454, Fig. III: E. L. A., age forty-one, nurse. One brother operated upon for malignant growth of bladder. Family history otherwise good. Had typhoid

FIG. III.

P. F., Pedunculated fibroid. Sm. F., Submucous fibroid. U. W., Uterine wall. C., Cervix. S. F., Subserous fibroid.

fever in 1897. Recovered fully and gained in weight to 150 lb. Now weighs 110 lb. Catamenia began at seventeen, regular twenty-eight-day type. Menorrhagia for five years, flowing seven days and soiling twenty to twenty-five napkins. Dysmenorrhea for

first two days, with nervous symptoms predominating. In 1907 suffered from neurasthenia which was thought to have been brought on by overwork. Prior to the onset of this nervous state there had been continued menorrhagia for several months and the patient had little opportunity for rest. After a period of six months of absolute rest, she recovered but was not sufficiently well to warrant a return to work. She continued to have periods of depression and indisposition, though milder in degree. For the last six months the loss of blood had been excessive both at the menstrual and intermenstrual periods. She had intermittent bearing-down pains across the lower abdomen, with a feeling of pressure in the pelvis. She suffered from headache, vertigo, insomnia and extreme irritability. There was palpitation, dyspnea upon exertion, constant fatigue, cold extremities, numbness and neuralgic pains. Appetite was poor and bowels were very sluggish. Frequency of urination only in the daytime. January, 1909, for the first time, she felt a tumor within her abdomen. She was emaciated, anemic and extremely nervous. Heart sounds normal, urine negative. The lower abdomen was occupied by a hard, irregular, movable tumor. Vaginal examination revealed the uterus indurated, irregularly enlarged and movable, together with the fibroid tumor above which extended upward two inches above the umbilicus.

Hysterectomy was done and the specimen marked 454 was removed, consisting of an enlarged uterus with wall 1.5 cm. in thickness. Filling the uterine cavity was a globular submucous tumor measuring 5 cm. in diameter. There were multiple subserous fibroids and a pedunculated mass attached to the fundus measuring 5 cm. by 12 cm. No degeneration was noted. There was a hydrosalpinx from one side. Both ovaries were apparently normal and were not removed.

Case 434, Fig. IV, represents a condition in which the submucous tumor was small, had undergone adenomatous degeneration and was associated with hemorrhage of a marked degree. J. K., age forty-six,

FIG. IV.

A. M., Adenomyoma. U. W., Uterine wall. U. C., Uterine cavity. C., Cervix.

four-para, normal labors. Last child born 1903. Two miscarriages. Father died of cancer of the liver. She was operated upon in Boston for chronic appendicitis in 1902. Only other illness was tonsillitis. Catamenia began at fourteen, twenty-eight-day type, normal in quantity until 1906, when she began to notice an increase in the amount lost at each menstrual epoch. This continued for one year; then there developed metrorrhagia as well as menorrhagia. At times bleeding persisted from two to five weeks and was not accompanied by pain at any time. November, 1908, there was a uterine hemorrhage of alarming

proportions accompanied by dizziness, syncope and extreme prostration. Bleeding occurred at varying short intervals and in varying amount since. With this history she was referred to me by Dr. Annie C. Macrae.

The patient showed a moderate degree of anemia. Uterus was about twice the normal size, symmetrical, in normal position, freely movable, but tender to pressure. Nothing abnormal was felt in either broad ligament. In view of her age, the character and degree of the uterine hemorrhage, with consequent loss in weight and strength, and the enlargement of the uterus, I advised hysterectomy, to which she assented. When the uterus was exposed there was observed an enlargement with an irregularity in the left cornua. Supravaginal amputation was done. The ovaries were apparently normal and were not removed. The gall bladder was found to contain many stones; cholecystostomy was done with the removal of twenty stones varying in size from a split pea to a common marble. It was subsequently learned that this patient had had two attacks of acute indigestion during the past year; both occurred at night after being in bed some time and were characterized by epigastric pain, vomiting and diarrhea.

Examination of the specimen showed the uterine cavity measuring 6 cm.; the walls measured 3 cm. At the left cornua was a roundish tumor 3 cm. in diameter, one border of which was directly beneath the mucosa. Histological examination of the growth showed a tissue made up of plain muscle fibers scattered throughout which were numerous glands lined by one layer of epithelium resembling the glands of the endometrium. There was also a simple hyperplasia of the glands of the endometrium.

The reluctance on the part of some women to submit to operation except as a "*dernier resort*" is responsible for the size and many complications accompanying fibroid tumors that we see at the operating table. After the gamut of medicine has been tried, electro-mechano- and psychotherapeutics are applied with renewed hope.

The extremes to which a patient will sometimes venture in this respect is well illustrated by the following case, referred to me by Dr. George Parlow:

No. 445, Fig. V: L. M., age forty-seven, married twenty-seven years, clerk. Negative family history. One child, 1883, normal labor. Menorrhagia for seven years. During this period she has required general tonic treatment at varying intervals. For two years the menorrhagia has been more severe. Metrorrhagia for six months. She has suffered increasing debility, headache, disturbed sleep, cold extremities, palpitation, nervous irritability, digestive disturbances and neuralgic pains. There was at intervals a respite for one or two months, and the severity of her symptoms was dependent upon the degree of hemorrhage. In 1907 a pelvic tumor was discovered and operation advised. Growth had increased to the size of a child's head. For six months there had been intermittent attacks of pain across the lower abdomen, accompanied by nausea, vomiting, obstipation, fever, distention and tenderness. They would usually subside after a flow of considerable purulent material from the vagina. The attacks would last from three to seven days and were occasionally ushered in by a chill. There was marked dyspnea upon exertion. She had been wholly incapacitated for work for two months. The blood examination was as follows: Hemoglobin, 25%; color index, 0.5; red cells,

2,400,000; white cells, 12,000; polymorphonuclear, 87%; small mononuclear, 2%; large mononuclear, 6%; transitionals, 2%; eosinophiles, 0.8; myelocytes, 0.4.

FIG. V.

U. W. Uterine wall. U. C. Uterine cavity. Sm. F. Submucous fibroid. C. Cervix. Ss. F. Subserous fibroid.

The red cells appeared undersize with marked achroma. Frequent variation in size of red cells. No normoblasts or megaloblasts. The pulse was 112 and of low tension. There was a soft blowing systolic murmur heard over the entire precordium and transmitted into the neck. Urine was normal; there was no edema of the extremities. A hard lobulated tumor with limited mobility occupied the lower abdomen. It extended about one inch above the umbilicus and well out to the iliac crest on the left side. Vaginal examination revealed the pelvis filled by a slightly movable tumor involving the uterus and its appendages. Tenderness behind and to the left side.

Although it was evident that there was a degenerating submucous myoma and an inflammatory infiltration in the pelvis, surgical interference with hemoglobin at 25% was not considered advisable. She entered my hospital on June 5, 1909, and was given $\frac{3}{4}$ gr. of citrate of iron daily by hypodermic injections and as much nourishment as she could take. For two weeks her temperature ran an irregular course, ranging between 99° and 102°. June 21 the hemoglobin was 52%. On the following day the abdomen was opened. A large fibroid tumor, together with an inflammatory mass, occupied the pelvis and lower abdomen. There were adhesions to the omentum, sigmoid and small intestine. On the left side the ovary had become sacculated and contained a quantity of pus. This was ruptured in its removal. There was a double pyosalpinx. A fibroid tumor involving the uterus with its adnexa was removed. Drainage. Recovery was rapid and uneventful. She left the hospital four weeks later with her hemoglobin at 80%.

Another case of less severity but where surgical interference was postponed for many years is the following:

No. 453, Fig. VI. H. C., patient of Dr. Mayhew, Vineyard Haven. Age, forty-five years. Negative family history. She began to have menorrhagia in 1901. One year later operation was advised for tumor of the uterus. Her secondary symptoms lasted only one week during each month and were chiefly of nervous origin. She improved for one year under general tonic treatment. Menorrhagia increased and she tried

electricity for about a year with some benefit. Then the vibrator was used and other forms of local treatment, with a view to "shrinking the tumor" until the menopause would establish itself and thus save her the ordeal of an operation. Instead, however, of a lessening in the degree of hemorrhage with dysmenorrhea, there was a steady increase, accompanied by dysmenorrhea. In addition, during the past year there have been frequent occurrences of intermenstrual hemorrhage. She had the common symptoms of secondary anemia, together with pressure upon the rectum, bladder and sacral plexus. There was a faint systolic murmur at the apex, not transmitted, and an

growth acts as a foreign body, and the "labor-like" pains so frequently complained of are due to this intermittent contraction of the uterus.

FIG. VI.

Sm. T. Submucous tumor. H. Hemorrhage. U. W. Uterine wall. C. Cervix.

accentuated pulmonic second sound. Hysterectomy was done, together with the removal of one enlarged ovary. Section of the excised uterus revealed a large submucous myoma with a sessile base nearly the extent of the long diameter of the tumor, 9 cm. The endometrium covering the tumor presented a blotchy appearance from the submucous hemorrhages. Sections of the tumor showed a plain muscle growth. This specimen shows in an interesting manner the enormous hypertrophy which the uterus may undergo in its effort to expel the tumor from its inner wall. The

FIG. VII.

U. W. Uterine wall. Sm. T. Submucous tumor. C. Cervix.

An excellent specimen of a submucous fibroid I recently obtained from a patient referred to me by Dr. John H. Gifford.

FIG. VIII.

Represents Fig. VII with inversion of uterus.

I. U. Inverted uterus. E. Endometrium. Sm. T. Submucous tumor.

No. 508, Fig. VII. M. T., age forty-six years, housewife. Family history negative. Two children, first born 1884; second, 1888; both labors normal. Catamenia began at twelve, twenty-eight-day type. Six

FIG. IX.

U. Uterus. F. T. Fallopian tube. O. Ovary.

years ago began to have menorrhagia with dysmenorrhea. In 1906 she was curetted for excessive flowing. This resulted in temporary benefit. During the past year the menstrual periods have been accompanied by an excessive loss of blood with the usual train of symptoms following. There has been dysmenorrhea and pressure upon the rectum and bladder. The patient had noticed a marked increase in size of the tumor in six months.

Hemoglobin was 70%. There was a soft systolic heart murmur of hemic origin. Uterus was enlarged to the size of a six-months' pregnancy, symmetrical and freely movable, cervix soft, os patulous, admitting the tip of the index finger, by which was felt the submucous fibroid. The abdomen was opened by a median incision, and the tumor exposed as in Fig. IX.

It presented a perfectly smooth symmetrical surface and everywhere yielded readily to pressure, though to a lesser degree than would be expected in the pregnant uterus. The color was not the healthy shade of a normal uterus, but more dusky, and enlarged veins were visible beneath the peritoneum. The broad ligament, as the photograph shows, was so stretched out of its normal relations that at the fundus the tube was a mere string. The ovarian ligament, which normally has its attachment to the fundus near that of the fallopian tube, and retains that relation during the physiological enlargement of pregnancy, was, in this specimen, widely separated from that position by the degree of stretching which occurred in the uterine wall. The dark areas at the fundus were due to bleeding from use of double hooks.

The growth of this tumor was so rapid that there evidently was not time for any marked degree of hypertrophy in the uterine wall to occur except at the upper anterior third. The remaining two thirds, including the entire posterior wall, was a mere sac.

An analysis of 700 cases of fibromyomata of the uterus from the Bender Laboratory, made by Ellis McDonald, showed the submucous tumor to have occurred alone in only 75 cases, or 10%, yet multiple fibroids were found in 299 cases, or 42.7%.

In 44 cases in which I have done hysterectomy for fibromyomata, the submucous tumor was found alone in 6 cases, or 13.6%, while its presence was accompanied by one or both of the other forms in 18 cases, or 40.9%. It was noted, therefore, in 24 specimens, or 54.5%.

Of these 24 cases, 2 had undergone hyaline degeneration, 2 were adenomyoma, 1 myxoma, 3 had undergone necrosis. Four specimens were not examined microscopically.

The submucous fibroid had been the cause of hemorrhage or dysmenorrhea or both in every case in which it was found; and in many instances it was present and overlooked when a cause for these symptoms had been sought for many months, and occasionally for years.

The operation employed in these cases was supravaginal hysterectomy with implantation of the round ligaments into the stump of the cervix and conservation of any healthy adnexa. The number of cases, 40, was my total, done consecutively, and with no deaths.

Montgomery advocates the enucleation of uterine fibroids when the growths are few in number and the structure of the uterus but little involved; when the growths are readily accessible

through the vagina or cervical canal; when the woman is under forty years of age, and when the tubes and ovaries are free from complicating conditions.

An examination of other structures both in the pelvis and abdominal cavity in the presence of a submucous tumor of any size would seem to be of such importance as to limit the vaginal operation to the removal of polypi.

As for the transperitoneal enucleation of submucous myoma, Case No. 436, a specimen which was removed from a married woman aged thirty-three years, would seem to present the indication for that procedure, and yet, when this method of removing fibroid tumors was employed in about forty cases reported by Dr. Montgomery, there were two deaths, some stormy recoveries and some subsequent morbidity. Therefore, in the light of our present knowledge, the comparative risks of the operations and the possibility of malignant degeneration of the tumor should be clearly presented to the patient.

If she is willing to accept the added risks, though very small, to preserve her uterus, it is up to the surgeon to act accordingly. The operation is her investment for the future. She should be enlightened and her wishes considered.

SUMMARY.

1. The submucous fibroid, clinically, may be considered a malignant growth.
2. It commonly occurs with one or both of the submural and subserous types, and is then easily diagnosed.
3. It occurs alone with relative infrequency, and is then usually not discovered until late.
4. Hemorrhage at the menstrual periods and between the periods, frequently with dysmenorrhea, characterizes its presence.
5. When it is of large size and sessile, hysterectomy is the operation of safety. In certain cases the choice of operation may be left with the patient.

I am indebted to Dr. Annie C. Macrae and Dr. Mary W. Marvell for the painstaking pathological examination of my specimens.

Clinical Department.

PERIODICAL HEMOPTYSIS.

BY THEODORE J. EASTMAN, M.D., BOSTON, MASS.,

Consulting Physician, Massachusetts Charitable Eye and Ear Infirmary; Assistant Visiting Physician, Long Island Hospital; Assistant Physician to Out-Patients, Massachusetts General Hospital, Boston.

THE close physiological connection between the female genitalia and the respiratory tract was recognized as far back as the day of Hippocrates, who said, "Hemoptysis in a woman is removed by eruption of the menses."¹ While this is not wholly correct to-day — perhaps from a greater prevalence of tuberculosis — still, we find occasional cases in which its truth is apparent.

Shortly before and often during menstruation, there is noticed in many women a congestion and