The Boston Medical and Surgical Journal

Table of Contents

December 16, 1915

ORIGINAL ARTICLES

DISINFECTION OF SEPTIC JOINTS. By Frederic J. Cotton, M.D., Boston.....	905
THE BACTERIN TREATMENT OF CERTAIN CHRONIC PYOGENIC DERMATOSES. By Charles C. Dennie, M.D., Kansas City, and John H. Buford, M.D., Boston.....	910
ALOPECIA SYPHILITICA. By Douglass W. Montgomery, M.D., San Francisco.....	913
EXAMINATION AND PROPHYLAXIS FOR SYPHILITIC PATIENTS AND THEIR FAMILIES: METHODS OF INVESTIGATION AT THE PSYCHOPATHIC HOSPITAL, BOSTON, MASSACHUSETTS, 1915. By Helen M. Wright, Boston. (Continued.).....	916
ESTIMATE OF THE COST OF SOCIAL WORK IN CONNECTION WITH EXAMINATION AND TREATMENT OF OTHER MEMBERS OF FAMILIES OF SYPHILITIC PATIENTS THROUGHOUT THE STATE, BASED UPON 147 CASES DEALT WITH IN EIGHT MONTHS AT THE PSYCHOPATHIC HOSPITAL. By Mary C. Jarrett, Boston.....	921

CLINICAL DEPARTMENT

FRACTURE OF THE TRANSVERSE PROCESSES OF THE FIFTH LUMBAR VERTEBRA. By Lloyd T. Brown, M.D., Boston, and W. J. Dodd, M.D., Boston.....	922
A CASE OF DERMATITIS MEDICAMENTOSA. By William S. Walsh, M.D., Providence.....	924
A CHEMICAL AID IN SECURING THE SPIROCHETA PALLIDA FROM SYPHILITIC LESIONS. By Henry D. Lloyd, M.D., Boston.....	925

HARVARD MEDICAL SCHOOL

LEGAL MEDICINE.....	925
---------------------	-----

BOOK REVIEWS

Occupational Affections of the Skin. By R. Prosser White, M.D., Edin.; M.R.C.S., Lond.....	928
--	-----

Diseases of the Skin and the Eruptive Fevers. By Jay Frank Schamberg, M.D.....	928
Diseases of the Arteries. By Sir Clifford Albutt, K.C.B., M.A., M.D., F.R.C.P., F.R.S., LL.D., D.Sc.....	928
Handbook of Physiology. By W. D. Halliburton, M.D., LL.D., F.R.C.P., F.R.S.....	928
A Text-book of Chemistry and Chemical Urinalysis for Nurses. By Harold L. Amoss, S.B., S.M., M.D., D.P.H.....	928
The Care of the Baby. By J. P. Crozer Griffith, M.D.....	929
The Practical Medicine Series. By Charles L. Mix, A.M., M.D., Emilius C. Dudley, A.M., M.D., and Herbert M. Stowe, M.D.....	929
Simplified Infant Feeding. By Roger H. Dennett, B.S., M.D.....	929

EDITORIALS

A NEW METHOD OF TABULATING MORBIDITY STATISTICS IN THE MEDICAL DIVISION OF THE HEALTH DEPARTMENT OF THE CITY OF BOSTON.....	930
MEDICAL LEGISLATION: AN UNJUST BILL.....	931
MILITARY TRAINING AND PHYSICAL DEVELOPMENT.....	932
EPIDEMIC ACIDOSIS IN CHILDREN.....	933
MEDICAL NOTES.....	933

CORRESPONDENCE

THE POWER OF ADAPTATION OF THE HUMAN MIND. (PARIS LETTER.) "S.".....	937
MEDICAL RELIEF IN BELGIUM. DRUG TRADE WITH GERMANY. W. G. P.....	939
EUTHANASIA IN WAR. W. B. Trull, M.D.....	940

MISCELLANY

MEMORIAL RESOLUTIONS FOR WILLIAM NOYES, M.D.....	938
BELGIAN PHYSICIANS' RELIEF FUND.....	940
NOTICES, RECENT DEATHS, APPOINTMENTS, ETC.....	940

Original Articles.

DISINFECTION OF SEPTIC JOINTS.*

By FREDERIC J. COTTON, M.D., F.A.C.S., BOSTON.

INFECTED joints have always been a reproach to our art; their treatment the counsel of despair.

A suppurating joint, infected from without or from within, has been counted among the lost, and not only joints, but lives—many lives—have been lost as well. Nothing could be worse than the story of joint drainage. Today the opening and drainage of joints should be listed not as an operation but as a crime. It is our duty today either to determine and declare a joint infection hopeless, and then to *lay the joint wide-open (not drain it)*, or else to bend all our efforts to an *optimistic attempt to save the joint*; not only to get it healed, but to preserve its function as well.

Years ago, about ten years, I was engaged in opening and washing-out gonococcus joints, as were many others. During the course of this work I learned as a purely empiric matter that such gonococcus joints, whether in the small minority showing positive culture, or the many that gave no viable cultures, that these joints did better if sewed up tightly, than if left to drain through loose suturing; while wick or tube drainage proved, as all have since recog-

nized, no less than disastrous to future function.

Not long after this series had crystallized in my mind the desirability of efficient disinfection and closure in these cases, there came to my hand some cases of really *septic* joints (for the gonococcus is but a facultative pus organism after all), joints with staphylococci particularly, and a few streptococcus joints. These I treated in the same way, sewing the capsule tight, and gradually evolved a definite technic: opening of the joint; irrigation for fifteen minutes (by the watch) with a full stream of 1-15000 corrosive sublimate†; a flushing-out of the excess with salt solution; then a water-tight closure of the capsule by mattress and single sutures in such a way as to leave the synovial membrane cut edge to cut edge, as accurately as may be. Usually the membrane is so swollen as to make this really possible.

The external wound is left wide open or nearly so, for it must be conceded that real disinfection of the external soft tissues is not practicable. After operation there may or may not be an effusion into the joint; if an effusion comes it goes again within one to three days, and is not accompanied by temperature or by any considerable discomfort.

I have not in these cases, with two or three exceptions, resorted to traction for the separation of joint surfaces. There has seemed to be no pain-factor constituting a clinical demand for such treatment, and I see no theoretical reason for it, for these joints show little or no spasm.

† As a rule the best way is to distend the joint; then let it collapse and empty; then distend again. Accurate reaching of all parts of the joint by a direct stream is almost impossible.

* Read before the Hennepin County Medical Society, June 7, 1915, at Minneapolis.

There have been about two dozen such cases treated by this technic, excluding the gonococcus cases, not less than fifteen of them representing infections by staphylococci; the majority staphylococcus infections. In looking up the cases for this paper, I have been rather appalled by my inability to find well remembered cases in the hospital records of years ago; cases of failure as well as successes.

There have been two failures: one a punctured wound of the knee in a male adult; treated according to this technic but with a recurrence of the streptococcus infection; laid wide open; patella laid back on thigh; later (after vaccine treatment) subjected to a cleaning-out of granulations from nearly intact cartilage and suture of divided ligaments, regaining in the end a good leg with 45 to 50 degrees of motion. The second case was a knee with an epiphyseal staphylococcal infection in a boy of three; cleaned out and sutured; later the joint broke down and an ankylosis resulted.

The rest have all done well; I cannot give nearly all, and will rest my case on six instances, covering seven pus joints disinfected and sewed up successfully.

CASE 1. Barbara S., aged 10; seen in consultation with Dr. John MacDonnell of South Boston. History of having hurt her knee in jumping down into a cellar and having a few days thereafter noticed tenderness and lameness gradually increasing. The fall was about two weeks previous, she had been in bed a week when seen, with great pain and steady fever. Physical examination showed a sturdy young Polish girl; nothing of note save the right knee, which showed great distention of the joint, tenderness and limitation of motion, with local heat, but without redness. I sent her into the City Hospital for immediate operation, as a septic joint, of the osteomyelitic type; one of the juxta-epiphyseal infections.

The same day, Jan. 16, 1909, I operated under ether. The joint was opened on the inner side and found full of thick, creamy pus. Usual irrigation with HgCl₂ followed by salt, and the joint closed in the usual way. Then through an incision on the outer side, a large popliteal abscess was opened and drained. The cultures taken from both joint and abscess were reported as staphylococcus pyogenes aureus by Dr. Mallory.

The temperature at entrance was 102.5 degrees, but three days later reached 99 and ran along about that level for some time. At eight days there was no fluid in the joint; tenderness was less; wound over joint nearly closed. Deep sutures held; there was no joint leakage. Two weeks after the operation, the joint was all right, but there was more pus from the popliteal space. This continued. An x-ray taken showed a sequestrum and it was decided to remove it.

March 15, 1909, under ether, I removed a sequestrum $2 \times 2\frac{1}{2} \times \frac{3}{4}$ inches from the femur, exactly at the back, starting at the epiphyseal line and running up. The temperature went up the next day and then down permanently.

March 28, 1909. Still draining from sinus behind;

joint wound healed; knee joint motion of 60 degrees. Discharged to the out-patient department.

About a year later, I removed a little bone chip with forceps; function of the knee had been perfectly good before that. After this chip came away, the sinus promptly closed.

I sent for this girl and examined her a year ago; five years after the operation. She has grown big and strong, and save for the shin scars, the two knees cannot be told apart; an absolute restoration, anatomically and functionally. She can sit on the heel of this leg.

This case was exhibited before the Clinical Surgical Congress October 28, 1915.

CASE 2. Pat. F.; seen in consultation with Dr. Thomas F. Carroll of Brockton; then of Norwood, Mass., Aug. 7, 1912. The patient, a man of 40, had an old scar and had had a childhood osteomyelitis, leaving some lameness and an old scar on the inner side of the thigh. The story was that he had then been some seven weeks ill; bedridden and feverish; that the knee had been opened by some doctor. Owing to his extreme deafness, detail data were scanty. There was, as Dr. Carroll had recognized, an infection of the joint as well as a recurrence of the old osteomyelitic trouble. The joint was full. The man's general condition was very poor indeed; an anemic factory worker to begin with, dragged down by seven weeks' of sepsis. Subnormal temperature at the time.

I got him into town, and into the City Hospital, and operated the next morning, under ether, with a three-inch long incision of the outer side of the knee. There was a large amount of thin pus in the joint, under much pressure. There was no change in the joint surfaces, beyond slight purple tinting and minimal swelling of the synovial membrane. The joint was irrigated in the regular way; 15 minutes of corrosive; then salt. It was then carefully sutured—the capsule with No. 1 chromic gut and the skin with silkworm gut,—then, on account of the presence of pus all about the wound, a sealed dressing was put over it. Then an incision on the inner side of the femur opened up the expected pocket in the condyle, which was cleaned of debris and packed. The wound into the joint leaked some serum, but never any pus, even from the skin wound. There was never any synovial leakage. This wound was protected from outside infection by intermittent sealing until it closed solidly Aug. 23. Cultures taken at operation showed staphylococcus pyogenes albus, both in the pus from the joint and in that from the bone. September 9 under ether, a sequestrum was removed from the internal condyle about $1 \times \frac{1}{2} \times \frac{1}{2}$ inches. October 10, 1912, he went home with a small bone sinus but with free joint motion.

Later reports from Dr. Carroll were of a persistent small bone sinus but never any recurrence of joint symptoms or of effusion in the joint. He never regained full motion, but got serviceable motion; as good motion as he had had before the knee joint infection.

CASE 3. Antoinette De L., aged 6; admitted to the City Hospital June 12, 1914, under Dr. David Scannell, for a septic elbow wound, ten days old. This appeared to clean up and June 22 she went to the out-patient department for dressings. On July 10, 1914, 38 days after the injury, she was readmitted and came to my service. The left elbow was tender and swollen with a granulating wound and

sinus, the sinus obviously going into the joint. There was motion of 30 degrees and there was little or no spasm. (It is remarkable how *little* spasm there is to a septic joint in comparison to a T. B. joint, for instance, owing, I think, to the fact that in sepsis, the process is in the synovial lining, not primarily in the articular ends of the bone). An x-ray showed haziness about the lower end of the humerus; no focal lesion.

July 12, 1914, operation under ether. Oblique incision cutting the triceps tendon; very little thin pus in the joint. Joint surface of humerus showed some loss of cartilage; cartilage of radius and ulna intact. Synovial membrane red, spongy and in spots ragged. Usual irrigation preceded by carbolic 95% to the raw humeral surface; excess cleared, as usual, with alcohol. Then the corrosive and then salt solution and closure of the capsule and suture of the triceps tendon. Then carbolic and alcohol to the outer wound and only two or three loose sutures in the skin. Cultures showed mixed infection with staphylococcus predominating.

July 20 (eight days) deep wound entirely solid; only serum from outer wound. July 24, wounds all healed; 45 degrees of joint motion. Discharged to the out-patient department for exercises. Seen two months later at which time there were very definite adhesions and motion was pretty much gone; child in excellent condition.

In this case the patient came to operation after there was real damage to the synovial joint surfaces, too late to prevent adhesion, but the joint disinfection and the aseptic closure of the joint worked perfectly well, despite the unfavorable conditions.

CASE 4. Lloyd S, aged 9. After a slight trauma, developed a knee-joint effusion with temperature. The joint was tapped and clear fluid obtained; a second tapping showed pus. About a week after the first effusion appeared, the joint was opened on either side and drains introduced. Nine days after this, he came into my hands, the drains still in situ. He had just come on a stretcher from middle Pennsylvania to Boston, and was not in the best of shape, though a healthy boy.

On Aug. 29, 1914, I operated under ether; laid the joint open on either side, excising the granulating wounds, in which the gauze drains lay; washed out as usual with 1-15000 corrosive, and then with salt solution, and sewed up the capsule tight, leaving the outer wound almost wide open.

The temperature fell directly and though a little suppuration developed after five days, near the upper end of the quadriceps pouch, it yielded to camphorphenol dressings promptly.

Restoration of motion to 45 degrees was prompt and painless. Here we met a check and on Nov. 28, 1914, under primary anesthesia, some adhesions were ruptured. From then on restoration of motion was rapid until 90 degrees was reached. Now motion is almost perfect; function as good as new.

In this case, owing to the long duration of infection, there was some loosening of ligamentous capsule, and an apparatus was worn for a time to counteract the tendency to backward subluxation of the tibia. There is now nothing left of this. This patient was shown to the Clinical Surgical Society last fall.

There is some question whether this infection was from within or without. There was, I understand, a bitter local difference in Pennsyl-

FIG. 1. CASE 4.

Taken at his home in the fall of 1914, and sent to me.

vania owing to the allegation that the first man's trocar had infected the joint. Cultures at the first opening are said to have been mixed staphylococcus and streptococcus. When the case came to me, cultures showed straight staphylococcus pyogenes aureus. I believe it to have been a case of hematogenous infection starting near the epiphysis, as in our first case, though in this case, no sequestrum developed. The important point in this case, like the last, is that the joint was an open infected joint, and in this case had been open for nine days with drains in the joint, yet real disinfection was possible.

In February, 1915, Dr. E. H. Place, in charge of our "South Department" infectious hospital, having heard of some of the cases, asked me to see a case there. He told me that, in all the years of his experience there, the South Department had never saved a septic joint, to say nothing of the mortality in these cases, and in convalescence from diphtheria and scarlet fever, such septic joints are not very unusual.

The case to be seen was a post-diphtheritic suppuration of the elbow.

CASE 5. The patient, Isaac G., 8 years old, was admitted for diphtheria Jan. 17, 1915. The infection had attacked the larynx as well as the fauces and tonsils. He was promptly intubed. So far as the K. L. infection went, he did fairly well, but on Feb. 8, 1915, had developed multiple arthralgia with a distinct septic type of temperature.

Feb. 10: temperature remained up; of septic type; fluctuation evident in left wrist; right elbow and right ankle; much pain.

Feb. 11th: I saw him and operated immediately under ether.

The wrist and ankle infections were merely peri-articular abscesses. The right elbow joint was, however, full of thick pus. It was opened on the outer side and irrigated and sewed up, according to my technic, the external wound being left entirely open. The culture was reported as pure staphylococcus aureus.

Feb. 16: The note says—"elbow healing up rapidly; no discharge; no limitation of motion; no discharge from right ankle; temperature gradually approaching normal."

FIG. 2. CASE 4.
Further pictures of Case 4. Taken at the same time as Fig. 1.

Feb. 18: Left hip swollen and tender; temperature slightly elevated.

Feb. 20: Operation on left hip. In this joint the capsule was already "blown-out" and gone. The joint was washed and drained but not handled ac-

cording to my technic; it could not be!

Mar. 7: Periarticular infection at the ankle opened up.

Mar. 20: Elbow wound healed; motion perfectly free.

FIG. 3. CASE 5.
On the left flexion of the disinfected right elbow,—on the right full extension taken about ten weeks after operation.

Apr. 23: Discharged to my ward in the main hospital. At this time the elbow joint was perfectly normal. The infected hip healed and did well but has nothing to do with the especial technic under discussion.

The photographs show the result in the elbow joint.

This was a pyemic joint in a child who was in wretched condition after a long siege of severe diphtheria. The result speaks for itself.

CASE 6. The second case at the South Department was Charles F., a boy of seven years, who entered the hospital Mar. 17, 1915, then being six days ill. He had a rather severe scarlet fever; typical. For ten days nothing happened; then after lysis his temperature came to normal.

Mar. 28: He had paracentesis performed on right ear.

Apr. 5: Left drum punctured.

Apr. 6: Right mastoid infection showed itself.

Apr. 7: Operation on right mastoid.

Apr. 16: Pain complained of in left elbow joint; some swelling and fluctuation; exquisite pain on motion.

Apr. 19: Left elbow joint opened under ether; about three ozs. of thick yellow pus were obtained. Joint dealt with in the usual way, washed and sewed up. Culture showed streptococcus pyogenes.

Apr. 21: Joint showed no signs of infection, though the temperature still remained up. At this time a bronchopneumonia appeared.

Apr. 22: Pain in the right knee; much tenderness to pressure or on motion; swelling and fluctuation.

Apr. 24: Pain on dorsum of right foot.

Apr. 26: Operation on left foot (a periarticular abscess) and on the right knee. From the knee about 5 c.c. of blood-stained pus was obtained; rather thin. The joint was washed and closed in the usual way. Cultures later showed a mixture of staphylo- and streptococci. At the same time an extra-auricular hip abscess was opened. The next day the temperature was normal.

May 2: The left elbow showed a perfectly mobile joint, with the external wound clean, if not aseptic. The right knee showed some recurrence of effusion and a good deal of tenderness to touch, but the process seemed to be essentially extra-articular.

May 11: Pulmonary consolidation appeared; temperature 101; pulse 130 to 170.

May 12: He died.

Autopsy showed not only pneumonia, but a basal meningitis. Both joints were clean; the elbow was as good as new in every way; it was not opened. The knee showed a bit of periarticular infiltration and an ounce or two of effusion; was opened up; fluid clear and proved sterile. Specimens taken from the joint showed no bacteria in the tissues by any of the various stains. The synovial membrane showed a little fibrin at the surface; a little inflammatory reaction; a trifling infiltration with leucocytes; that was all that we could find. Dr. F. B. Mallory kindly went over the slides with me, and expressed it as his estimate of the intensity of the process that it was about what you would expect after an irrigation with corrosive or other moderate irritant.

Here is a case, most unfavorable in every way, unable to handle his finally fatal general infection, yet, despite the feebleness of resistance, two serio-

ously-infected joints were *disinfected* one after another and remained clean. One returned absolutely to normal condition; the other stayed sterile and was nearly normal before his death from meningitis. Unfavorable as the result was, as to survival, no more conclusive test of the efficiency of this method of joint disinfection should be asked for.

Just what happens after disinfection I do not profess to know clearly; the procedure has been one of gradual empiric growth, based purely on the clinical findings. Whether the corrosive used remains behind in sufficient quantity to impair the growth of bacteria, or whether the synovial infections we have so long dreaded are, after all, only *surface* joint infections, at first, is hard to say. I am inclined to the latter view, but dare not omit the corrosive.

Those of us whose lot it has been to watch the horrid progress of drained joints, know that it is not the pus in the joint so much as the pus steadily draining from the joint reservoir into the soft tissues of the thigh, behind and above the knee, for example, that kills. We shall all, I think, incline to the hypothesis that the joint itself is more tolerant to sepsis than the books give it credit for; to believe that surface disinfection plus natural resistance may be the answer. However this may be, the fact seems established that this technic suffices for an efficient sterilization of joints infected not only by the facultative pus organisms, but also by the frankly pyogenic bacteria.

In the milder cases (as in the gonococcus joints) the advantage of an open washing is mainly in the removal of debris that contributes to later adhesions; in the septic cases it seems that actual removal of infection is accomplished.

There is one very definite limitation to this method. I have not yet worked out any scheme for using this scheme in infection of the closed joints, hip, shoulder, ankle, etc.; joints that are non-distensible by pus or irrigation fluid. Given such a joint, the capsule is already gone before we reach the case. Disinfection may be tried, but closure of the joint is not practicable. Such successes as I have had in the treatment of such cases by disinfection with camphophenol dressings have no bearing on the matter in hand.

In such cases there is much to be said for Murphy's formalin glycerin injections, if the process can be diagnosed before capsule rupture; later I can see little chance even for this method. In the distensible joints, on the other hand,—the knee with its quadriceps pouch, the elbow with its subtricipital joint space, pus can collect in considerable amount, enough to be obvious, long before there is tension enough to threaten the integrity of the capsule.

In these cases, I believe the technic here presented to be the procedure of choice. All septic products present, fibrin as well as pus, are removed. The efficiency of disinfection is, I think, pretty well shown. The results are not pretty good only, but nearly always perfect.