and in all cases in which catarrhs of the tube or acute otitis come under our observation so as to prevent a later development of adhesive processes.

(2) Mechanical treatment by air douche, catheter, and bougie, combined with internal use of iodine and extensive application of hot air and steam externally as well as *per tubam*.

(3) Intra-tympanic operations, not only in the most advanced cases, but also where other treatment gives a decided but not lasting relief.

(4) Use of direct pharyngoscopy in order to better detect changes around the ostium pharyngeum of the tube.

Finally, I would like to express the wish that when experimenting new methods of treatment only such cases may be recorded that are absolutely unquestionable as to their diagnosis, and that the records be furnished with the complete functional tests. By this we may reach the position of being able to control the value of different methods on a large and more uniform material. Furthermore, an exact exploration is desirable of the relation existing between the localisation of troubles of the conducting apparatus and the function of hearing. This might lead us to further development of our operative measures.

SOCIETIES' PROCEEDINGS.

BRITISH MEDICAL ASSOCIATION—SECTION OF LARYNGOLOGY, OTOTOLOGY, AND RHINOLOGY.

Meeting at Birmingham, July, 1911.

MR. F. MARSH, Birmingham, *President, in the Chair.*

DISCUSSION ON ADHESIVE PROCESSES IN THE MIDDLE EAR.¹

Dr. ANDREW WYLIE (London) said: The leaders of this debate have covered most of the points in this subject, but I am sorry that Mr. Yearsley has given us no definite form of treatment, and Dr. Frey has perhaps given us too many. Mr. Yearsley has mentioned that he considers otologists must combine with electricians to provide some form of artificial aids to hearing, and I agree that the sooner this is done the better. I wish to-day to bring before your notice the treatment of this condition by hot and also hot medicated air. I have been treating such patients for over four years with considerable success. Since the meeting at Belfast my instrument for administering this treatment has been modified and improved. It has a little chamber where drugs such as benzol,

See p. 626, et seq.

menthol or turpentine can be inserted, and the hot air blown through it and through the Eustachian catheter into the middle ear, or externally through the meatus. One great difficulty with this treatment is that the catheter becomes very hot and has to be withdrawn. I tried ivory and gum-elastic catheters without success, but Meyer & Meltzer have made me a catheter of a vegetable fibre which does not get hot and can be boiled and sterilised. Physicians and surgeons have found that heat in any form relieves and cures thickened conditions in joints and other parts of the body. Otologists also know that patients suffering from chronic adhesive processes improve when sent to reside in a dry climate; therefore surely the application of dry hot air to this thickened membrane and hypertrophied tissue in the middle ear should relieve it. I have notes of over 1000 cases, and 420 of these were cases of post-suppurative deafness (all with a thickened membrane and some with perforations). About one third of these cases have been decidedly improved. I do not say cured, but feel convinced if they will continue having the hot medicated air blown through the Eustachian catheter at intervals the improvement will continue. The amount of heat depends on the patient; when the tympanic membrane becomes red its application should cease. All my cases have been catheterised and Politzerised many months before the hot air was applied. I would suggest that all should try this form of treatment.

Dr. W. JOBSON HORNE (London) reminded his listeners that even in the worst of the cases of deafness under discussion there was a residuum of hearing, and that it was important to give hope, and to encourage the patient to cultivate that residual hearing. Therefore in what appeared to be the most hopeless case one was justified in urging a course of treatment. Otherwise patients were liable to become despondent, and worse than despondent, if the subjective noises became very distressing. It was important also not to lose sight of the general health of the patient, but in prescribing tonics one had to be guided by the blood tension. As regards the treatment of nasal obstruction no one could lay down hard and fast rules about the generality of patients. One indication, however, for nasal treatment was marked relief following the use of the Eustachian catheter; but even should the use of the Eustachian catheter yield a negative result, that was not, in his experience, a contra-indication of nasal treatment being carried out. In all these cases deafness was of much longer duration than the patients realised, therefore prophylaxis and special attention to the naso-pharynx in early life were strongly indicated. It was remarkable and highly suggestive how in some of these cases of chronic dry non-suppurative catarrh the supervention of pus resulted in a marked improvement in hearing. He could call to mind more than one case, but one case in particular, in which suppuration of the ears followed upon influenza in the case of a patient who had been deaf for many years with chronic, dry, non-suppurative catarrh, with the result that considerable credit, quite undeserved, was attached to himself for the improvement in hearing that followed. In fact, it would seem as if a little pus at times was helpful.

Dr. WATSON-WILLIAMS (Bristol) referred to the vascular and lymphatic interconnections between the naso-pharynx, the Eustachian tube and middle ear, and with these one should perhaps include the posterior ends of the inferior turbinals. This interconnection explained the importance of removing the infective lymphoid tissue aggregations in the naso-pharynx we term adenoids, not only when associated with nasal obstruction, but even in the complete absence of nasal symptoms,

provided the middle ear was involved. He felt, too, that when definite mechanical obstruction existed in either one or both nasal passages, their removal in the class of case under consideration was of value, because it tended to get rid of the associated catarrhal conditions. When one had tried and failed to obtain improvement from Politzerisation and other means, and one's thoughts turned to the propriety of removing the drum and malleus, he suggested that it was well first to simply perforate the drum, and if, so long as the perforation remained patent, the hearing power was definitely improved, there was a likelihood that the more complete removal of the drum and malleus would be permanently beneficial. If stapedio-vestibular adhesions were present the removal of a drum could be of little or no use, whereas if they were not present, one could usually obtain improvement from Politzerisation, Eustachian catheterisation, otomassage and so forth, hence the cases calling for removal of the drum and ossicles were few and far between.

Mr. H. F. MOLE (Bristol) had always been sceptical of the use of fibrolysin, and never recommended it, and recently he had seen two cases—one a medical man, who gave himself injections for six months without benefit, and another case carefully tested before the treatment recommended by an aural surgeon in Germany, and three months later, after the treatment, in which no benefit, subjective or objective, had resulted. He asked if it was possible to tell which cases of malleus adherent to the inner wall will be improved in hearing by mobilisation, as he had seen several cases where the malleus had been mobilised and kept freely movable by massage with Siegle's speculum, but in which no improvement in hearing took place.

Dr. W. S. SYME (Glasgow) said: With reference to the treatment of nasal conditions I do not think that the fact of obtaining no improvement by catheterisation should deter us from treating nasal conditions where these exist. Indeed it is a matter of fairly common experience that many of such cases show decided improvement in hearing after appropriate operations on the nose and naso-pharynx. In cases even where there is an amount of hypertrophy of the posterior extremities of the inferior turbinals which does not give rise to nasal obstruction, but which is keeping up some naso-pharyngeal catarrh, this hypertrophy should be removed. The same applies to pathological conditions in the posterior accessory nasal cavities, and with especial force to remains of adenoids. Regarding the length of the piston stroke of the otomasseur, it is my opinion that a longer and stronger stroke should be used after a period of use of the shorter stroke. The increase should be gradual, and of course the condition of the membrane with special reference to the presence of atrophic portions should be carefully observed. The usual electric masseurs do not in my opinion permit of a sufficiently strong stroke. A watch should be kept for phenomena such as vertigo, increase of tinnitus and so on. Fibrolysin I have found useful in post-suppurative changes in the tympanum. The treatment should be prolonged and of course combined with other methods. Probably, however, the fibrous adhesions in ordinary adhesive catarrh do not differ from those found after suppuration. Their formation is probably due to ulceration of the surface of the thickened mucosa. By the apposition of two ulcerated surfaces adhesion is obtained, and ultimately this adhesion undergoes organisation and fibrous formation. It is probable, therefore, that the reason fibrolysin is more valuable in post-suppurative adhesions has to do with the amount of fibrous tissue. A more prolonged use of it might be valuable in ordinary adhesive processes. With regard to climate and altitude in the

treatment of middle-ear adhesive catarrh I have been in the habit of advising a fairly high and dry locality as the best. It is advisable that reference should be made in connection with the deafness of children to the results of medical school inspection with reference to the presence of adenoids and enlarged tonsils. In most of the statistics it will be observed that the proportion of adenoids to enlarged tonsils is absurdly low, and as specialists we know that the proportion is really very much higher. As a result it follows that the tonsils are removed by general practitioners, and the adenoids, to which attention had not been drawn by the medical inspector, and which in most cases are of most importance, and this is especially so with reference to deafness, are left. The results as regards the hearing as well as the general health are, as one would expect, disappointing.

Dr. L. H. PEGLER (London) said he doubted if we knew all that was important to be known about the pathology of non-suppurative middle-ear deafness—that form which could be excluded from otosclerosis in its typical shape, and in which intra-tympanic adhesions were presumed to exist. If true organic changes had taken place in some of the exceptional cases one met with, it was difficult to explain the great relief in the deafness from intra-nasal treatment in such cases. He constantly saw examples, as all did, in which a nasal operation for obstruction restored the hearing apparently by improvement in ventilation and drainage, and on this principle he thought we might make the path for the catheter the criterion. If this could not be effected, or only by causing pain and perhaps bleeding, it seemed advisable that the aurist should by surgical means render the passage of the instrument no longer a matter of inconvenience to the patient. But he would like to refer to a case of curable deafness that had recently occurred in his practice, where catheterism, though done under difficulties, had been satisfactory, but the benefit to the hearing by that procedure insignificant. A gentleman complained principally of deafness in the left ear, so that he could not converse, especially out of doors, with his companion on that side of him. The right ear was slightly affected, but cleared up under inflation. The left-sided nasal obstruction consisted of a contact between the anterior third or more of the inferior turbinal with a horizontal ledge on the septum. The watch was heard just short of contact. Conversation one yard away; bone-conduction not increased. Membrane thin, slightly atrophic. The nasal breathing was not too free, especially at night, though not specially complained of. In May last the anterior portion of the left inferior turbinal was resected under cocaine, and when the patient left the nursing home a day or two later he reported that his hearing was restored, so that he could converse comfortably. An interval then ensued during which the patient left town for the country, but when he returned a month later it was found that the improvement had not been maintained, and the passage in the inferior meatus, which had appeared sufficient at the operation, had not kept clear, owing to redundant tissue formation. A sufficient portion of the stump, therefore, was removed under ethyl chloride, and the air-way permanently cleared, all contact being abolished between septum and turbinal. This happened early in July, and a day or two after this last procedure the normal hearing was once more regained. Hearing for the watch did not exceed half an inch, but conversation in a low voice or clear whisper was heard six yards off with the face turned away. The speaker could not attribute this result to either ventilation, drainage, or change in the lymphatic circulation, but felt inclined to suggest a nervous influence. This could

be effected through Meckel's ganglion, which communicated with the tympanic nerve supply by the facial and glosso-pharyngeal, through the Vidian, and with the inferior turbinal by its palatine branches. Another route would be through the nerve to the tensor tympani from the otic ganglion receiving impulses from the turbinal branches of the maxillary division of the fifth through connection with the mandibular division, joining the former at the Gasserian ganglion. The speaker recalled a somewhat similar result in a young hospital patient, upon whom he had operated for septal deflection and dislocation in two stages, but in whom the aural affection was suppurative otitis media. There might be some analogy between the mechanism in these instances and that of the rapid curative effect of adenoidectomy in the deafness of children, which was not yet understood. We were also familiar with the good effect of operations for nasal obstruction upon asthenopia and referred head pain, which was obviously due to nervous influence. The speaker concluded by extolling the air-reservoir apparatus as an aid in catheterising obstructed Eustachian tubes, especially when the air was medicated. He also heated the compressed air in cold weather by means of an electric coil made for him by the Sanitas Company, but he could not understand, as a pathologist, how hot air could affect genuine fibrous adhesions, and thought some other explanation for the benefit derived from warm and moistened vapour must be sought.

Dr. BRONNER (Bradford) said that most people had off and on attacks of middle-ear catarrh with or without consequent adhesive processes. It was therefore very important that these attacks should be prevented by general treatment. Sleeping in rooms with open windows without adequate covering of the head was a very common cause of nasal catarrh and middle-ear disease. Local hyperæmia was the best method of preventing contraction of the tissues in the middle ear. Hot air by the Eustachian catheter or external meatus, and suction from the external meatus controlled by the speculum were most useful. Elderly people should take iodide of potassium for long periods of time; this not only prevented the formation of connective tissue so common in old people, but regulated the blood-pressure and improved the tinnitus. In many cases the nebuliser (with iodine, menthol) attached to the American compressed air apparatus and the catheter was very useful, as it cured the catarrh of the Eustachian tube and of the middle ear.

Dr. DAN MCKENZIE (London) said that a number of methods of treating these conditions by the induction of hyperæmia were in vogue. The hot-air treatment had already been mentioned. In addition he himself had recently been trying the effect of Bier's treatment by means of an elastic band round the neck and had found improvement in some cases. He had heard several speakers say that they did not try certain methods of treatment (the removal of nasal spurs, etc.) because they did not see how any benefit could arise from them. This, he submitted, was an injudicious mental attitude in the case of a disease the ætiology and pathology of which were not exactly known to us. The wiser plan was to try all the methods proposed without prejudice. He agreed with Dr. Frey with regard to the beneficial effects of coarse vibration applied to the spine or skull in the treatment of tinnitus. He concluded by an appeal for the prompt and early use of artificial aids to hearing in order to prevent or postpone the inevitable atrophy from disuse of the nerve elements in the labyrinth or brain.

Mr. G. SYDENHAM asked if anyone had tried the method of introducing medicaments into the middle ear by means of direct hypodermic

injection through the membrana tympani, which method he thought more exact as regards the certainty of introduction of a definite quantity. He had tried fibrolysin amongst others injected in this way, and thought in a few of the cases that a definite improvement had resulted. In only one case had suppuration followed, and this was due to faulty technique, though, strange to say, in this case the hearing improved.

Mr. HERBERT TILLEY (London) said that his experience as to the probability of cure or relief of deafness in the class of case under discussion depended almost entirely on the result of inflation of the middle ear by the air-bag or by means of the Eustachian catheter. If such means produced an immediate improvement in the hearing, even though slight, then any pathological defects in the nose or naso-pharynx should be dealt with, the general health of the patient cared for, and a good result anticipated. If, on the other hand, no relief of the deafness followed inflation, he much doubted the value of any intra-nasal treatment so far as the ear was concerned, although removal of nasal obstructions or pathological conditions of the naso-pharynx was often productive of much relief to the patient in so far as those regions were concerned. The patients often expressed themselves as "much better," but if we carefully compared the capacity of hearing before and after such operations it would generally be found that the hearing remained the same, especially when the comparison was made weeks or months after the treatment had been carried out. In his experience this result applied to all forms of treatment, *e. g.* injections of sterilised oils or alkaline fluids into the tympanic cavity, electro-massage of the membrane by an air column, the use of the Eustachian bougie, fibrolysin, or surgical intervention. He had had no success with fibrolysin, and thought that great care should be used before otologists made statements as to its value. He himself would like to see an external scar or a Dupuytren's contraction cured by fibrolysin, and then it would be easier to conceive that it might be useful in adhesive conditions in the complicated cavity of the tympanum, where many other factors had to be taken into consideration. He thought the various otological societies would render excellent service if members who had the time would collect some half dozen cases of deafness due to adhesive conditions in the middle ear, allow them to be carefully tested by two or three other members, and then submit such patients to any of the various modes of treatment which had been advised, and finally, after three to four months, have the hearing again tested. By such means we should soon obtain definite results of positive or negative value. He had used the bougie treatment in cases where the Eustachian tube was obstructed, and had come to the conclusion that if the tube was much narrowed and had been thus constricted for six to ten months, no amount of dilation would ensure permanent patency or relieve the accompanying deafness, at any rate for more than a few hours.

Dr. J. GIBB WISHART (Toronto) had been accustomed to apply heated air for a number of years in selected cases, and in many instances with marked benefit. The compressed air was supplied from a tank and not with a hand-bulb, thus more thorough application to the tympanic cavity was secured. As he had experienced the difficulties with the over-heating of the catheter related by Dr. Wylie, he would be glad to give a trial to the fibre catheters recommended. Dr. Wishart had seen the Holmes pharyngoscope demonstrated by its inventor in Atlantic City recently, and the picture obtained of the mouth of the Eustachian tube and its adjacent parts was a revelation. No matter how skilfully the rhinoscopic mirror were used, the picture obtained could never be exact

owing to the relative positions of the canal and the mirror. A knowledge of the exact condition of the mouth of the tube and the fossa of Rosenmüller was essential to a full understanding of the course to be followed in treatment, and it might well be that the treatment would be revolutionised by the further use of the Holmes pharyngoscope. It had one advantage that should be emphasised, in that while it was introduced through the opposite nostril, all instrumentation could be made directly under the control of the eye.

MR. HENRY J. DAVIS (London) said that the evidence in favour of treating nasal abnormalities in cases of deafness, such as that under discussion, was overwhelming. One saw cases every day where hearing was benefited by so doing. If nasal ventilation was not free hearing became impaired, and why should it be otherwise? So simple a remedy as an alkaline spray would improve patients by clearing the nose; they would volunteer the statement themselves. Regarding intra-tympanic fluid injections through the Eustachian catheter, for three years he had made observations on the results of this treatment. He could not go into the matter now, but briefly he would divide these under four heads: (1) Injections of fluid (diluted aqua menth. pip.); (2) injection of alkaline fluid (*lotio alkalina*); (3) injection of acid fluid (lactic acid 5 per cent.); and (4) oily solutions (*aquol*, *benzol-aquol*, *paroleine*, with or without menthol). His conclusions were that the greatest temporary improvement was obtained by No. 2, but that patients would often benefit by the acid solution injections when they did not improve with the others. In oily solution No. 4 improvement was sustained for a longer time, and this is what one would expect, as the oily solutions were not so readily absorbed. If the tympanic cavity was dry conduction was not so perfect as if it was moist. He agreed with Dr. Bronner that iodide of potash was of great value in some cases, not merely because it produced absorption of tissue, but because it made the tympanic cavity moister; to old people gr. iij a day would suffice.

The PRESIDENT, in summing up, thanked the introducers for the comprehensive way they had dealt with the subject. They were specially indebted to Dr. Hugo Frey, of Vienna. There seemed to be general agreement that if improvement was obtained by inflation, any nasal or naso-pharyngeal condition interfering with drainage or aëration should be dealt with, and a number of the speakers were in favour of this procedure even if no improvement followed inflation. All were agreed upon the importance of prophylaxis. It seemed at present that intra-tympanic injections and massage must be looked upon as the mainstay of treatment. Hot air would probably be more extensively employed after the good results reported by Dr. Andrew Wylie and the exhibition of his improved and very portable apparatus. The position of operative treatment remained practically the same—there had been no great advocacy for it, and the results obtained seemed variable. It was generally felt that a more accurate diagnosis and more careful record of cases were desirable before definite indications could be laid down. The members were agreed that more attention should be given by otologists to aids to hearing, and he was glad to hear that the opener of the discussion (Mr. Macleod Yearsley) was making experiments in conjunction with a skilled electrician.

MR. MACLEOD YEARSLEY, in reply, said: I am glad Dr. Frey has not hesitated to deprecate loose diagnosis and insufficient clinical data. It is impossible to be too careful in relating cases in which new treatments have been tried. I am also pleased to hear him say that prophylaxis is

as important as later treatment ; I believe that it is even more important. In reply to Dr. Wylie, I do not think he is justified in saying that I put my foot down on *all* treatment. I think I indicated those methods which are established as of proved efficacy, and dwelt most upon those whose value is doubtful. I am especially interested in Dr. Wylie's hot-air apparatus, but I am doubtful as to the wisdom of using turpentine ; is it not possible that the inflammatory reaction set up may result, later, in further adhesive processes ? In reply to Mr. Mole and one or two others, I have found mobilisation useful. I have only done it in a limited number of cases, which showed the malleus was absolutely fixed, with plus bone-conduction, and when the deafness has not been of long standing. The improvement has been marked. As regards fibrolysin, I recently asked a distinguished general surgeon as to his experiences. He told me that, in Dupuytren's contraction, it was best to divide the adhesions and rub the drug in. This may explain Dr. Sydenham's good results. I think that, as to length of the piston stroke in using otomassage, I have been influenced by continental warnings as to the danger of damage to the stapedio-ineudal joint. I agree with Dr. Syme that a dry, moderately high climate is best suited for these cases ; in the majority of them the neighbourhood of the sea is contra-indicated. I can assure Dr. Pegler that electrical aids are becoming cheaper ; one which has just appeared only costs £2 10s. I may say that I have met with an expert telephonist with whom I am going to work at these instruments after the vacation. Dr. Bronner doubts the effect of small spurs. Surely they act by impairing nasal drainage, so that nasal secretions can set up a catarrhal condition by constant irritation. I would like to suggest to Dr. McKenzie that the domain of the pure otologist depends upon one's idea of the locality of the opening of the Eustachian tube. Some American otologists have asserted that it is at the tip of the nose. The explanation of Dr. Tilley's hypothetical case lies, I think, in the fact that the nasal operation acts in two ways—by improving the air-way and so the general health, and by arresting the process. As regards the electric nasopharyngoscope, I indicated in opening the discussion that I believe it will prove a valuable aid both to diagnosis and treatment.

AUSTRIAN OTOLOGICAL SOCIETY.

June 26, 1911 ; *Monats. f. Ohren.*, year 45, No. 8.

PROF. V. URBANTSCHITSCH *in the Chair.*

Abstract of the Proceedings.

FOUR FURTHER CASES OF COMPLETE UNILATERAL DEAFNESS AFTER SALVARSAN.

BY O. BECK.

CASE I.—A man, aged twenty-one. Definite lesion. Wassermann positive. Examination of the ears showed their condition to be normal. February 16, intra-venous injection of salvarsan. Next day, rigors. Vomiting, headache for two hours. Moderate Herxheimer. On the 19th the sore was healed over. February 24, 0.4 grm. again of salvarsan intra-