declared them to be varieties of the *Filaria palpebralis*, Wilson, 1884; a nematode worm commonly affecting the eye of the horse. These filariae have also been found on cattle in France, Belgium and India.

Leiper summarizing the reports of these cases believes that, according to the description, the parasites differ from filaria worms in several important respects, both as regards their morphology and development. These worms have been recently (1915) grouped by Railliet in one family; the *Thelaziidae*, divided in three genera: *Thelazia*, *Ceratospira* and *Oxyspirura*. All parasites found in the eyes of mammals are now confined to the genus *Thelazia*. The other two genera contain the worms which live in the eyes of birds. In the cases reported above, the parasite was certainly of the *Thelazia* genus; although it is not certain if it was the *T. lacrymalia* or the *T. callipoeda*.

Cabaut has observed two cases of *filaria loa*. In the first the parasite was moving rapidly under the skin of the lower lid, from where it was removed by means of a threaded needle passed around the worm; the thread being immediately tied in order to capture the parasite. A small incision on the skin afforded a way to pull out the filaria. Another worm was seen under the conjunctiva in the same eye some days afterward and partially removed. Examination of the blood in day time demonstrated great quantities of embryos surrounded by a sheath and marked eosinophilia. In the second patient the parasite was also observed under the conjunctiva. Attempts at extraction were only partially

successful, due to the swiftness of the worm.

Pacheco Luna describes some disturbances of vision in patients suffering from a disease known in Guatemala as "Coast erysipelas" and due, according to the researches of Dr. R. Robles, to the infection with a variety of filaria, probably the *Onchocerca volvulus*, Leuckart, 1893. The doubts still existing about a definite classification are due in part, to the fact that none of the European authors mention among the symptoms produced by this parasite the ocular disturbances, which have been the most important in the author's cases.

These disturbances are: photophobia and diminished vision as subjective signs, and objectively a keratitis especially marked in the part disclosed in the palpebral fissure, and consisting of small dot-like whitish, superficial infiltrations similar to the keratitis superficialis punctata of Fuchs, and the central subepithelial keratitis of Adler. There is no conjunctival or ciliary injection. Iris normal. These lesions run a chronic course lasting for years and when recovery is to be made they diminish, but do not disappear completely, leaving permanent infiltrations.

If the disease goes on unchecked the iris becomes involved and a chronic, insidious iritis sets in, with contraction of the pupil and great diminution of sight. At last the dots become more abundant in the lower half of the cornea and form a uniform, diffuse infiltration giving this membrane the appearance of ground glass. The pupil is stretched downward and becomes irregular and occluded.

HYGIENE.

JOHN A. McCaw, M. D.

DENVER, COLORADO.

This section deals with the hygienic aspects and prophylaxis of conditions and diseases, the other aspects of which are dealt with in preceding sections. It covers the literature from January, 1917, to October, 1918.

LIGHTING IN ITS RELATION TO THE EYE.—In Ferree's work the object has been to compare the effect of different lighting conditions on the eye, and to

find the factors in a given lighting situation which cause the eye to lose in efficiency and to experience discomfort. Tests were made to determine the

eye's aggregate loss in functional activity and to analyse this effect.

Powell discusses the color of artificial light, and its relation to the medical profession; and concludes that the Mazda C-2 lamp gives light that approximates the sunlight. This result is obtained by a blue glass globe that permits the uniform transmission of the various colors of light. The lamps are standard and are listed in sizes from 75 to 500 Watts or from 60 candle power to 550 candle power. This lamp is particularly applicable in the chemical laboratory: In microscopic work, in ophthalmology and in examination of X-ray plates. **Luckiesh** writes on color in lighting and investigation of diffusing glassware; and **Worthing** on rating lamps as to filament brightness with fluctuation.

Kerr says that the maximum brightness which the eye is prepared for, is the brightest sky, apart from the actual image of the sun. This amount is generally taken as 2.5 to 3 foot candles to the square inch, and some such amount has been suggested as the standard for any visible source of lighting. **Kerr** thinks that the indirect system of lighting is not advantageous where small objects or textural surfaces have to be examined. It would be good in a bank, but bad in a school.

Direct lighting is the most economical method for handling objects or manufacturing processes. But the light source must be kept out of the field of vision, and shaded down to 3 candle power per inch for any visible surface, or it will cause strain and fatigue. Excessive or defective contrast may be equally troublesome. The New York Commission standard of 1 to 20, and 1-100, requires further investigation. Flicker is more evident to the peripheral retina than to the macula, and is due to alteration in light and shade slow enough to invoke continual efforts at adjustment. It is in this, perhaps, that pupillary contractions are most to be regarded.

Prosser holds the opinion that the color of the surroundings has an enormous effect on the recovery of the sick, especially those who are depressed

from one reason or another, or are suffering from shell shock. To remove from the patient the idea of being shut in is of utmost importance. So by a color scheme for the ward he endeavors to give the impression of out of doors. The effects are said to be beneficial.

Marks enumerates the advantages of good light in factories and the present status of factory light (especially quoting from reports of British Committee on Health of Munition Workers). His summary of general requirements for good illumination in factories is as follows:

(1) Sufficient illumination should be provided for each workman irrespective of his position on the working space. (2) The lamps should be properly selected and so installed as to avoid or minimize strain on the eyes of the workman. The type of lamp should be adapted to height of ceiling and closeness of work. (3) The lamps should be operated from sources of supply which will insure continuity of service and steadiness of light. (4) Adequate illumination should be provided from overhead lamps so that sharp shadows may be prevented as much as possible. (5) In addition to the illumination provided by overhead lamps, individual lamps should be placed close to the work if they are absolutely necessary, and in such cases the lamps should be provided with suitable opaque reflectors. The lamps should be placed if possible without the field of vision.

INJURIES OF EYE BY LIGHT AND OTHER RADIATIONS.—**Burge's** conclusions on the injurious effect of ultra-violet radiation on living tissue are: That ultra-violet radiation kills living cells and tissues by changing the protoplasm to form an insoluble compound. The effective region of the spectrum in changing the living material of the cells lies between 254 $\mu\mu$ and 302 $\mu\mu$. An opacity of the lens can be produced in fish living in solutions of calcium chlorid or calcium nitrat, by exposing the eye of the fish to radiations from a quartz mercury vapor lamp. Abnormal quantities of the salts of calcium and sodium silicat in the cells of

the eyelids and of the cornea increase the effectiveness of ultraviolet radiation in producing anterior eye trouble.

Verhoeff and Bell in their work on the "Pathological Effects of Radiant Energy on the Eye" conclude as follows: "Abiotic action for living tissues is confined to wave lengths shorter than 305μ , at which length abiotic effects are evanescent, while for shorter wave lengths they increase with considerable rapidity.

Liminal exposure capable of producing photophthalmia to the extent of conjunctivitis, accompanied by stippling of the cornea is in terms of energy 2×10^6 erg-seconds per square centimeter of abiotic radiation, of the character derived from the quartz lamp or the magnetite arc. About 2 1-2 times this exposure is required to produce loss of corneal epithelium.

The abiotic action on the cornea and conjunctiva produced by any radiating source follows the law of inverse squares, and is directly proportional to the total abiotic energy received. After exposure of the eye to abiotic radiations, there is a latent period before any effects, clinical or histologic, become perceptible.

The combined effect of repeated exposure to abiotic radiations is equivalent to that of a continuous exposure of the same total length, provided the intermissions are not long enough to establish reparative effects. Actual abiotic damage to the external eye renders it temporarily more sensitive to abiotic action.

SNOW-BLINDNESS.—Daland's conclusions after a visit among Eskimos are:

- (1) The Eskimo possesses no immunity from snow-blindness.
- (2) Snow-blindness occurs on cloudy days or dark days as well as on sunny days.
- (3) One attack of snow-blindness predisposes to another.
- (4) Snow-blindness occurs in animals. Ross saw snow-blindness in a bear in the arctic region.
- (5) Eskimos show great cleverness in making goggles for prevention of snow-blindness.
- (6) There is association of conjunctivitis and corneal erosions with chorioretinitis in snow-blindness.
- (7) The ultraviolet

ray causes conjunctivitis, corneal erosions, and possibly chorioretinitis.

MOVING PICTURES.—Berry made a special study on the relation of the moving picture to defective eye-sight. He states; that with proper illumination of the auditorium and proper projection on the screen, after good mechanical preparation of the pictures, there is small hazard to eye-sight.

Kerr thinks constant fixed attention with the eyes in a dark adapted condition is unusual and soon gives rise to strain. The practice of remaining in the hall and seeing the same series of films over and over again should not be allowed, especially by children. The question of the length of such entertainment for children is of importance. The best position to view the pictures from is between 20 and 30 ft. from the screen, on a level with the center. If viewed from nearer than 20 feet or from below or the side futile efforts at accommodation are induced. The amount of flicker depends a good deal on the skill of the operator and the quality of the film.

Wilson has pointed out that an investigation carried out in Glasgow schools by H. Wright Thompson, in which 50,000 children were examined, showed that while the proportion of real ocular defects was fairly constant, the proportion of children with defective vision varied from 53 per cent in the crowded and poorer parts of the city to 20 per cent in the outskirts where the children enjoy good hygienic conditions. The patients attending Harman's clinic were of a respectable artisan class and lived under good hygienic conditions. The only novel factor to account for the increased number of children with defective vision without any organic ocular defect seemed to be the cinematograph.

Pollock believes, that with frequent attendance at moving picture shows, children from 4 to 8 years of age develop a tendency to convergent squint, in the absence of errors of refraction; and in older children it causes congestion of the optic nerve and eye strain.

Bahn's conclusions are that moving pictures under favorable conditions do

not cause as much fatigue as the same period of concentrated reading. Most persons who complain that moving pictures cause eye discomfort have some ocular defect. Under favorable conditions the pictures act as a test of distant eye endurance.

With regard to the lighting of moving picture houses, Kerr makes these suggestions: (1) That the standard of illumination of the screen be fixed at a minimum of 1 foot candle. (2) That the material of the screen should be good matt-white screen. (3) In illumination of the theater, extreme contrasts between the screen and the general surroundings should be avoided. The ratio of 1:100 would imply brightness of surroundings of not less than 0.01 to 0.02 foot candles. For illumination of the seats 0.02 to 0.05 foot candles has been suggested. Any lights indicating exits should be screened so that their brightness does not exceed 3 c. p. per square inch.

EFFECTS OF GLARE.—Kerr states that *glare* is more important than abnormal refraction or the provision of spectacles. He points out that eye strain due to its influence becomes increasingly obtrusive after the twelfth year: It is especially insistent in reading, writing, and figuring; and is particularly distressing under artificial illumination. He urges as an absolute rule that no naked filament, mantle or flame should be permitted; nor any source of light with greater intrinsic brilliancy than three foot candles per square inch. The influence of paper in producing glare is explained. The blackboard, as a source of glare, comes in for attention.

ILLUMINATION FOR FINE HAND WORK.—Pol thinks that the practical conclusion is that it is impossible to set up a definite standard for minimum illumination; the individual conditions vary so widely. If a minimum is to be selected the tests described point to 288 candle power meters, for vision of 0.8 to 0.5.

EYE STRAIN AND HIGHER EDUCATION.—Heard presents the subject of hygiene of the eyes in relation to the school life of the child, and carries it

up to the field of higher education in college and university.

CLASSES ON THE CONSERVATION OF VISION.—Irvin reports that the children assigned to this class are those whose eye-sight is so defective as to make continuous use of the book type of ordinary size inadvisable; and whose distant vision with glasses is more than 6/60 and less than 6/15. Children with high refractive errors, which are likely to be progressive unless all eye strain is removed, are also regarded as suitable candidates tho their vision exceed 6/15. The aims are first to instruct the pupils with a minimum of eye strain; second to conserve the vision they possess; third to provide such vocational guidance, and if necessary vocational training, as will enable them to fill the most useful places in the community their powers will permit.

OCULAR WORK AND REST IN CHILDHOOD.—Cutler states in the introduction to his paper that it is not based on data, contains no narration of cases or of results, and deals with matter essentially commonplace. But he speaks of children's eyes in relation to their work, with especial reference to certain tendencies, the early recognition of which is vital to normal development.

UNUSUAL DANGERS.—Jackson in an editorial comment on the great number of persons blinded in the Halifax disaster, points out the dangers in certain emergencies of standing facing window glass. Sudden explosion, or wind may produce injury leading to blindness. He also points out the great danger there is in the free use of the air-gun in the hands of children.

PROTECTIVE GLASSES.—Greeff describes two substances, *cellon* and *triplex glass*, which have proved useful in the manufacture of war glasses. Cellon is a derivative of cellulose and related to celluloid; but is not inflammable and not affected by acids or acrid gases. It is used in very thin plates in sand glasses, gas masks, airships, etc. Triplex glass consists of two thin plates of plate glass, with a layer of cellulose between, which prevents wounds of the skin and eye when they

are splintered. The official military snow spectacles, which have proved useful, are made of simple smoky gray glass with a lateral wire sieve attachment. For gas masks, plates of cellon are mostly used.

Higgins recommends triplex glass for spectacles, as a substance that will not break and splinter. He advises every one who has to wear glasses for distance, especially those who shoot, to have them made of triplex glass. Such glasses guard against injuries from small foreign bodies, such as shot; and so far as he is able to judge, there is no difference in refraction between triplex and ordinary glass. **Giuseppi's** iron goggles having slits, transverse and slanting, permit ample vision. A circular piece of woolen goods is fastened around the part that serves as a lens. This absorbs sweat dripping from the forehead.

Cruise examined 320 eyes injured in war, and comes to the conclusion that slightly over 50% of injuries to the eyes in war are preventable. His percentage agrees with the findings of **Morax** and **Moreau**. Both **Cruise** and **Morax** agree that some sort of visor or shield would prevent these injuries. **Cruise** invented a steel mesh visor that fastens to the helmet for this purpose.

Walter, with large experience along beaches and roads of Florida, finds that amethyst tinted **Crookes'** and noviol glasses are better than amber tinted ones. The lightest shade which gives relief from irritation is always given.

Richardson evolved the idea of making use of the *stenopeic slit* in spectacles to wear in the movies. He found them very beneficial. They have been tried and found highly efficient for other purposes than just the movies. They have been tried by pilots and by the captains of several large liners, and they have found that in moonlight or on a sunny day they can see farther with the spectacles than they could without them. The United States Government has made a number of tests, on the firing line and in the aviation field, and the efficiency of the marksman is greatly improved by the use of these spectacles.

Kirkpatrick, to protect the eye after cataract extraction, has a special form of goggles made of aluminum. The shield is made to fit the orbital margins. The two shields are held together by a piece of tape which serves as a bridge and is kept in position by a tape passed round the head above the ears. Each has a large opening in it. For the eye operated on this opening contains a lens of amber glass; the other opening is left free. This sort of protection has been substituted for the dressing the day after operation.

The different types of goggles for protecting the eyes in industrial processes, **Luckiesh** enumerates as:

Goggles for protection against flying materials; Goggles for protection against gases, fumes, and liquids; Dust goggles. "The goggle has the greatest possibilities for the saving of the sight of men engaged in hazardous industrial occupations."

"The requirements of goggles for protection in these operations are that they shall provide sufficient reduction of the intensity of the light, be effective in absorbing the ultraviolet rays, keep the eye cool, and transmit as much as possible of the visible spectrum without color distortion."

Valuable data on this subject are given in a recent publication entitled, "Glasses for Protecting the Eyes from Injurious Radiations," by **Coblentz** and **Emerson**. "The data given by these investigators show that of the infrared rays emitted by a furnace heated to 1,000° to 1,100° C., (1) 99% are obstructed by gold plated glasses; (2) about 95%, by sage-green or blue-green glasses; (3) about 60 to 80% by very deep black glasses, and (4) about 60% by greenish-yellow glasses. At higher temperatures these data would be somewhat different."

The glass, which, in the opinion of **Crookes**, provides most satisfying protection is sage-green in color, which in a plate 2 mm. in thickness, is opaque to 98% of the heat radiation, absorbs the ultraviolet rays beyond any possibility of their proving objectionable; and at the same time transmits 27.6% of the incident light.