*The Principles of Crop Production.*LECTURE DELIVERED BEFORE THE CHEMICAL SOCIETY ON
NOVEMBER 18TH, 1915.

By EDWARD JOHN RUSSELL.

IN any discussion of the principles of crop production it is necessary to begin with the year 1840. By that time it was definitely known that plants consist mainly of organic matter along with a little mineral matter—phosphorus, calcium, potassium, sodium, etc.—to which, however, very little importance was attached. The practical man knew that farmyard manure was the great fertiliser; he also knew that other substances, bones, salt, etc., had, in certain circumstances, considerable fertilising value. The most obvious facts were the large amount of organic matter in the plant and the large amount of organic matter in the best manures; and it is only natural that chemists and physiologists should have connected these, and argued that the object of the manure was to furnish organic matter for the plant.

By a brilliant stroke Liebig, in 1840, brushed aside this obvious connexion and declared that the true function of the manure was to provide, not organic matter, but the mineral constituents which the chemists had ignored. The first step, he said, was to find out what mineral constituents the plant contains, and then to supply these substances in a suitable form. If any one of them is lacking the soil is rendered infertile, and matters will not be put right until that one is added. Thus the whole art of manuring was reduced to an exact science. Liebig was a brilliant writer and secured many disciples. Farmers were told that they need only use a few pounds of simple salts to make their crops grow. It was the first time they had heard the fairy tales of science. We can only faintly realise the utter incredulity with which they received the idea that crops could be raised by small doses of chemicals sold in bags from a factory, in place of the cartloads of farmyard manure that had always been regarded as necessary and sufficient. It was as if farmers had been told that they themselves could dispense with the beef, bread, and beer that Cobbett had said was their Divinely prescribed diet, and could live instead on tablets of chemicals.

The comments of the local wits were pretty caustic; some of them are still preserved in the countryside, but they are hardly suitable for the polite audience that I have the honour of addressing to-night.

To make matters worse, the prescriptions failed in practice, and the downfall of the first manurial principle seemed complete. The Rothamsted experiments showed that Liebig's ash constituents

FIG. 1.

Yield of wheat on the Broadbalk plots, average of sixty-one years' results.

When nitrogen compounds are withheld the yield is little better than on the unmanured plot. Complete artificial fertilisers give a full crop which is fully maintained to the present time, and in this case is better than that given by farmyard manure.

Broadbalk Wheat. 61 years (1852-1912)

gave little better crops than no manure at all. Liebig had left something out; it was necessary to add nitrogen as well before complete growth could be obtained.

The critics urged that the effects would only be temporary; that in time the land supplied with "artificials" would give out. Ex-

perience has shown that this is not so; similarly good results have been obtained at Rothamsted over the long period of more than sixty years (Fig. 1).

Part, therefore, of Liebig's principle is perfectly correct: the mineral constituents are indispensable and must be supplied to the plant. The mistake was to suppose that they were sufficient. We may take it as established that crops can be grown satisfactorily and indefinitely by supplying proper quantities of suitable compounds of nitrogen, phosphorus, and potassium. This we can call our first principle. Difficulties arise, however, directly one tries to develop it in practice. Trouble began with the attempts to find out what are suitable quantities to use. Liebig had supposed that the requirements of a crop could be gauged by the composition of the ash. Lawes and Gilbert showed that this was not the case. Thus the ash of the turnip crop contains a considerable amount of potash but only little phosphate; according to Liebig, it should have required mainly a potassic fertiliser. Lawes and Gilbert showed, however, that it required phosphates and not potash, and they concluded that the special requirements of a crop could only be discovered by actual trial.

This view was developed in the 'sixties in a series of brilliant lectures by Ville. After numerous experiments (he says "many thousands"), he drew up the following list, showing the special need, or, as he called it, "the dominant," for each crop:

Ville's List of Dominants.

Nitrogen	for	Cereals.
"	"	Beetroot.
Potash	"	Potatoes.
"	"	Vines.
Calcium phosphate	"	Cane-sugar
No dominant	"	Flax.

In order to ascertain the special needs of the crop on a particular soil, he grew the plants on a series of plots, one of which was given the complete manure, whilst the others each had one constituent left out. Thus for wheat he obtained the following results, and, therefore, concluded that on this soil wheat requires a good supply of nitrogen, less phosphorus, and still less potassium:

	Crop per acre. Bushels.
Normal manure	43
Manure without lime.....	41
" " potash	31
" " phosphate.....	26½
" " nitrogen	14
Soil without manure	12

The method, of course, is perfectly sound, and it has been very widely adopted. It is, however, frankly empirical, and empirical work is never very inspiring, so that for a long time soil work came rather to a standstill.

It has several times happened in the history of agricultural chemistry that the new illuminating idea wanted to revivify the subject at a stagnant period has come in from some outside technical problem that had to be solved. So it was here. The growth of the towns and of stricter ideas on public health had brought into prominence the need for better sewage purification, and it was imperative that the problem should be dealt with somehow or other.

Schloesing and Müntz found that satisfactory purification involved the conversion of ammonia into nitrate, which could be attained by allowing the sewage to trickle very slowly through a tube filled with sand or limestone. There was nothing specially novel in this: the change of ammonia to nitrate in the soil was a well-known phenomenon. It had received a special name, nitrification; some people thought it chemical, others physical, but few troubled themselves about it, and it passed into the category of Things Accepted. The remarkable feature about the investigation of Schloesing and Müntz was the beautiful way in which they developed an observation that most people would have passed over. The conversion of ammonia into nitrate did not set in directly the sewage was allowed to trickle down the column; an interval of twenty days was necessary before it began. Why was this delay? They argued that the process could not be purely chemical or purely physical, otherwise it ought to set in at once. The only explanation seemed to be that it was brought about by living organisms which required time to establish themselves and multiply before setting about their work. If this hypothesis were correct, they argued, the whole process ought to be brought to a standstill by adding a little chloroform—and this was found to be the case. Further, it should start when a little extract of soil was added—this also happened.

This piece of work was brilliantly followed up, and led to most unexpected results. It was seen at once that the soil was not a mere inert mass, but that it was teeming with life and pulsating with change. The number of bacteria is enormous, running into millions per gram, and the question is raised: How do these organisms live? They must have food, and they must have energy. We are therefore forced to go back to the soil and study it as a medium for the life of a soil population.

A very cursory examination shows that the soil forms only a thin layer; underneath it lies the subsoil, which is wholly different

in colour, texture, and especially in its behaviour towards the plant (Fig. 2).

Yet there was not always this difference. When the soil was first laid down it was all like the subsoil, and whenever a new surface becomes exposed, either by landslips, cliff-falls, etc., it is always the subsoil type that appears. The first vegetation has no great supply of plant nutrients, but plants suited to the conditions nevertheless spring up. They take what they can from the crude soil, they take carbon dioxide from the air, they synthesise sugars, starches, cellulose, proteins, etc., deriving the necessary energy from sunlight. When the plants die they fall back on the soil and return to it all that they took, and a good deal more of new material besides. That introduces a fundamental change.

The new material thus added contains stores of energy and food substances suitable for the bacterial population, which forthwith flourishes. Decomposition goes on, nitrates and other substances are produced, and the conditions are made more favourable for the growth of a new race of plants. One of the most obvious changes is the formation of nitrates, but other products are formed as well. It is proving exceedingly difficult to trace out full details, but the following is in the main accurate:

Unfortunately, not much is known about the details, but the reaction is extremely important. The initial products are of little value to the crop or the soil. The final products are invaluable for plant nutrition, and some of the intermediate products are very valuable for the soil. This, therefore, is the reaction on which plant nutrition depends, and it is of the highest importance that it should proceed rapidly and smoothly. Where for any reason it does not, the soil becomes unproductive. This is usually the case on waste lands, commons, heaths, &c. In the Bagshot heaths, for example, there is too little water and too much acid for successful bacterial action, and decomposition does not proceed sufficiently

FIG. 2.

Soil

Sand

Subsoil

Plants grown in soil, sand, and subsoil respectively, all without manure, showing the marked differences in behaviour towards the growing plant.

FIG. 6.

(See page 1848.)

Pot No.

47

55

63

72

79

Tomatoes supplied with increasing doses of manure.

Pot 47.—No manure.

Pots 55 to 79.—Increasing dressings of manure. This increases the amount of growth up to pot 72, but it depresses growth in pot 79, where too much is given. The middle pot, 63, is best for fruit.

[To face page 1842.]

quickly. The land is simple waste and produces no agricultural crops.

Scientific crop production depends largely on controlling this reaction. Three things are necessary: the conditions—the air supply, water, temperature, etc.—must be favourable; the organisms must be of the right kind; and the supply of raw material—plant residues—must be kept up.

We shall see later on how the favourable conditions are obtained. Hitherto little has been done to control the organisms beyond improving the conditions, but beginnings have been made in the direction of inoculation and partial sterilisation. The supply of raw material is kept up in several ways; probably the oldest is to leave the ground alone, so that it covers itself with wild vegetation, which is then ploughed in. This formed part of the Mosaic Law*; it was the regular mediæval custom in our own country, and it is practised to this day in Connemara. It is too haphazard for modern use, however, and so nowadays the farmer grows a special crop with the express intention of ploughing in all or part of it. Clover, or a mixture of clover and rye grass, is very common, and has the advantage that some can be cut for hay and the rest ploughed in, and the further advantage that the clover plant is associated with one of the nitrogen-fixing organisms.

Fig. 3 shows the beneficial results of ploughing in clover on the Agdell field at Rothamsted.

Other crops instead of clover can be used; for example, mustard, lupines, vetches. A considerable degree of control is obviously possible, because different plants not only yield different amounts of organic matter, but also contain somewhat different compounds.

It is not necessary that the residues should be ploughed in as such. The plants may be fed to animals, and the animal excretions—which represent the parts the animal did not retain or convert into gas—can be ploughed in in the form of farmyard manure.

The second broad principle of crop production is, then, that the biochemical decompositions in the soil must proceed smoothly and rapidly.

New difficulties arise as soon as one begins to develop this principle. Reverting to the scheme on p. 1842, it is seen that the decomposition of proteins may proceed in two ways, either ending with nitrate or with nitrogen. Now the nitrate ending is desir-

* Leviticus, xxv., 3—5.—“Six years thou shalt sow thy field, and six years thou shalt prune thy vineyard, and gather in the fruit thereof; but in the seventh year . . . thou shalt neither sow thy field, nor prune thy vineyard. That which groweth of its own accord of thy harvest thou shalt not reap . . . : for it is a year of rest unto the land.”

1844 RUSSELL: THE PRINCIPLES OF CROP PRODUCTION.

able enough, but the nitrogen ending is highly undesirable. Yet this happens directly the process is speeded up too much. The more intense the cultivation becomes the more serious are these losses; they are bad on the prairies, but still worse under conditions of intense cultivation. To some extent this is inevitable; it is equally true of engines, but just as the engineer has increased the efficiency of engines, so the agricultural chemist has to increase the efficiency of the nitrogen utilisation processes.

FIG. 3.

Comparative effect of clover or bare fallow on the succeeding crops in the rotation.

Total produce in 100 lb. for clover, wheat and barley, and in 100 cwt. for roots.

The difficulty of investigation arises from the fact that these losses, whilst considerable on an acre of land, are very small in the quantities of soil that one uses in laboratory experiments, only a few c.c. of nitrogen being given off per 100 grams of soil in a week. The way round a difficulty of this sort is to find a parallel case where the loss of nitrogen is considerably greater, and this is furnished by the manure heap.

Here we are dealing with the same kind of action: the hydrolysis

of protein effected by micro-organisms. A very interesting result has been obtained.

After several months' storage under unaerobic conditions the whole of the nitrogen is recovered, although there has been some transformation of protein and amides to ammonia. Directly air is admitted, that is, under aerobic conditions, there is a loss which is apparently not ammonia, and can only be attributed to gaseous

FIG. 4.

Showing the amounts of nitrogen present as ammonia, as amide, and in other forms (for example, protein) in farmyard manure stored under different conditions. The lengths of the columns compared with the first show the various losses and transformations.

Nitrogen changes in farmyard manure

nitrogen. Still greater losses set in when we began with anaerobic conditions and follow on with aerobic conditions (Fig. 4).

This experiment was done in the laboratory with farmyard manure kept in carefully closed vessels, but we have been able to show that similar results are obtained with soil and also with ordinary manure heaps. Here, therefore, we have reproduced the reaction by which these losses arise. In agricultural practice the losses are aggravated by leaching and become very serious,

1846 RUSSELL: THE PRINCIPLES OF CROP PRODUCTION.

amounting at the lowest estimate to several million pounds sterling per annum in manure alone, quite apart from the soil. Leaching troubles can be obviated, but this loss by gaseous nitrogen cannot be dealt with until we know just what is happening. The simplest hypothesis is that something formed under anaerobic conditions reacts with something formed under aerobic conditions, giving rise to gaseous nitrogen.

Unfortunately, the purely chemical work on the decomposition of protein has not gone far enough to enable a full working hypothesis to be mapped out. The decomposition does not stop at amino-acids; under bacterial action there is a further change to bases and acids. These are under investigation in several chemical laboratories, but the results have not yet helped us much. Here, therefore, we have an economic problem of the first importance waiting for the solution of a chemical problem which, at first sight, seems rather academic and remote from practice.

These biochemical changes, important as they are in crop production, do not end the matter. The soil comes directly into the reaction; it is not simply the place where the reaction takes place—the beaker, so to speak, where plant nutrients are formed. Some of its mineral constituents come into play. The calcium carbonate neutralises acids produced during the decomposition. The clay and some of the other constituents possess colloidal properties, so that all these reactions proceed on a jelly-like surface and not in a fluid medium, and they are liable to be affected by all the complications produced by surface actions.

The plant plays an even more active part. Its roots absorb some of the products—the nitrates, the phosphates, etc.—and might therefore be expected to hasten the whole process; but this does not happen. On the contrary, the growing plant appears to retard it, and nitrate is always formed in higher quantities on fallow than on cropped land, even after allowing for what is taken by the crop:

	In 1911		In 1912	
	Fallow land.	Cropped land.	Fallow land.	Cropped land.
N as nitrate in top 18" of soil, June	58·7	15·3	46·1	12·8
Nitrogen in crop	—	22·6	—	6·1
Total.....	58·7	37·9	46·1	18·9
Deficit in cropped land	—	15·8	—	27·2

Whether the growing plant affects the nature of the change or only the rate is not yet known. The essential point is that, so far as plant nutrients are concerned, neither the soil nor the plant plays an entirely passive part. The soil is not an inert medium, and the plant is not a mere passive bucket into which the products

of the reaction are drawn; each plays an active part, disturbing both the reaction and the distribution of the products.

That leads us to the growing crop. Just as the recognition of the living agencies in the soil altered our whole conception of the soil itself, so the recognition that the plant is a living thing has broadened our conception of the factors necessary for plant growth. In much of the literature of the 'seventies and 'eighties it is tacitly assumed that the whole art of crop production is a question of manuring. Agricultural chemists, in short, had got rather into a groove. When this had happened before, they had been jerked out by the necessity of solving a sewage problem. On this occasion the new light came again from the outside, but this time from a study of the quality of American cigars. Whitney

FIG. 5.

Showing the relationship between the composition of the soil and the quality of the crop. Soils containing small amounts of clay produced the finest qualities of tobacco ("wrapper" tobacco); as the amount of clay increases, the quality falls off, although the crop becomes larger.

Cigar Tobacco.

was then with the United States Weather Bureau, and naturally came across many instances of the striking effects of climate on both the quality and quantity of crops. It is a commonplace among growers that high quality is often associated with certain climatic conditions. He recognised this, and further observed that in the tobacco-growing States good quality tobacco could be produced on certain soils but not on others, no matter what scheme of manuring was adopted, and he brought this observation into line with his Weather Bureau work by supposing the climate of these soils to be that required for the production of high quality. By "climate of the soil" he meant temperature, moisture content, air supply, etc., and in a very ingenious paper he shows that these factors can be correlated with the size of the soil particles (Fig. 5).

This led to the recognition that the type of the soil is an im-

portant factor in crop production, which has had some extremely interesting developments. Mechanical analyses to determine the type became an indispensable part of the routine of soil analysis. The perspective was restored, the fact emphasised that the plant not only wants food, but also proper water supply, air supply, and temperature.

Now there is a very simple rule that applies to all these factors. Plant growth increases with increasing supply of any one of them, but this only happens so long as the supply of every other factor is adequate. When anything is lacking the increase in growth is not kept up, and additional supplies give no extra crop. Finally a stage is reached when extra supplies may do harm, either by direct

FIG. 7.

Increases in amounts of the various factors necessary for plant growth do not cause indefinite increases in growth. After a time some other factor becomes insufficient and operates as a limiting factor.

General Relation between any particular Factor and Plant Growth.

injury or by cutting out another indispensable substance. This is shown in the tomato experiments of Fig. 6 (page 1482), where successively increasing doses of sodium nitrate are applied in the four pots 55, 63, 72, and 79, although no further growth is obtained in 72 and 79 because of the insufficient water supply. The conditions for favourable growth are all present in pots 72 and 79 excepting only this one, but it effectually prevents the plant from taking full advantage of the good conditions. All this is expressed in a generalised form in the curve of Fig. 7, which thus represents our third principle of crop production. It has only recently been revived in agriculture, although the fundamental idea is old; it can be found in the writings of the political economists of the Malthusian school; it was used in a special form by Liebig in his

"Law of the Minimum"; it was developed by Horace Brown and F. F. Blackman. In its full generalised form it is proving extremely useful.

Problems of soil fertility generally have to be approached from this point of view. Whenever a case of infertility has to be studied the first question to settle is: What is the limiting factor? And the next: How can this limiting factor be put out of action? As a rule the limiting factor is one of the following:

Limiting factor.	Put out of action by :
Wetness	Drainage, liming
Dryness	Irrigation, suitable cultivations
	Addition of organic matter
Lack of temperature	Drainage and cultivation
Sourness or acidity	Liming or chalking

The removal, although simple in principle, may be very difficult in practice: it has often proved to be the rock on which many beautiful schemes for increasing food production have been wrecked. The laboratory experiment may show conclusively that a certain treatment increases plant growth. Every other condition except the one under investigation is eliminated by making it as perfect as can be, and the result may be thoroughly sound from the physiological point of view. It is, however, wholly unjustifiable to assert that such treatment will increase the growth of crops in the field, because one cannot, in the field, eliminate all limiting factors and make every other condition as good as in the laboratory. Even if one succeeds with the soil—which one can very rarely do—there still remains the weather, which may always confound the calculations.

We have seen that, broadly speaking, three general principles of crop production can be laid down:

1. The plant must have a sufficient supply of all necessary nutrients, especially of nitrogen, potassium, and phosphorus.

2. The biochemical decompositions in the soil must proceed smoothly and quickly.

3. All the requirements of the plant must be satisfied. Any one left unsatisfied constitutes a limiting factor preventing further growth. Increases in any one factor give increases in growth until something else proves insufficient and becomes a limiting factor.

No doubt future investigators will find other principles besides these three. It has been suggested that besides the nutrients proper there are certain directive agents akin to the remarkable substances recently investigated by animal physiologists. Armstrong was the first to direct the attention of plant physiologists to the effect of hormones in regulating the vital processes of the plant,

1850 RUSSELL: THE PRINCIPLES OF CROP PRODUCTION.

and the idea is proving very helpful in forming a mental picture of what goes on during growth. It has aroused considerable interest among agricultural chemists, because it suggests the possibility of increasing the effectiveness of the favourable conditions in the soil. Indeed, quite recently a substance has been produced on the large scale, which claims to owe part of its value to the possession of some such properties. As the matter is still under investigation, I do not propose to discuss it further this evening.

We go back, then, to our three established principles. Each of these can be recognised broadly in every case of crop production, but considerable difficulties arise when one tries to develop

FIG. 8.

Effect of phosphates on yield of oats.

Mitscherlich.

Equation for curve

$$\frac{dy}{dx} = (A - y) K.$$

any of them; there are so many factors involved and their interaction is so complex. I can best illustrate this by taking one of the factors in some detail, and I will choose one that has received very much attention from chemists, namely, the supply of phosphates.

Phosphates are indispensable for plant growth, and well conducted physiological experiments have shown a simple connexion between the amount of phosphate supplied and the amount of growth. Mitscherlich's results are plotted in Fig. 8; the plants were grown in sand, so as to eliminate various disturbing factors. But such simple results are never attained in soil. To begin with, there is always some phosphate already present. At first sight it

looks easy enough to take account of this, and simply add it on as a constant in the equation. It has proved almost impossible, however, to give any precise value to the amount of phosphate in the soil that is of any use to the plant. Ville showed years ago that the amounts revealed by chemical analysis are far beyond anything the plant can ever get:

Quantities of Potash and Phosphates in Soil and in Crops
(Ville at Vincennes, 1864).

	Lbs. per acre.		
	In soil.	In four crops of wheat.	In three crops of beetroot.
Phosphoric acid	1,581	62	132
Potash	2,025	102	288
Lime	34,674	—	—

In spite of the large amounts in the soil, the plants began to starve for want of added phosphates, and Ville rather gloomily concluded that "chemistry is powerless to throw light on the chemical properties of the soil." One could hardly expect chemists to acquiesce in that view, nor did they. They faced the situation quite squarely; they had found too much phosphorus and potassium in the soil: they proceeded to try again and find less. Instead of using strong acids they used dilute acids; several were suggested, and by a happy inspiration Bernard Dyer selected 1 per cent. citric acid as being the most suitable; although that was twenty-one years ago, 1 per cent. citric acid still holds the field in this country.

The part extracted by dilute acids was called the "available" portion to distinguish it from the "unavailable." The new method at once proved very helpful; it was used by Wood in his Norfolk experiments (T., 1896, 69, 287). Difficulties, however, soon began to arise. It was found impossible to assign any definite value to the amount of available phosphate present. Variations in the conditions of the experiment gave wholly different values for the amount of "available" phosphate, whilst in the case of nitric acid the longer the acid acted the less phosphoric oxide (P_2O_5) was extracted.

Now that gave the clue to the problem. It is obvious that there must be two actions going on: a direct solvent action and a reverse action, resulting in the withdrawal from the solution of the dissolved phosphoric oxide. The direct reaction was studied by extracting the soil by a diffusion method; this eliminates the reverse reaction, and the various dilute acids now behave very much alike. The direct solvent action, therefore, is much the same for all dilute acids.

The reverse action was studied by swamping the direct action. Sodium phosphate was added to the soil and the acid was then allowed to act. The whole of the added phosphate was never recovered, in spite of its solubility and the presence of the acid. Some of it was withdrawn from the solution by the soil. By varying the amounts of added phosphate a curve was obtained showing how the phosphate was distributed between the soil and the solution, which turned out to be the ordinary adsorption isotherm, similar in type to that obtained with charcoal and dilute

FIG. 9.

acids (Fig. 9). The constants are not the same for the different acids, and from these curves it is possible to go back and explain the apparently erratic action of the different acids on the soil.

Thus it appears that when phosphate is added to the soil for the purpose of increasing the growth of a crop it does not simply stop in the soil, waiting for the plant to take it up. It reacts with the soil; it is adsorbed, and the amount available for the plant at any time depends on the adsorption relationships. There is, in short, a competition between the plant and the soil for the

phosphate. Fig. 10 gives the curves for a clay and a sandy soil, which show that adsorption is greater for clay than for sand; in other words, the clay competes for the phosphate more vigorously than does the sand. An amount, therefore, which is sufficient for the plant growing in a sandy soil proves inadequate on a clay soil. This has thrown light on an interesting problem in manuring, for it has long been known that clay soils stood in more need of phosphatic manures than sands. The field results bring out this fact: the yield of barley on the heavy Rothamsted soil falls when

FIG. 10.

phosphates are omitted, but it does not react nearly so quickly on the Woburn sand.

It seems a far cry from the logarithmic curve expressing an adsorption isotherm to the management of barley and turnips, but the connexion is really simple and direct.

This, however, does not settle the matter. The plant is a living thing, and consequently its requirements are not rigidly constant, but vary with the conditions. There are very good grounds for supposing that the plant actually requires more phosphate on a

clay soil than on a sand. These are the effects produced by phosphates:

Promotion of early growth,
 „ „ root development,
 „ „ early ripening,
 Specially active on clay soils,
 „ „ in wet regions,
 „ „ for shallow rooting, and quick growing plants, for
 example, swedes and turnips.

It is possible that some simple connexion underlies all these, but no one has yet discovered it.

FIG. 11.

Yields obtained at Saxmundham on plots receiving (a) artificial fertilisers alone, (b) artificials and farmyard manure. The effect of withholding phosphates is seen to be far more marked in absence of farmyard manure than where it is added.

Saxmundham Results 1910-13.

Again, seeing that the need varies with the conditions, it is clear that if the conditions are altered the needs may change. When, for instance, a dressing of farmyard manure is applied, some of the properties of the soil are altered; it becomes more porous and more retentive of water, and phosphates may behave differently from what they did before. That is well shown on the Saxmundham plots (Fig. 11).

It is unnecessary to go any further. The point I want to bring out is that the simple and incontrovertible statement that phosphates increase plant growth proves very complex when applied in practice. So it is with the other factors. They can be disentangled and investigated, but it is not yet possible to put them together again and predict the resultant. One cannot set out from first principles and reconstruct the normal case of crop production; the factors are too numerous and too complex.

Yet something has got to be done. The technical chemist has the advantage over his colleague in the purely scientific laboratory that he cannot shelve an inconvenient problem. He is compelled to do something. The practical man brings in his problem with such insistence that, for shame's sake, one is bound to make an endeavour to solve it. A method has been evolved, an empirical method, which, whilst not very rigid, has at least the merit that it works. It consists in going into the field and finding out the actual agricultural properties of the soil by observations, inquiries, and direct field experiments; these have to be repeated for two or three years because the first results may only have been a trick of the weather, but if the same result is obtained for several seasons running, one may be sure of being right.

All that is old; it is, of course, Ville's method over again. The new part consists in trying to extend the results to other soils. For this purpose a soil survey of the area, usually the county, is arranged. The surveyor goes over the ground with a geological and orographical map and divides it up into areas of similar soil and vegetation characteristics. He then takes samples of soil typical of each area and makes fairly complete chemical, physical, and bacteriological studies of them. Simultaneously he collects information from the farmer about their agricultural properties, and, if necessary, arranges for proper field trials.

In this way a collection is made on the one hand of the agricultural properties, on the other of the chemical, bacteriological, and physical data, of typical soils. It is obvious that the possession of these standard soils helps the analyst and expert adviser very considerably; if a farmer asks for information it is much easier and safer to compare his soil with the standard than to attempt any absolute measurements. Moreover, these soil surveys are greatly facilitating advisory and analytical work.

They do far more than that, however. The normal case of crop production can never be decided on purely laboratory methods because there are always two or three varying factors, whereas in the ideal laboratory experiment there is only one factor varying. We are not, however, confined to the ordinary laboratory methods.

Statisticians have to deal with problems involving two or three variables, and they have worked out a method—the method of correlation—which, when intelligently applied, gives valuable results. It is hoped to apply this to crop production. The necessary masses of data are slowly being accumulated, and it is anticipated that very interesting results will be obtained.

The ordinary laboratory method, however—the one factor method—may still on occasions work satisfactorily. It sometimes happens in Nature that one of the various interacting factors overshadows all the rest and virtually eliminates them, so that here too it is possible to apply laboratory methods with satisfactory results.

For example, on a certain type of clay soil the whole situation is controlled by the circumstance that phosphates are almost absent, whilst the need of the plant for phosphates is particularly great. The addition of basic slag in these circumstances has caused most remarkable improvement. The best instances are seen at Cockle Park, and the results are given in their Bulletins.

Another illustration is furnished by our work on the partial sterilisation of soils. We had several times had occasion to carry out the fundamental experiment showing that the decompositions cease once the soil has been heated sufficiently to kill all organisms. On one occasion, however, the autoclave was out of order, and the soil was only heated sufficiently to kill the active forms and not the spores. To our surprise, the bacterial activity was now much greater than before. Other methods of partial sterilisation had the same effect. These results are reflected in plant growth, which is markedly increased by partial sterilisation. Other experiments, which I need not now describe, showed that any treatment of the soil which is detrimental to life produces the same effect, sometimes, of course, with secondary effects as well. Conversely, it was shown that any method of treatment favourable to life, such as increased temperature and moisture content and addition of stable manure, leads, sooner or later, to a falling off of bacterial activity. The simplest explanation of the phenomena is that the soil population can roughly be divided into two groups: one favourable to the production of plant food, the other not. The useful population is, on the whole, more resistant to adverse circumstances than the harmful organisms, and therefore survives more drastic treatment. Hence any method that kills some, but not all, of the soil population effects an improvement and leads to good results. A continued spell of favourable conditions, however, enables the harmful organisms to establish some sort of superiority. This hypothesis throws important light on the behaviour of the

soil in natural conditions, and it reveals another factor in crop production.

We have not yet succeeded in making much of it in the normal case; indeed, we have hardly attempted to do so, for the reason I have already given: there are so many interacting factors. There are, however, cases where this one factor largely dominates the situation. In glasshouses run at a high pitch, where the soil temperature and water content are high, and where large dressings of organic manures are used, the bacterial efficiency falls off so much that the plants begin to suffer. The soil, in the picturesque language of the practical man, is said to become "sick." This sickness proved so difficult to deal with in practice that the problem was given up; the soil was thrown out and new soil brought in to take its place.

As soon as the problem came to our notice it was easy to suggest a remedy. The reduction of bacterial activity seemed clearly due to an excessive development of the detrimental organisms. It was only necessary to adopt partial sterilisation to get rid of these and to give the useful organisms a better chance of action. The basis of a suitable method was already in existence; steam had been used to kill insect pest in the houses, and by suitable modification this process was successfully used for the treatment of sick soils.

In cases of this sort, where only one factor is concerned, one is back to the old laboratory conditions in which one was brought up and trained to think, and consequently there is always considerable hope of success. It is fortunate for us that these cases puzzle the practical man more than the normal case, so that he rather tends to bring them in as specially difficult problems, and any little success on our part bring us more credit than we really deserve.

There is, however, a very real advantage. The most fruitful ideas for working out the development of our subject have often been got from abnormal cases brought in by the growers. Practical men have the great advantage that they are compelled to keep their eyes open for Nature's problems; they cannot shirk them, or they find their crops suffering and themselves losing money. The close association of Science with an industry is, therefore, a great advantage, because it brings in new problems which, if properly investigated, may prove extremely valuable in opening up new fields of knowledge. There is an exhilarating freshness about all this work that one often misses in the more academic investigations.

All the same, while speaking in praise of applied science, one

1858

BRADY AND DUNN :

must recognise that science cannot be applied until it is developed. We have seen, and instances might have been multiplied, how the hydrolysis of protein throws light on the proper management of a manure heap, and how the adsorption isotherm worked out for charcoal and dilute acids clears up a difficulty in the manuring of turnips. It is impossible to set any limit to the value of good work in science honestly carried out. The fact is that Science and Creative Industry are one and indivisible, and any attempt to divorce them may only end in disaster.