THE VISIBILITY OF RADIATION IN THE BLUE END OF THE SPECTRUM.

By L. W. Hartman.

THE determination of the visibility of radiation in the short wave-length region of the spectrum is difficult, owing principally to the small amount of light available when using ordinary methods and sources. The optical pyrometer method, which had previously been employed¹ in obtaining this function for the red end of the spectrum, minimizes this difficulty, and has now been used to obtain values of the visibility in the blue end out as far as wave-length 0.400μ . In contrast to results obtained by the flicker photometer, this method involves the direct-comparison principle.

The effect of stray light and of extreme color-difference was eliminated to a very large extent by the use of a piece of blue glass, of known spectral transmission, in the eyepiece of the telescope. During any one run the brightness was kept constant, but experiments were also made with values of brightnesses as seen through the blue glass ranging from 0.000033 to 0.0013 candle per square centimetre. A small field was employed, and tests indicated no evidence of a Purkinjé effect. Corrections were made for slit-widths, for the dispersion and selective absorption of the prism and lens system, for scattered light, and for the absorption of the blue glass.

Measurements were made by twenty observers, all of whom were trained in laboratory work. The results, together with the published data of other investigations are given in the table.

Wave-lengths, $\mu\mu$	Mean visibility of twenty subjects	Mean values given by Nutting*	Mean values given by Coblentz and Emerson
410	1.7	9.5	24
420	11.4	17.1	42
430	32.6	30.3	59
440	61.6	58.0	71
450	100	100	100
460	153	168	137
470	240	266	202
480	376	392	305
490	620	566	474
500	905	828	770

*These values, kindly furnished by Doctor Nutting, differ slightly from his published data, owing to the redetermination of the distribution of energy in the spectrum of the acetylene flame used.

¹ *Astrophys. Jour.*, vol. 42, 1915, p. 285.