FIG. 4. Showing the arm before and after operation.

arm rotate inward and, as stated above, this rotation takes place at the point of fracture. This position may be corrected by passing a broad piece of adhesive around the hand, extending about four or five inches beyond, where another piece of adhesive, being reversed, is made to overlap the first piece and then passed along the

back, thus holding the hand and arm, and preventing rotation. (Fig. 3.)

I wish to report a case of cubitus varus in a girl ten years of age. She fell from the cradle when a baby, striking the left elbow, and ever since has had a gun-stock deformity. For some time previous to her consulting me, the fingers had been tender and numb, and she would often drop things from her hand. There had also been pain and tenderness in the elbow, so that it had been considered suspicious of tuberculosis.

An osteotomy just above the external condyle was done April 14, 1913, and with the restoration of the proper carrying angle, the symptoms in elbow and hand completely disappeared. Fig. 4 shows the arm before and after the operation.

Therapeutic and Preventive Medicine.

OBSERVATIONS ON THE DIAGNOSIS AND TREATMENT OF DIABETES.*

By MARTIN M. BROWN, M.D., F.A.C.S.,
NORTH ADAMS, MASS.

CONCERNING the diagnosis and treatment of this malady, I have nothing original to offer, but will present a few facts gleaned from various sources. All authorities are agreed that the condition of diabetes mellitus is increasing; that it is a disease largely, but not exclusively, of the well-to-do. The malady occurs at all ages and in both sexes. Heredity plays a large part, and is traceable in at least 20%. Observers are agreed that functional or organic disturbances of the pancreas are the main causative factors. Experimental observation upon dogs shows that complete extirpation of this organ is invariably followed by diabetes. The conclusion is drawn that the pancreas supplies an internal secretion essential to carbohydrate metabolism. Evidence has been accumulated which shows that other organs besides the pancreas exert an influence upon this process. Disturbances of the thyroid gland, parathyroid and suprarenal capsules are many times associated with glycosuria.

Normal sugar is present in the blood from 0.05% to 0.12%. In diabetes the amount varies from 0.16 to 1.16%. The kidneys are able to hold back sugar when it is present in normal quantity, but when the sugar concentration is increased, the kidneys are no longer able to hold the sugar back. Why does sugar accumulate in the blood? Two theories suggest themselves; one, that the diabetic cannot oxidize sugar; the other, that too much sugar is made. Normally, sugar is derived from carbohydrates, which make a very large amount of our food. That sugar may be formed from proteins has been demonstrated by many experiments.

Transient glycosuria, besides being caused by diabetic tendency, may be due to injuries, to feb-

*Read before the Berkshire District Medical Society, November 4, 1915, as part of a Symposium on Diabetes.

rile infectious diseases, to certain poisons, and to the excessive intake of sugar. In my own personal observation, I have met with several cases of transient glycosuria, due, undoubtedly, to the excessive use of sugar. In these instances, a proper regulation of diet in a short time rendered the patient sugar-free.

The symptoms associated with diabetes are familiar to you all. The diagnosis is established when repeated examinations of urine show the presence of sugar. Generally speaking, we become suspicious that a person may be suffering from glycosuria when he complains of excessive and frequent urination, unnatural thirst, pruritis, vague affections of the nervous system, and digestive disturbances which are not otherwise explained. The presence of boils and furuncles are always suggestive of diabetic conditions. It is a well established fact that all patients suffering from this disease have a marked lessened resistance, and the presence of sugar in the blood renders them especially prone to all kinds of infections.

The ideal treatment for any disease is the removal of the cause. In diabetes this is not possible, for the cause is unknown. The condition should not be treated in a routine manner; cases differ and must be treated individually. In no disease is the patient to be warned more against quackery and quack remedies, for the harm of improper treatment cannot be well retrieved. Diabetes is a disease against which the plan of battle cannot be laid for a month or a year, but for a decade, or possibly a lifetime.

Having determined that the patient has diabetes, after several urinary examinations, the next question to be considered is whether the case is mild, moderately mild, or severe. In mild cases, glycosuria disappears from the urine as soon as carbohydrates are eliminated from the diet. It is moderately severe when the sugar disappears upon restriction of the diet, but reappears when carbohydrates in minimum quantities are added. It is severe when sugar cannot be removed from the urine by eliminating all carbohydrates from the diet and restricting the intake of protein food. The essential point of the dietetic treatment consists in bearing in mind the impaired tolerance of the diabetic for carbohydrates by not overtaxing this tolerance. The power of assimilation for carbohydrates is not only conserved in the majority of cases, but increased.

Defective non-combustion or oxidation of carbohydrates, as well as of proteids, in diabetes brings about a condition known as acidosis, due to the presence of beta-oxybutyric acid, which accumulates in the blood, and together with its derivatives, acetone and diacetic acid, is eliminated in the urine. Acetone is also eliminated in the breath. When this condition is present, the patient is liable to develop diabetic coma. This coma is usually preceded by premonitory symptoms, such as restlessness, irritability, dizziness, pain in the limbs, headache and drowsiness.

The treatment of acidosis will depend largely on the previous history of the case. If the patient has been under competent observation, and has developed ketonuria on a strict diet, largely of proteids, he needs carbohydrates. Sodium carbonate should be given in large doses, frequently repeated until careful urinary tests show freedom from this condition. The treatment of diabetic coma must be prompt, energetic and intelligent. Free bleeding is oftentimes advisable, free elimination from the bowels with bitter salines, and the introduction into the blood, by every way possible, of alkalis.

The tendency of all diabetics to become infected with various maladies calls frequently for surgical intervention. No diabetic is considered a good surgical risk. In every case requiring surgical intervention, preliminary treatment, if possible, should be instigated. The urine should be made sugar-free, and the condition of acidosis eliminated; strictest aseptic precautions should be used. It is well to administer alkalis before an operation, whether diacetic acid is or is not present in the urine. The fear of operation upon diabetics has, in a great measure, subsided, and no up-to-date surgeon hesitates to resort to surgical methods of treatment in any diabetic case, when indicated.

My experience in the use of drugs or special remedies, with the exception of one preparation, is unsatisfactory. The continued and intelligent use of trypsin in certain selected cases of a pancreatic type seems to have a beneficial influence. The small quantity of gold and arsenic act as a good tonic for the anemic condition, and the pancreatic ferments, theoretically at least, supply the needed defects, and materially aid in the digestion of carbohydrates.

CASE REPORTS.

CASE 1. Mrs. J. C., married, age 62 years. Under observation past 10 years. Passing large quantities of sugar in urine when first came under my observation. Diet restricted. Trypsin administered. Gradual improvement. At no time has sugar been absent from urine, but greatly lessened. Has not taken tablets for about 5 years, but keeps in fairly good health by restricted diet.

CASE 2. Mrs. S. W., married, age 60 years. Under observation past five years. Passing continuously large quantities of sugar in urine. Restricted diet; use of trypsin. Still using tablets, and finds that condition becomes much worse if omitted.

CASE 3. Mr. C., widower, about 50 years of age. Under observation 3 years. Sugar continuously present in large quantities. Unable to restrict diet to any degree. Takes trypsin tablets faithfully. Urine sugar-free at end of two years. Discontinued tablets, sugar returned. Sugar-free when resumed tablets. General health first class, but gradual loss in weight.

CASE 4. Master L. S., 10 years of age. Under observation one year. Acute manifestation, no apparent cause. Attempted to restrict diet and

placed on trypsin tablets. No improvement. Referred to Dr. Joslin of Boston. Placed in Deaconess Hospital and Allen treatment administered. Became sugar-free in about one week. Allowed to return home in six weeks. Remained sugar-free when instructions were carefully followed. Sugar present at repeated examinations during past three months, at times. Sugar present in small quantities at present time in spite of starvation treatment one day a week, and restricted 5% diet. Outlook unsatisfactory.

CASE 5. Mrs. R., age 58 years. Sugar present continuously in large quantities. Treatment—restricted diet, use of trypsin tablets. Material improvement, but at no time sugar-free. General health at present time much improved over that of two years ago.

REFERENCES.

- Riesman, D.: *Therapeutics of Internal Diseases*. Forchheimer, Vol. 11, p. 717.
 Allen, F. M.: *The Treatment of Diabetes*. BOSTON MED. AND SURG. JOURN., June, 1915.
 Dicalafoy Text-Book of Medicine, Vol. 11, p. 1802.

THIRTEENTH RUSH SOCIETY LECTURE.*

(ABSTRACT)

Investigative and Scientific Phases of the Diabetic Question With Their Probable Relations to Practical Problems of Clinical Medicine, by Dr. Frederick M. Allen of the Rockefeller Institute for Medical Research.

Dr. Allen emphasized the necessity of keeping in mind the distinction between glycosuria and diabetes. The one was a symptom; the other a disease. One was glucose in the urine; the other a defect in metabolism. Defining diabetes, the author recognized the increasing agreement on the part of investigators that diabetes was deficiency of the internal secretion of the pancreas. This doctrine, it was said, went back to early observations of gross changes in a few cases of clinical diabetes, finally followed by Von Mering and Minkowski's production of acute diabetes by total pancreatectomy. Sandmeyer, by means of atrophy of a pancreatic remnant, produced a type of chronic diabetes comparable to the unusual human cases due to atrophy of the pancreas. A more accurate reproduction of human diabetes was afforded by removal of most of the pancreas, leaving the remnant with its normal blood supply and duct communication with the bowel. De Renzi and Reale first reported such an experiment in 1892. This form of diabetes was later rediscovered independently in four different places about the same time. The first in point of publication were Thirloix and Jacob, who observed that when a sufficient portion of a dog's pancreas was removed, leaving the remainder in its normal connection with duct and blood vessels, this remnant retained its normal appearance and supplied enough pancreatic juice to maintain the animal's nutrition, yet diabetes developed and ran a gradual downward course much like the human disease. Their second important observation was that in suitably prepared animals this downward progress could be produced or hastened by over-feeding with carbohydrate. The second in point of publication was Helly, who studied only the glycogen, finding that in this form of diabetes the animals retain more or less glycogen, here resembling human diabetes. The third was the work of Allen, which dealt with various points concerning this type of diabetes, compared it with the normal and with other forms of diabetes and glycosuria, and also, thanks to the coöperation of Professor F. B. Mallory, described, for the first time, the specific

changes in the islands of Langerhans in experimental diabetes. The fourth in point of publication was Homans, who discovered this form of diabetes in cats, and made the most accurate microscopic studies.

By this method, human diabetes, Dr. Allen stated, could be imitated in animals in the following particulars; (1) Mode of onset, sudden or gradual; (2) the influence of diet; (3) some individual idiosyncracies; (4) the diminished permeability of the kidney for sugar; (5) the effect of nervous influences; (6) the effect of traumatism; (7) a temporary stage during which there was glycosuria with little or no polyuria, thus resembling diabetes decipiens; (8) complications in the cachectic stages, generally not identical with the human complications, but rather an increased tendency to corresponding canine afflictions. There was no evidence that complications were due to increased blood-sugar.

The chief points of alleged difference between clinical and experimental diabetes, it was pointed out, had been the following: (1) The dextrose-nitrogen ratio, which is Lusk's figure of 3.65 in the severest human diabetes, whereas removal of the dog's entire pancreas gave only the Minkowski ratio of 2.8. Total pancreatectomy in the human subject, he said, should theoretically give a 3.65 ratio. It was well known, he observed, that phloridzin gave a 3.65 ratio in some species, and a 2.8 ratio in others; likewise the difference of species might explain the different ratios in diabetes. (2) The marked increase of metabolism in totally depancreatized dogs. To the author it seemed probable that partly depancreatized dogs with severe diabetes would not show such a great increase; it was now known that some severe human diabetics showed an increase of metabolism. (3) Anatomical changes in the diabetic pancreas. Weichselbaum had described so-called hydropic degeneration and other lesions in a series of 186 diabetic autopsies. Other investigators had not been convinced of the existence of any specific alterations in the diabetic pancreas. It was obvious that such advanced and generalized changes as were found in the pancreas in experimental diabetes were seldom, if ever, present in human cases, because they were too obvious to be missed. The classical work in the microscopic study of experimental diabetes was that of Homans. Using the Lane-Bensley technic, Homans proved that only the Beta cells showed the exhaustion and degeneration, while the Alpha cells remained intact. Homans had found some similar changes in at least a portion of the islands in human diabetes, and it would seem that the most promising prospect for the microscopic investigation lay in this direction. The functional element in human diabetes was also to be considered which, it was said, was emphasized by the results of treatment. The rapid clinical changes possible in human diabetes seemed to the author to indicate functional rather than organic alterations in the pancreas. Back of human diabetes might be, perhaps, a disorder of the abdominal nerves, possibly disturbing not only the function of the pancreas, but to some extent that of other organs. Though the kidneys were generally normal microscopically in diabetes, functional changes were frequent. Further study of the pancreas obtained quickly after death and fixed and stained by the best methods was regarded as important for deciding this question, and physicians who could obtain such material would render a service by sending for study. (4) A fourth difference was that found in ketonuria or acidosis. It was an old statement that the diabetic patient died in coma and the diabetic animal in cachexia; but by stopping the over-feeding of diabetic patients coma might be less frequent. Also, under suitable conditions, diabetic animals seemed to be subject to this intoxication. The present status of the theories of ketonuria or acidosis were discussed. The fact that severe diabetics were cleared up in these respects by fasting, whereas normal persons or milder diabetics developed ketonuria

* Delivered in Thomson Hall of the College of Physicians of Philadelphia, Jan. 21, 1916, at 8.30 P.M.

on fasting, along with various other experimental and clinical observations, threw the existing theories into considerable confusion. Ketonuria and acidosis seemed to the author to be not explainable by mere lack of carbohydrate, or even by habituation or individual idiosyncrasy. The fundamental nature of the process was in doubt. Diabetes, Dr. Allen stated, might probably be defined as a specific deficiency of the power of assimilating food. The accumulation of reserves, as when the body gained weight, diminished tolerance and aggravated diabetes, whereas the reducing of the reserve stores by low diet or by exercise increased tolerance and checked diabetes. Also one kind of food lowered the tolerance for another kind of food, as adding carbohydrate to a diabetic diet might give rise to ketonuria, and adding fat to a diabetic diet might give rise to glycosuria. The history of diabetic therapy consisted essentially in an interweaving or alternation of two principles: (1) The restriction of the sugar-yielding elements of the diet, namely carbohydrate and protein; (2) diminution of the total caloric value of the diet. The fasting treatment was in harmony with the observed benefits of these methods in the past. This had been previously outlined and details were to be published in monograph form. The principle underlying it was a well-established one in all therapeutics,—rest of a weakened function. No positive means of strengthening this function aside from rest has yet been discovered.

Medical Progress.

PROGRESS IN GYNECOLOGY.

By STEPHEN RUSHMORE, M.D., BOSTON.

(Concluded from page 282.)

OPERATION FOR PROLAPSE OF THE UTERUS.

Mayo²³ discusses several forms of operation for prolapse of the uterus with associated pelvic relaxation. He advocates interposition of the uterus for cystocele with the first or second degrees of descensus of the uterus, but rejects it for the third and fourth degrees. For certain conditions after fifty, in which there is considerable protrusion after hysterectomy, a complete obliteration of the vagina has proved adequate and satisfactory.

If, with considerable prolapse, the bladder is not extensively torn from its uterine supports, with resulting cystocele, some modification of the Kocher operation is to be recommended. In these cases the chief object to be attained is to fix the uterus permanently. Sutures as ordinarily placed, tend to cut out in time and the uterus tends to come down. This has occurred in a sufficient number of cases to need special means for its prevention.

The operation of Mayo is based on Murphy's modification of Kocher's abdominal hysteropexy—splitting the uterus in the median line from the fundus to the internal os, and excising the endometrium. Each lateral flap of uterine wall is then turned outward from the median line and sutured between the rectus and its anterior sheath by chromic catgut and silk sutures. It is

the placing of the uterine tissue beneath the fascia, which, according to Mayo, gives a better closure than the Murphy modification. The closure is in the usual manner.

In the cases where there is considerable cystocele, the uterine supports being also much relaxed and the uterus itself small, better opportunity for plastic support of the pelvic structure is given if the uterus is removed. The recommended incision appears as an inverted V on the anterior vaginal wall, the apex near the external urethral meatus, and the arms of the V being extended laterally and posteriorly to encircle the cervix and meet in the median line posteriorly. After opening the peritoneal cavity and delivering the fundus anteriorly, the broad ligaments are clamped with heavy straight clamps from above downward, two on each side. The incisions laterally to remove the uterus vary in their position according to the relaxation of the tissues. If not much relaxation is present, they may be made so as to leave in a slice of the uterine wall on each side. If there is considerable relaxation, part of each broad ligament may be removed with the uterus. Suturing, with continuous mattress of chromic catgut, begins posteriorly at the tips of the second clamps, which are rolled out so as to approximate the peritoneal surfaces and give no raw area in the peritoneal cavity. The suture passes back and forth behind the forceps, completely through the ligaments, tightening them. When the round ligament is reached, it is caught into the angle of dissection, where the bladder wall has been separated from the anterior wall of the vagina. "The suturing extends backwards on each side from this point, catching into the broad ligaments and then on each side into the angle of the depth of the dissection, thus compelling the bladder to rest on the broad ligaments." This is the essential step in the cure of cystocele. The rest of the closure is by running suture, the mucous membrane being approximated by submucous catgut. Perineal restoration completes the operation.

INVERSION OF THE UTERUS.

Roberts²⁴ presents an interesting and somewhat comprehensive account of inversion of the uterus, including historical notes, and a description of two cases seen by himself. Of these cases, one was relieved by taxis, a procedure recommended by Hippocrates; and the second cured by operation, the treatment recommended by Themison and Soranus. He gives references to the exhaustive studies of Crosse, Thorn and others, and has contributed a very readable paper.

Buck²⁵ reports two cases of inversion of the uterus on which he operated and notes briefly the technic employed. The anterior wall of the uterus was split from vagina to fundus. This free incision does not materially increase the loss of blood and facilitates eversion. He did