A discussion on "The Supposed Risks Attending X-Ray Treatment of Ringworm" was held by the Dermatological Section of the Brit. Med. Association, July 30, 1909, and reported in the *British Journal Dermatology*, Dec., 1909, p. 391. Dermatitis and permanent alopecia have resulted in treatment of a few cases, but the most important question was the effect of the X-rays upon the brain. In conclusion it was the general consensus of opinion that the X-ray had no effect upon the brain when properly used in the treatment of ringworm.

J. M. K.

TREATMENT OF VERRUCAE PLANTARES.

R. L. Sutton (*Jour. of Cut. Dis.*, Apr., 1909).

The author used carbon dioxide according to Pusey's method for freezing the warts. He applied a pencil of the snow for thirty to sixty seconds at first, and after thawing re-applied for a second time of thirty seconds. Boric acid was dusted over the surface. No anesthetic was required. The method was used in four cases with excellent results.

J. M. K.

EXAMINATION OF THE URINE IN SKIN CASES. (*Derm. Centralb.*, Apr., 1909, p. 194.)

The author concludes that there is a close connection between outbreaks of the skin and internal disorders after examining the urine in 200 skin cases.

In pruritus he finds deficient acidity, and phosphates, and has obtained good results in the treatment by using phosphate preparations.

In chronic eczema the urine is concentrated

with an increase in acidity and nitrogenous excretion; there is also a slight diminution in chlorides.

In psoriasis the condition is practically the same as that found in chronic eczema, except more pronounced.

In acne and rosacea there were variations in concentration, and indican was found in 50 per cent of the cases.

In urticaria indican was found as a rule and in greater amount than in acne; otherwise the results resembled those found in pruritus.

In prurigo of Hebra, as it is so often associated with urticaria, he looked for increased indican, but did not find it.

Albumen was found in 2 per cent of the cases.

J. M. K.

TREATMENT OF ERYSIPELAS.

By Barclay (*Amer. Jour. Derm.*, Jan., 1909).

The disease can be controlled and in many instances aborted. The quick administration of calomel and large and repeated doses of quinine—half a drachm daily—are advocated. Tincture of iron is also a valuable remedy. But the most important treatment for the control and cure of erysipelas lies in local measures. The following formula has never failed, in the writer's hands, to control the spread of the disease, cut short its duration, and prevent infiltration and ulceration of the skin: Sodiri boratis, Ziv.; phenolis, Zss.; glycerine, Ziv., equal grs. Zvi. Apply hourly. Cloths saturated with lotion are constantly applied over the affected area. The writer has not observed ulceration for thirty years.

J. M. K.