labels should be privileged to conduct their interstate traffic in their so-called medicines, admittedly worthless, because Congress did not intend to reach them.

Nor does it seem to me that any serious question arises in this case as to the power of Congress. I take it to be conceded that misbranding may cover statements as to strength, quality and purity. But so long as the statement is not as to matter of opinion, but consists of a false representation of fact—in labeling the article as a cure when it is nothing of the sort from any point of view, but wholly worthless—there would appear to be no basis for a constitutional distinction. It is none the less descriptive—and falsely descriptive—of the article. Why should not worthless stuff, purveyed under false labels as cures, be made contraband of interstate commerce,—as well as lottery tickets? *Champion v. Ames*, 188 U. S. 331.

I entirely agree that in any case brought under the act for misbranding,—by a false or misleading statement as to curative properties of an article,—it would be the duty of the court to direct an acquittal when it appeared that the statement concerned a matter of opinion. Conviction would stand only where it had been shown that, apart from any question of opinion, the so-called remedy was absolutely worthless and hence the label demonstrably false; but in such case it seems to me to be fully authorized by the statute.

Accordingly, I reach the conclusion that the court below erred in the construction that it gave the statute, and hence in quashing the indictment, and that the judgment should be reversed.

I am authorized to say that Mr. Justice HARLAN and Mr. Justice DAY concur in this dissent.

Correspondence

Knots

To the Editor:—In THE JOURNAL, April 8, is an article entitled, "An Efficient and Time-Saving Method of Tying the Square Knot." In the *South Carolina Medical Journal*, October, 1909, appeared an article in which I described a method of tying knots identical with the one described by Drs. Castle and Ryfkogel. The maneuvers are reversed, however, that is to say, the first turn of their knot is brought down by my second maneuver, etc. Their description also refers to carrying out these maneuvers with both hands simultaneously in order to tie the friction knot. Strange to say, I also anticipated them in describing this one.

B. B. STEEDLY, Spartanburg, S. C.

To the Editor:—Being interested in knots, I attempted to tie this one according to the directions and discovered an error. If the first half of the knot is tied as directed and as shown in Figures 1, 2, 3, 4, I do not get the result as shown in Figure 5. Instead, hands A and B must be crossed to make this half of the knot lie down flat so that it will hold while the second half is tied. This is an awkward position for pulling up the first half which must be snug. This can be obviated by starting with ligature C in hand B and ligature D in hand A. This change will cause the first half of the knot to lie down flat while the second half when tied as given in the directions will come into place properly when the hands are crossed. This is not such an inconvenience in the case of the second half as it does not need to be pulled so snugly.

W. H. BAILEY, Kansas City, Mo.

To the Editor:—In March, 1909, I presented in *Surgery, Gynecology and Obstetrics* an original method of tying surgical knots rapidly and accurately. I have used the method continuously since 1901. The "one-finger" principle described by Drs. Castle and Ryfkogel is identical with mine. The method of application differs merely in the fact that they make first the knot I described as the second, while their second half is my first knot. (Practically, it depends entirely on the manner the hand happens to catch the ligature, whether the one knot is made first or last.) For these reasons I respectfully claim priority in devising and publishing this

rapid-tying method. I am very much in accord with Drs. Castle and Ryfkogel in their arguments. This method of making knots will do all they assert and more.

Slight errors are unquestionably due to the artist and not to the surgeon. The context makes this clear. For instance, Figure 4 does not represent any part of a knot. In Figure 5 the hands are shown as having interchanged ends. If this were not done, the hands must be held crossed to show this step in making the knot.

E. P. QUAIN, Bismarck, N. D.

[The above extracts from letters were referred to Drs. Castle and Ryfkogel. The former replies.]

To the Editor:—The principles of producing the square knot which Drs. Quain and Steedly published are identical with those we recently gave. Our application of these, which by the way is merely composed of the movements which have long been employed to form the well known "crochet knot," is less cumbersome than Dr. Quain's. Further, we have given the information necessary to obtain a knot which has two turns in each half, which some surgeons deem very important.

The article that appeared in THE JOURNAL, April 8, 1911, was written over three years ago, since which time it has been lying on the shelf. It was devised, written, and published without the slightest knowledge of any other person's work. I compliment these other authors on their ingenuity and ask your pardon if this occurrence has caused you any embarrassment.

As to Dr. Bailey's criticism: The principle of tying the knot is vividly shown. It would be confusing to mention the end of the ligature that should be held in the grasp of either hand. That is too obvious. If the person tying is the surgeon and if he retains his needleholder in his right hand, thereby using his left hand to tie the free end, one can readily understand that the conditions will be different from those wherein the assistant ties the free end of a continuous suture.

If any one desires to learn to tie this knot by these movements he will have no trouble in learning the method from the article as it stands. He will soon learn to adjust his ligatures in accordance with the conditions governing the situation, i. e. if he is tying with his right or left hand or with both hands, whether he is tying the free end of a continuous suture from the assistant's side of the table or from the surgeon's, or if he is tying ligatures.

H. EDWARD CASTLE, San Francisco.

Queries and Minor Notes

ANONYMOUS COMMUNICATIONS will not be noticed. Every letter must contain the writer's name and address, but these will be omitted, on request.

TREATMENT OF COAL-OIL POISONING

To the Editor:—Please give the treatment of poisoning from coal-oil. I do not remember having noticed a description of this particular form of poisoning in any of the text-books.

F. S. SPEARMAN, M.D., Whiting, Iowa.

ANSWER.—The treatment of coal-oil poisoning consists in washing out the stomach or administering emetics and using purgatives. In cases of collapse, warm baths with cold effusions are recommended; the usual respiratory and cardiac stimulants are also indicated.

DIGESTIBILITY OF SOUR AND SWEET MILK

To the Editor:—Kindly give me information concerning the relative digestibility of sour and of sweet milk, both for adults and for infants.

C. B. PURVIS, M.D., Brookline, Mass.

ANSWER.—Sour milk may be consumed in two different forms. The one is the naturally soured milk, the other buttermilk. The difference between the two is, first, that sour milk contains the full amount of butter-fat, and, second, that the coagulated casein is not churned in sour milk. The churning of the casein undoubtedly adds to the digestibility, since the more solid curd of sour milk has to be broken up in the stomach. Both sour milk and buttermilk may be considered to be partially predigested as compared to sweet milk and both are therefore more easily digestible, buttermilk to a higher degree than sour milk.