THE EFFECT OF BENZENE ON THE PRODUCTION OF ANTIBODIES *

LUDVIG HEKTOEN

From the Memorial Institute for Infectious Diseases, Chicago

Selling¹ found that in rabbits benzene in suitable doses causes profound destruction in the hemopoietic structures, especially of the marrow, followed when recovery takes place by prompt regeneration, which is complete in from 10 to 20 days. During the height of this effect there develops a marked leukopenia, involving the granular cells relatively more than the lymphocytes. Fatty changes may develop in the kidneys and the liver, and hemorrhages in the lungs and pleura, the stomach, etc. Some reduction of red corpuscles results, embryonal red cells may appear in the blood, and the blood platelets may be reduced also,² marrow giant cells being destroyed.³ In view of this more or less selective action of benzene on tissues and cells that are concerned in the production of antibodies and in the defense against infection, it lies near at hand to investigate the course of infection and the production of antibodies in animals under the influence of benzene.

Winternitz and Hirschfelder⁴ found that rabbits in which leukopenia had been produced by means of benzene were little resistant to pneumonia from intratracheal insufflation of pneumococci, there being only very few leukocytes in the exudate. Kline and Winternitz⁵ published further studies on the importance of the leukocytes in the reaction against pneumonia as observable in benzenized rabbits, but they did not study directly the influence of benzene on the production of pneumococcal antibodies.

White and Gammon⁶ inoculated rabbits with tubercle bacilli and then gave benzene by way of inhalation. They found that rabbits so treated were less resistant to tuberculosis than rabbits which had not received benzene, but they offered no basis for any explanation of this effect.

In a study of the inflammatory reactions in the ears of rabbits with marked benzene leukopenia, Camp and Baumgartner⁷ noted that there apparently was no restraint on the entrance and growth of bacteria in such animals.

* Received for publication January 7, 1916.

¹ Ziegler's Beiträge, 1911, 51, p. 567.

² Duke: Arch. Int. Med., 1913, 11, p. 100. Hurwitz and Drinker: Jour. Exper. Med., 1915, 21, p. 401.

³ Duke: Arch. Int. Med., 1913, 11, p. 100. E. Frank: Berl. klin. Wehnschr., 1915, 52, p. 961.

⁴ Jour. Exper. Med., 1913, 17, p. 657.

⁵ Ibid., 1914, 18, p. 50; 1915, 21, p. 320.

⁶ Tr. Assn. Am. Phys., 1914, 19, p. 332.

⁷ Jour. Exper. Med., 1915, 22, p. 174.

Schiff⁸ found that in guinea-pigs small doses of benzene—0.01 c.c.—intra-peritoneally injected before the injection of antigen increased the reactivity to the second dose of antigen, whereas larger doses of benzene—0.03 c.c.—decreased the reactivity. Schiff mentioned several explanations but favored the view that the effect most likely is due to the action of benzene on the tissues which produce antibodies.

Rusk⁹ was the first to publish results of direct observations on the effect of benzene on antibody-production. He recorded that in rabbits the injection of benzene in doses of 1 c.c. for each kilo of weight, either before or at the same time as the antigen, reduced greatly the formation of lysin for sheep corpuscles and of precipitin for horse serum.

Simonds and Jones¹⁰ found that in rabbits injected with benzene in doses of about 1 c.c. per kilo of body weight the production of lysin for dog corpuscles and of agglutinin and opsonin for typhoid bacilli was reduced, the reduction being most marked in the case of lysin and least marked in the case of opsonin. In their work, with occasional exceptions, benzene reduced the number of leukocytes in the blood, especially the polymorphonuclears, and sooner or later evidence of damage to the erythroblastic centers appeared in the form of stippled and nucleated red cells, but in both these effects there were observed sharp individual variations.

In connection with other experiments I have been making observations for some time on the action of benzene and certain related substances on the production of antibodies.¹¹ It was thought that substances acting so directly on blood-making structures would affect the output of antibodies, in the elaboration of which these structures appear to take the leading part.

For the study of this action, benzene in an equal quantity of olive oil, was injected subcutaneously into rabbits, which had already received, or subsequently were to receive, 30 c.c. of sheep blood intra-peritoneally. Quantitative determinations were then made of the newly formed lysin and precipitin. Sheep blood in the quantity mentioned was selected for antigen because I knew from other work that usually it calls forth in otherwise normal rabbits the production of considerable, tho somewhat variable, quantities of lysin for sheep corpuscles and of precipitin for the proteins in sheep blood. Undoubtedly the injection of a considerably smaller quantity of sheep blood would have served the purpose. In order to guard against mistakes of interpretation on account of inherent variation in response to the antigen, each experiment included several animals as nearly as possible of the same age and weight, and the estimations of the antibodies studied extended over the entire period covered by the antibody curve.

⁸ *Ztschr. f. Immunitätsf.*, 1914, 23, p. 61.

⁹ *Univ. Calif. Publ. in Pathol.*, 1914, 2, p. 139.

¹⁰ *Jour. Med. Research*, 1915, 33, p. 197.

¹¹ A brief report is published in *Tr. Chicago Path. Soc.*, 1915, 9, p. 308.

The preparation of benzene used in all the experiments was "Benzene Merck. Highest chemical purity—crystallizable."

The details of the following experiments will serve to illustrate the methods of procedure and the principal results.

In the tables and curves the figures referring to "Lysin" represent the highest dilution of the rabbit serum in question in which distinct lysis of sheep corpuscles was produced. The tubes were incubated for 2 hours and then placed in the ice-box until the next morning. In each test the total quantity of the mixtures was 0.6 c.c., of which 0.2 c.c. was 5% suspension of sheep blood; 0.006 c.c., guinea-pig serum as complement; and the rest, rabbit serum and salt

TABLE 1

THE EFFECT ON LYSIN AND PRECIPITIN IN RABBITS, OF 8 CONSECUTIVE DAILY INJECTIONS OF 1 C.C. BENZENE, BEGINNING ON THE 4TH DAY AFTER INJECTION OF SHEEP BLOOD

Days After Injection of Sheep Blood	Rabbit 1		Rabbit 2		Rabbit 3		Control	
	Lysin	Precipitin	Lysin	Precipitin	Lysin	Precipitin	Lysin	Precipitin
2	384	0	384	0	384	0	768	0
4	3072	0	3072	0	192	0	768	0
6	12288	200	6144	10	192	0	6144	50
8	12288	800	6144	400	768	50	6144	800
9	6144	200	6144	800	1536	100	6144	1600
11	6144	200	12288	200	3072	25	12288	3200
12	1536	100	768	200	384	25	12288	6400
13	1536	50	1536	100	192	25	6144	12800
15	3072	0	3072	50	Died		3072	12800
16	3072	0	3072	0			3072	12800
17	3072		3072				3072	12800
20	1536		1536				3072	12800
23	768		768				3072	12800
28	768		768				3072	12800
36	768		384				3072	3200
42	768		384				1536	1600
49	768		384				1536	800
58	48		48				96	200

The figures under "Lysin" give the highest dilution in which distinct lysis of sheep blood was produced.

The figures under "Precipitin" give the highest dilution of water-laked sheep blood with restored salt content in contact with which the rabbit serum formed a definite precipitate.

solution. Inasmuch as the estimations of lysin were made at frequent intervals during the course of an experiment, and not all at one time, the figures can not be regarded otherwise than as only approximately correct with respect to the concentration of the lysin in the rabbit serum at different bleedings because of the undoubted variations in the activity of the guinea-pig serum used as complement and in the resistance of the sheep corpuscles employed in the different sets of tests. The results are believed to be accurate enough, however, to answer the requirements in this case.

The figures in the tables and curves referring to "Precipitin" give the highest dilution of water-laked sheep blood with restored salt content in contact with which—"ring" or "zonal" method—the serum of the rabbits formed a definite precipitate after 1 hour at room temperature. As the same sheep blood was used in all the estimations, the figures may be regarded as fairly correct, especially since they practically always were made under almost identical conditions.

In the first experiment the introduction of benzene was begun 4 days after the injection of sheep blood.

Four rabbits of the same litter, weighing from 1,500 to 1,600 gm., were given each 30 c.c. of sheep blood intraperitoneally. Four days later the injection of benzene was commenced in 3 of the animals, the 4th serving as control.

TABLE 2

THE EFFECT ON LEUKOCYTES, LYSIN, AND PRECIPITIN IN RABBITS, OF 10 TO 14 CONSECUTIVE DAILY INJECTIONS OF 1 C.C. OF BENZENE, THE FIRST ON THE SAME DAY THAT 30 C.C. OF SHEEP BLOOD WERE INJECTED

Days After Injection of Sheep Blood	Rabbit 1			Rabbit 2			Rabbit 3		
	Lenko- cytes	Lysin	Precip- itin	Lenko- cytes	Lysin	Precip- itin	Lenko- cytes	Lysin	Precip itin
1			...			...			...
2	4500		...	5000		...	3000		...
3	5000		...	4500		...	2800		...
4	3200	1536	10	4000	1536	10	2400	768	10
5		6144	50		12288	50		1536	50
6	3200	12288	300	3800	6144	100	2500	1536	50
7		6144	300		300			1536	300
8	3900	6144	300	3500	12288	1200	2200	1536	300
9		6144	300		6144	1200			300
10	2800		...	4000	12288		2200	1536	300
11		6144	600		6144	1200		3072	100
12	1800	6144	300		6144	1200	2600	1536	100
13		6144	300			1200		3072	100
14		3072	300		12288	1200		3072	100
15	2900	1536	300	3900	6144	1200	2800	3072	100
16		1536	300			600			...
17			...			600			...
18	2800	1536	100	5000	6144	600	3000	3072	100
19			...		6144	600			...
20	4800	1536	100	5400	6144	600	3200	1536	100
22	4200	1536	100	5600	6144	600	3900	1536	50
25	4000	768	100	5300	6144	300	8000	1536	50
30	4600	768	100	6000	6144	200	6400	1536	50
35	4200	768	100	5600	6144	100	4600	1536	50
40	4600	768	100	6500	3072	100	2300	3072	50
47	4425	768	100	5800	1536	100	3800	1536	50
54		768	100		3072	200		768	50
61		48	50		1536	1600		384	50
68			...		768	100		768	50
75			...		768	100		768	50
82			...			0		384	0
92			...			...			...

Rabbits 1, 2, and 3 received 14, and Rabbits 4 and 5, 10 injections of benzene.

The figures under "Lysin" give the highest dilution in which distinct lysis of sheep blood was produced.

Daily injections of 1 c.c. in 1 c.c. of olive oil were given subcutaneously for 8 days. The animals were bled every 2nd day or so until the 17th day, and thereafter at longer intervals, the amount of blood withdrawn each time being from 2 to 3 c.c. In the benzene rabbits the leukocytes rapidly fell to 2,000 or less, the number rising after a few days to fall again before coming to rest at about the normal level.

The results as to lysin and precipitin are given in Table 1. The chief effect of the benzene is seen to be reduction of the precipitin titer. In Rabbit 3, which died 15 days after the injection of sheep

blood, only a small amount of lysin was produced. The results obtained with the serum of the control in this experiment correspond fairly well with the titer of normal rabbits in general injected with 30 c.c. of sheep blood, the lysin titer being, however, rather lower than usual (see Control, Table 2).

TABLE 2—Continued

THE EFFECT ON LEUKOCYTES, LYSIN, AND PRECIPITIN IN RABBITS, OF 10 TO 14 CONSECUTIVE DAILY INJECTIONS OF 1 C.C. OF BENZENE, THE FIRST ON THE SAME DAY THAT 30 C.C. OF SHEEP BLOOD WERE INJECTED

Rabbit 4			Rabbit 5			Control	
Leuko- cytes	Lysin	Precip- itin	Leuko- cytes	Lysin	Precip- itin	Lysin	Precip- itin
6900	96	0	6100	96	0		
10200	24	0	18400	192	0	96	
8900	24	0	11000	192	0	768	
7000	884	0	6800	192	0	12288	
2200	6144	50	3000	1536	0	12288	400
2400	24576	50	4200	1536	25		
2400	24576	50	3100	768	50	24576	1600
3000	24576	200	4300	1536	50	49152	3200
4300	12288	800	5300	1536	50	49152	6400
3600	12288	3200	8900	1536	50		
6500	12288	3200	6100	1536	50	49152	12800
5100	12288	3200	6600	1536	0		
5100	12288	3200	6700	768	0	49152	6400
3400	6144	6400	3900	768	0		
2600	3072	3200		768	0	24576	6400
.....	3072	3200		768	0		
.....					0	24576	1600
.....	3072	3200		768	0		
.....	Died						
.....				192	0	6144	3200
.....						3072	800
.....						1536	200
.....						768	200
.....						768	200
.....						768	200
.....						768	200
.....						768	0

The figures under "Precipitin" give the highest dilution of water-laked sheep blood with restored salt content in contact with which the rabbit serum formed a definite precipitate.

In the next experiment the administration of benzene was commenced on the same day that the sheep blood was injected and then continued daily in some animals for 10 days, in others for 14 (Table 2). Practically the same results were obtained as in the first experiment; namely, marked reduction in the precipitin titer and in some rabbits reduction in lysin also. In Rabbit 4, which received 1 c.c. of benzene per kilo daily for 14 days, considerable reduction of leukocytes was observed, but in this animal the amounts of lysin and precipitin were not

much less than in many nonbenzenized rabbits. In Rabbit 5, however, which was treated in the same way, the leukocytes seem to have been reduced less and the lysin and precipitin more, a divergence of results that one might regard as due to a natural variation in the power to produce antibodies.

The following experiment illustrates the result when benzene is injected daily for some days both before and after the injection of sheep blood. The figures of Table 3 appear to show that in rabbits

TABLE 3

THE EFFECT ON THE PRODUCTION OF ANTIBODIES OF INJECTIONS OF BENZENE (1 C.C. PER KILO) FOR SOME TIME BEFORE AND AFTER THE INJECTION OF THE ANTIGEN

Days After Injection of Sheep Blood	Rabbit 1			Rabbit 2			Rabbit 3			Rabbit 4		
	Leu- ko- cytes	Lysin	Pre- cip- itin	Leu- ko- cytes	Lysin	Pre- cip- itin	Leu- ko- cytes	Lysin	Pre- cip- itin	Leu- ko- cytes	Lysin	Pre- cip- itin
	5000			4600			5000			6100		
	4500			2700			5000			4500		
	3500			5300			6200			7200		
	3700			2300			5400			6800		
	4500			7500			6800			6600		
	6300			5300			6300			5100		
	1400			2000			6000			5600		
1	3000			4200			8000			9600		
2	1800			5400			4800			4600		
3	5600			7400			10200			9400		
4	3400	384	100	3400	192	0	8000	1536	0	6800	1536	0
5		384	0		192	0		3072	0		3072	0
6	5200	1536	50	4400	768	0	5400	6144	100	4400	6144	0
7	4600			4500			10000			2600		
8	3000	1536	50	3200	1536	0	5000	6144	600	2000	3072	0
9		768	100		768	0		6144	1600		6144	1600
10	1000			5000			4400			2600		
11	3400	1536	100	6000	768	0	8400	6144	1600	2400	6144	1200
12	1600			7200							Died	
14		1536	0		384	0		3072	1600			
16	9000*	768	0	3000	384	0	2600	3072	800			
18	8400*	768	0	4800	384	0	5000	1536	400			
21	1800	768	0	3000	384	0	4400	1536	400			

* Suppuration about gangrenous area in skin.

The figures under "Lysin" give the highest dilution in which distinct lysis of sheep blood was produced.

The figures under "Precipitin" give the highest dilution of water-laked sheep blood with restored salt content in contact with which the rabbit serum formed a definite precipitate.

so treated the amount of antibody set free may be very small, even tho the reduction of leukocytes is not extreme. The same general result is illustrated also by the record of 1 member of a group of 6, the other members of the group dying too early to permit extended observations, the results so far as they went, however, having the same bearing.

The rabbit weighed 1,030 gm. It was given 1 c.c. of benzene for 8 consecutive days; the leukocytes not falling, it was given 2 c.c. on each of the next 2 days, and for 3 days 3 c.c. each day. Two days later it received intraperitoneally 30 c.c. of sheep blood. The leukocytes remained at about 6,000

until about 5 days after the last injection of benzene; then their number fell to 3,500; subsequently the number remained between 4,000 and 5,000 for about 25 days and then rose to 6,000 or more. The amount of precipitin formed after the injection of sheep blood was insignificant; only on two days, the 6th and the 15th, did the titer reach 100. The amount of lysin produced was also small, the highest titer being 768 on the 20th day. On the 56th day after the first injection of sheep blood 30 c.c. of sheep blood were injected again and large amounts of lysin and precipitin were produced with more promptness than in the fresh animal.

TABLE 4
THE EFFECT OF BENZENE WHEN INJECTED AT THE HEIGHT OF ANTIBODY PRODUCTION

Days After Injection of 30 c.c. Sheep Blood	Injection of Benzene, c.c.	Rabbit 1		Rabbit 2		Rabbit 3	
		Lysin	Precip- itin	Lysin	Precip- itin	Lysin	Precip- itin
1		24		0		0	
2		24		0		0	
3		384		384		768	
4		3072		6144		6144	
5		12288	100	49152	50	12288	100
6		24576	400	24576	400	24576	400
7		12288	600	24576	600	24576	400
8		24576	600	24576	600	24576	1200
10		12288	2400	24576	600	24576	2400
11		12288	2400	12288	2400	24576	4800
12	0.3	6144	2400	24576	2400	12288	4800
13	0.3	12288	2400	24576	2400	12288	4800
15	0.5	3072	1600	3072	2400	1536	4800
17	0.5	768	1600	3072	3200	1536	6400
18	0.5	384	1600	3072	3200	768	6400
20	0.5	384	1600	3072	6400	768	3200
22	0.75	3072	2400	12288	9600	3072	3200
25	0.75	1536	1200	12288	9600	3072	3200
28	0.75	1536	1200	6144	4800	3072	3200
32	0.75	1536	1200	6144	2400	3072	9600
35			1200	6144	9600	3072	36000
42		768	1200	3072	19200	Died	
50		768	1200	6144	12800		
59		768	1200	768	3200		
70		768	1200	768	4800		
77		768	600	768	2400		
84		768	300	768	600?		
94		768	300	768	640		
108		Died		768	200		
113				384	100		
129				384	800		
145				768	200		
161				768	200		
174				192	100?		
186				384	0		
208				384	0		
216				384	0		

The figures under "Lysin" give the highest dilution in which distinct lysis of sheep blood was produced.

The figures under "Precipitin" give the highest dilution of water-laked sheep blood with restored salt content in contact with which the rabbit serum formed a definite precipitate.

The effect of benzene on the production of antibodies when production was at or near the height of the curve, was tested also. Rabbits were injected intraperitoneally with 30 c.c. of sheep blood and on the 12th day or so injections of benzene were commenced. In the first

experiment of this kind all the animals died soon after the injections of 1 c.c. per kilo of benzene were started, and on this account in the next experiment (Table 4) the initial quantity injected was reduced to 0.3 c.c. per kilo and then gradually increased as shown in the table. There was no definite decrease in the leukocytes after the injections of benzene. The results indicate some temporary reduction of lysin in the blood but no definite effect on the precipitin. In all the animals the production of lysin and precipitin persisted longer and the precipitin underwent much more fluctuation than is the case under ordinary circumstances. Just before it died, Rabbit 3 of this experiment gave a marked sharp rise in precipitin content.

TABLE 5
INJECTION OF BENZENE AT THE HEIGHT OF ANTIBODY PRODUCTION

Days After 2nd Injection of 30 c.c. Sheep Blood	Injections of Benzene	Lysin	Precipitin	Remarks
3		12288	200	No reduction of leukocytes
5		49152	800	
7	1.5 c.c. per kilo	49152	6400	
8	1.5 c.c. per kilo	49152	25600	
9	1.5 c.c. per kilo	12288	19200	
10	1.5 c.c. per kilo			
11	2 c.c. per kilo	12288	12800	
12	2 c.c. per kilo	24576	19600	
13	2 c.c. per kilo	12288	25600	
14			25600	
15		12288	25600	
17		12288	25600	
19		12288	25600	

In several other experiments of the same kind benzene has had a similar effect, even when given in much larger doses. Table 5 gives the results in a rabbit which had received 1.5 c.c., eventually 2 c.c., of benzene per kilo each day for several days during the height of the production of antibodies after a second injection of 30 c.c. of sheep blood. Reference may be made also to the injection of benzene without any recognizable effect in rabbits treated with toluene during the early stages of the production of antibodies for sheep blood. In 2 rabbits, which received 12 consecutive daily injections of toluene, approximately 1 c.c. per kilo, the 1st on the same day that 30 c.c. of sheep blood were injected, the injection some 2½ months later of 13 daily doses of 1 to 1.5 c.c. of benzene per kilo had no marked effect on the number of leukocytes, the course of antibodies, or the general health of the animals so far as ascertainable. The details of this experiment will be given later.

When doses smaller than 1 c.c. per kilo of benzene were given to rabbits injected with 30 c.c. of sheep blood, the reduction in antibodies was less pronounced, and 0.005, 0.01, and 0.025 c.c. per kilo, given daily for several days both before and after the injection of the antigen, appeared to have but little appreciable effect. There were no indications that the smaller doses just mentioned—0.01 c.c. and 0.005 c.c.—cause any stimulation of the production of antibodies so as to lead to a greater concentration in the serum than in the control rabbits even tho such doses may induce a moderate degree of leukocytosis. In dogs, however, relatively small doses of benzene appeared to increase the production of antibodies (Table 6). It would be of interest to test more thoroughly the effect of small doses of benzene when given as indicated and also during the height of the antibody curve.

TABLE 6
THE EFFECT OF BENZENE IN SMALL DOSES ON THE BLOOD AND ON THE FORMATION OF LYSIN
IN THE DOG

Days After Injection of 1 c.c. of a 10% Suspension of Goat Blood per Kilo of Weight of Dog	Injections of 0.2 c.c. of Benzene per Kilo of Weight of Dog	Lysin (The figures give highest active dilution of dog serum)	Red Corpuscles	Leukocytes
	+	24		
	+	24		
1	+	24	7896000	30300
2	+	48	7598000	34450
3	+	768	8334000	29000
4		6144		
5		6144		
6	+	24576	7560000	20350
7	+	98304	6240000	27000
8	+	98304	7132000	30950
9		49152	6034000	35850
10		49152	6040000	31625
11		49152	6589000	29400
13		49152		
18		24576		
23		24576		
30		12288		

In making the tests the dog serum was heated to 58 C. for 30 minutes; each mixture measured 0.6 c.c., 0.2 c.c. being a 5% suspension of goat blood; 0.0125 c.c., guinea-pig serum as complement; the rest, dog serum and salt solution.

Besides this study on rabbits, I have made a few observations on the action of benzene on the dog and on the white rat.

In dogs the subcutaneous injection of 1 c.c. of benzene for each kilo of weight may cause at first leukocytosis, sometimes as high as 60,000 to 70,000, reduction of leukocytes taking place only after many injections and the development of fatty changes and hemorrhages. On account of the severe local and general effects no extended experiments

have been made. Table 6 gives the results of an experiment in which 0.2 c.c. of benzene per kilo of the body weight was injected several times; not only did leukocytosis develop, but the number of red corpuscles remained high and the production of lysin for goat corpuscles reached a very high point. Usually the highest point reached by the lysin after the injection of 1 c.c. of a 10% suspension of goat blood per kilo of the weight of the dog in otherwise normal animals is from 12,288 to 24,576.

In the white rat doses larger than 1 c.c. for each kilo of weight were required to depress the number of leukocytes in the blood. The most pronounced results were obtained after from 4 to 5 daily doses of from 0.6 to 0.8 c.c. of benzene with olive oil to make 2 c.c., injected subcutaneously in rats weighing from 130 to 160 gm. Usually the leukocytes began to fall after the 2nd or 3rd injection, and in a day or two the number might run less than 2,000; a gradual return to normal set in about 3 or 4 days after the injection of benzene was stopped. The injection of sheep blood (10% suspension) in the dose of 5 c.c. per kilo of weight at the end of such benzene treatment was followed by a much smaller production of specific lysin than followed in the case of rats injected with sheep blood but not with benzene. It was noted, however, that the lysin titer of the serum of benzenized rats might continue above the normal a little longer than in the untreated rats. The death rate was high in experiments on rats with benzene, even when much smaller doses than those mentioned were given.

In the course of this work many of the observations by Selling and others on the effect of benzene, in doses of 1 c.c. per kilo of weight of rabbit, on the leukocytes in the peripheral blood and on the general condition have been verified. In most cases a few injections given daily caused a distinct leukopenia, affecting the granular cells much more than the lymphocytes, and frequently also some diminution of the number of red corpuscles. In animals with extreme leukopenia the death rate was high, and in the rabbits that died while under the influence of benzene the bone marrow was poor in cells. After the first injections of benzene there sometimes occurred a preliminary leukocytosis of moderate degree. A rise in the number of leukocytes followed by a second fall before return to the normal level, as emphasized by Weiskotten, Schwartz and Steensland,¹² was observed in a number of animals.

¹² Jour. Med. Research, 1915, 33, p. 127.

In animals with distinct benzene leukopenia there was nearly always a marked reduction in the amount of lysin and precipitin, but the general form of the curves described by these substances in passing into and out of the blood resembled that of the curves in nonbenzenized rabbits, the lysin reaching the high point earlier than the precipitin, but the last phase of the curve—the gradual return to the normal—sometimes persisted longer in the benzenized animals (Charts 1 and 2).

Exceptionally rabbits were encountered in which many consecutive daily injections of 1 c.c. of benzene per kilo appeared to have little or no effect on the number of leukocytes in the peripheral blood. In these animals the relative number of granular leukocytes might be diminished while the relative number of lymphocytes was increased; in some of the resistant rabbits the production of antibodies did not seem affected nearly so much as in leukopenic rabbits.

At the height of antibody-production benzene may have little, if any, effect (Tables 4 and 5) either on antibody concentration in the serum or on the leukocytes. It appears as if rabbits at this time may have an increased resistance to benzene, and it is noteworthy too that in rabbits injected with benzene at this time, the antibodies persist longer and may undergo greater fluctuation in concentration than in rabbits not so treated.

Hamburger¹³ finds that in the test tube small amounts of benzene promote phagocytosis, but in rabbits with benzene leukopenia, no such effect is demonstrable. As benzene leukopenia is established, the leukocytes in the blood may appear more or less injured (as observed by Smith¹⁴ in the blood of leukemic patients treated with benzene) and their phagocytic activity becomes reduced, the cytophagic index for staphylococcus under the opsonic influence of normal rabbit serum having been found as low as 0.5 to 0.3 with leukocyte counts of about 1,800. Hence we may conclude that benzene may lower the anti-infectious powers of the body in at least 3 ways: by reduction of antibodies, by reduction of the number of leukocytes, and by reduction of the phagocytic activity of leukocytes. That the unguarded use of benzene in the treatment of leukemia may be associated with danger on account of its lowering the resistance to infection, is hardly to be doubted, especially in view of the course of events in some cases so treated.

The obvious explanation of the depression of antibody-formation under benzene is damage to the marrow and the lymphatic structures,

¹³ *Lancet*, 1916, 1, p. 37.

¹⁴ *Jour. Am. Med. Assn.*, 1915, 64, p. 1734.

Chart 1. Precipitin Curves: Curve 1 = Rabbit 2, Table 1. Curve 2 = Rabbit 3, Table 2. Curve 3 = Rabbit 3, Table 3. Curve 4 = normal control rabbit.

which we have strong reasons to believe are concerned directly with this formation.

Pfeiffer and Marx¹⁵ and others¹⁶ have shown that various antibodies can be demonstrated in the bone marrow, spleen, and lymph glands before they can be demonstrated in the blood; under certain circumstances the removal of the spleen diminishes the amount of antibody produced;¹⁷ when the blood-making organs are stimulated, as in acute anemia, the production of antibodies is increased; when the blood-making tissues are destroyed, as by Roentgen ray, the production is diminished;¹⁸ and finally, it has been shown recently by Carrell and Ingebrigtsen¹⁹ and others²⁰ that in the presence of antigenic substances, cultures of the spleen and bone marrow outside of the body may manufacture antibodies (lysin, agglutinin, precipitin).²¹

Chart 2. Lysin and Precipitin Curves: Fine line = lysin. Heavy line = precipitin.

Just how benzene interferes with the elaboration of antibodies, whether by direct injury to cells, which seems likely in view of its destructive effects on the tissues concerned, or possibly by modification of enzyme action without necessary injury to cells, are questions that at present seem difficult of solution. That benzene, in spite of its affinity for the nervous tissues according to Joachimoglu,²² acts on the

¹⁵ Ztschr. f. Hyg. u. Infektionskrankh., 1898, 27, p. 272.

¹⁶ Cantacuzene: Ann. de l'Inst. Pasteur, 1902, 16, p. 552; 1908, 22, p. 54. Tsurumi and Kohda: Ztschr. f. Immunitätsf., 1913, 19, p. 519.

¹⁷ Deutsch: Ann. de l'Inst. Pasteur, 1899, 13, p. 689.

¹⁸ Hektoen: Jour. Infect. Dis., 1915, 17, p. 415. Simonds and Jones: Jour. Med. Research, 1915, 33, p. 183.

¹⁹ Jour. Exper. Med., 1912, 15, p. 287.

²⁰ Lüdke: Berl. klin. Wchnschr., 1912, 49, p. 1034. Przygode: Wien. klin. Wchnschr., 1913, 26, p. 841; 1914, 27, p. 201.

²¹ For more complete discussion of the earlier literature see Hektoen, Harvey Society Lectures, 1909-10, p. 177, and Gay and Rusk, Tr. 15th Internat. Cong. on Demography and Hygiene, 1913, 2, p. 378.

²² Biochem. Ztschr., 1915, 70, p. 93.

elements concerned in the elaboration of antibodies and that the leukocyto-genic centers are concerned in this elaboration is indicated not only by the reduction of antibodies in benzene leukopenia but also, it seems to me, by the resistance to these effects of benzene at the time when the production of antibodies is at or near its height, and by the leukocytosis and increased lysin-formation observed in the dog under the influence of small doses of benzene. As pointed out by Simonds and Jones, the morphologic effects of benzene do not indicate that certain particular elements are concerned in the elaboration of antibodies to the exclusion of others, because the action is too general in that, while benzene affects the leukocyto-genic centers and the leukocytes the most, it also acts, directly or indirectly, on the lymphocytes and the erythroblastic elements. The x-ray, which also reduces the production of antibodies, does not permit any morphologic differentiation because it too is diffuse in its action, altho in a different order of intensity. Possibly the study of the action of substances closely related to benzene may yield results of interest with respect to these questions. The work of Milnikowa and Wersilowa²³ on the action of phenylhydrazin(and of hydroxylamin hydrochlorid) on the development of agglutinins for typhoid bacilli is limited to the demonstration that the effect in the doses used is one of restraint.

The results of the investigations into antibody-formation just cited indicate that the final elaboration of antibodies takes place outside the circulation, and the results of transfusion experiments by Carlson and myself point in the same direction.²⁴ Assuming that such is the case, we have to consider the possibility that in benzenized rabbits the absorption and fixation of the antigen may be disturbed and delayed. Von Heinrich²⁵ explains the depressive action of the x-ray by assuming that it prevents the lymphatic apparatus from binding the antigen. We know that after the injection of considerable amounts, antigen may persist in the blood for some time, apparently even after the appearance of specific antibodies;²⁶ hence the question arises whether the course of the antigen in the blood differs in the case of benzenized animals from that in normal animals. I have made some observations on this point which show that in rabbits injected intraperitoneally with 30 c.c. of sheep blood the blood serum may contain sheep protein in such form as to

²³ *Centralbl. f. Bakteriol.*, I, O., 1912, 66, p. 520.

²⁴ *Jour. Infect. Dis.*, 1910, 7, p. 319.

²⁵ *Centralbl. f. Bakteriol.*, I, O., 1913, 70, p. 421.

²⁶ References bearing on this fact are cited by Zinsser, *Arch. Int. Med.*, 1915, 16, p. 223. See also Gay and Rusk, *Univ. Calif., Publ. in Pathol.*, 1912, 2, p. 59.

be demonstrable by means of the precipitin test for from 10 to 12 days and sometimes longer, in one instance even on the 20th day, the test in this case being made with an antiserum of a titer of 1:20000—that is, it would cause a precipitate in a dilution of sheep blood of 1:20000. I have not met any indications, however, that under the circumstances outlined, sheep protein is demonstrable by the precipitin method any longer in the blood of benzenized rabbits than in the blood of rabbits not injected with benzene, at least when antiserum of the titer of 1:12000, or thereabout, is used.

On account of its bearing on the question of what elements in the marrow and lymphatic tissues are concerned in the production of antibodies, the result of the following experiment may be of interest at this point:

In conjunction with Professor A. S. Loevenhart of the University of Wisconsin, I determined the production of lysin for sheep corpuscles in white rats which were kept in an atmosphere of 10% oxygen by means of the special apparatus devised by Professor Loevenhart to maintain atmospheres with controllable oxygen content. The rats under low oxygen pressure during the entire period of antibody-production developed an increase of hemoglobin and red cells of from 20 to 35%—total hemoglobin being increased 43%—as compared with that of the controls, but no demonstrable increase or decrease in the amount of lysin, as compared with that of controls. Hence, increase in red corpuscles under these conditions does not seem to be associated with changes in the production of antibodies as might be expected in view of the stimulating effect that acute anemia is known to have on this production; but of course without corroborative evidence this result alone, altho suggestive, does not permit one to exclude the erythrocytogenic centers from consideration as possible producers of antibodies.

The ultimate mechanism of the action of benzene, not only in restraining the production of antibodies, but in causing injury of a selective nature to marrow and lymphatic cells and leukocytes, invites further study.

SUMMARY

In rabbits the repeated injection of benzene in doses of 1 c.c. per kilo of rabbit weight at about the same time that sheep blood is injected greatly reduces the production of specific precipitin and lysin.

In considerably larger doses benzene has a like effect on the production of lysin in white rats.

The reduction of antibody-formation under these circumstances is associated with grave lesions in the marrow, with leukopenia, and other changes characteristic of benzene intoxication, the leukocytes in the rabbit being of reduced phagocytic power.

In the dog, benzene (0.02 c.c. per kilo) may cause leukocytosis associated with increase in production of lysin for goat corpuscles.

So far as can be determined by the precipitin method, the course of the antigen in the blood appears the same in benzenized as in non-benzenized rabbits.

At the height of antibody-production the injection of benzene appears to have but little effect on the leukocytes of the blood and on its antibody content, the precipitin especially persisting longer and with more fluctuation than otherwise.

Benzene may lower the resistance to infection by reduction (1) of antibody-production, (2) of the number of leukocytes, and (3) of leukocytic activity.

That benzene acts on elements that elaborate antibodies and that the leukocytogenic centers are concerned in this elaboration is indicated (a) in the rabbit, by the reduction of antibodies and of leukocytes and by the resistance to these effects when antibody-production is at or near its highest activity as measured by the concentration of antibodies in the blood, and (b) in the dog, when suitable doses are given, by leukocytosis and increased formation of lysin.